

SOQUEL DEMONSTRATION STATE FOREST

GENERAL FOREST MANAGEMENT PLAN



CALIFORNIA DEPARTMENT OF FORESTRY
AND FIRE PROTECTION

SOQUEL, CALIFORNIA
November 5, 2014



**SOQUEL DEMONSTRATION
STATE FOREST

GENERAL FOREST

MANAGEMENT PLAN**

This final General Forest Management Plan is intended to set the goals and to guide the development of programs and facilities on Soquel Demonstration State Forest (SDSF). It was approved by the SDSF Advisory Committee on January 14, 2014 and by the Board of Forestry on November 5, 2014.

**CALIFORNIA DEPARTMENT OF FORESTRY
AND FIRE PROTECTION**

**SOQUEL, CALIFORNIA
March 4, 1998
Revision date: November 5, 2014**

Angela Bernheisel, Forest Manager

SDSF ADVISORY COMMITTEE:

California Board of Forestry and Fire Protection – Matt Dias
California Department of Fish and Wildlife – Terris Kasteen
California Department of Parks and Recreation – Tim Hyland and Chris Spohrer
County of Santa Cruz – Matt Johnston
The Nature Conservancy – Larry Serpa
Neighborhood Representative – Patricia Marland (Secretary)
Registered Professional Forester – Geoffrey Holmes
Resource Conservation District of Santa Cruz County – Steve Butler (Vice-Chair)
Stewards of Soquel Forest – Patty Ciesla (Chair)

THE ORIGINAL PLAN WAS PREPARED BY:

Thomas Sutfin, Forest Manager
Richard Eliot, Assistant Forest Manager
Sheryl Nives, Forestry Aide

AUTHORS OF THE 2012 UPDATES:

CAL FIRE:

Angela Bernheisel, Forest Manager
Jill Butler, Forester
Pete Cafferata, Forest Hydrologist
Anna Cameron, Office Technician
Julie Dufrense, Forestry Aide
Lawrence Erickson, Fire Captain
Jeff Leddy, State Forests Biometrician
Elizabeth Lucas, Forestry Aide
Jack Marshall, Forest Pathologist (retired)
Bob Motroni, Senior Wildlife Biologist
Thomas Sutfin, Forest Manager (retired)
Chuck Whatford, Associate State Archaeologist

OTHER:

Jim Robins, *Alnus* Ecological
Patty Ciesla, Stewards of Soquel Forest

**KEN PIMLOTT
DIRECTOR
CALIFORNIA DEPARTMENT OF FORESTRY
AND FIRE PROTECTION**

**J. KEITH GILLESS
CHAIR
BOARD OF FORESTRY
AND FIRE PROTECTION**

**EDMUND G. BROWN, JR.
GOVERNOR
STATE OF CALIFORNIA**

**JOHN LAIRD
SECRETARY FOR RESOURCES
THE NATURAL RESOURCES AGENCY**

TABLE OF CONTENTS

CHAPTER 1: INTRODUCTION..... 3
CALIFORNIA'S DEMONSTRATION STATE FOREST SYSTEM..... 3
THE GENERAL FOREST MANAGEMENT PLAN..... 3
FUTURE PLANNING..... 4
CHAPTER 2: MANAGEMENT GOALS 5
RESOURCE PROTECTION AND ENHANCEMENT 5
DEMONSTRATION AND EDUCATION 6
RESEARCH..... 6
TIMBER MANAGEMENT..... 7
RECREATION 7
CHAPTER 3: PROPERTY DESCRIPTION..... 9
LOCATION AND BOUNDARIES 9
ADJACENT OWNERSHIP..... 9
HISTORY OF OWNERSHIP 9
CLIMATE..... 12
SOILS AND GEOLOGY 12
WATER RESOURCES 14
ZONING AND GENERAL PLAN 15
CHARACTERISTIC FEATURES 16
CHAPTER 4: ADMINISTRATION 17
THE LEASE..... 17
CAL FIRE ADMINISTRATION..... 17
FUNDING AND TAXES 18
ADVISORY COMMITTEE 19
SAFETY 20
LAW ENFORCEMENT 21
ACCESS NEEDS..... 21
LAND ACQUISITION PRIORITIES..... 22
COOPERATION WITH ADJACENT LANDOWNERS..... 23
COOPERATION WITH OTHER AGENCIES AND INSTITUTIONS..... 23
CHAPTER 5: BIOTA..... 25
INTRODUCTION 25
HABITAT COMMUNITIES 25
FUNGAL RESOURCES 30
FERAL PIGS 30
BULLFROGS 31
CORVIDS 31
INVASIVE SPECIES 32
PLANT SPECIES OF CONCERN 33
SPECIAL-STATUS WILDLIFE SPECIES..... 34
MANAGEMENT GUIDELINES 37
PLANNED ACTIONS..... 38

CHAPTER 6: FISHERIES	39
INTRODUCTION	39
HABITAT AND RESOURCE CONSIDERATIONS	47
AQUATIC INVERTEBRATES	50
MANAGEMENT GUIDELINES	53
PLANNED ACTIONS.....	54
CHAPTER 7: WATERSHED ASSESSMENT	56
PHYSIOGRAPHY OF THE WATERSHED.....	56
BENEFICIAL USES.....	56
LAND USE HISTORY	60
CUMULATIVE WATERSHED EFFECTS ASSESSMENT	60
MONITORING AND ENHANCEMENT	65
MANAGEMENT GUIDELINES	66
PLANNED ACTIONS.....	66
CHAPTER 8: DEMONSTRATION AND EDUCATION	67
INTRODUCTION	67
DEMONSTRATION PROGRAM.....	67
FORESTRY EDUCATION	69
INTERPRETATION.....	70
VOLUNTEER PROGRAM.....	71
FORESTRY EDUCATION CENTER.....	72
CHAPTER 9: TIMBER MANAGEMENT	74
INTRODUCTION	74
HARVEST HISTORY	74
CONIFER VOLUME INVENTORY	75
GROWTH.....	77
VEGETATION TYPE	78
TIMBER SITE CLASS	78
SILVICULTURE.....	78
OLD-GROWTH AND LATE-SUCCESSION MANAGEMENT	82
HARVEST SYSTEMS	86
HARVEST PLANNING.....	87
TIMBER OPERATIONS MONITORING	92
TIMBER MARKETS.....	92
HARDWOOD MANAGEMENT	93
DEMONSTRATION AND RESEARCH PROGRAMS	94
MINOR FOREST PRODUCTS.....	95
MANAGEMENT GUIDELINES	96
PLANNED ACTIONS.....	98
CHAPTER 10: RESEARCH	99
INTRODUCTION	99
COMPLETED PROJECTS.....	99
RESEARCH PRIORITIES	104

MANAGEMENT GUIDELINES	106
PLANNED ACTIONS.....	106
CHAPTER 11: RECREATION.....	107
INTRODUCTION	107
RULES, REGULATIONS AND POLICIES	107
RECREATION STUDY AND PLAN	108
PARTNERSHIPS	109
REGIONAL CONTEXT.....	110
PUBLIC ACCESS AND STAGING AREAS.....	112
EXISTING FACILITIES	112
RECREATION PROGRAMS	118
MANAGEMENT GUIDELINES	119
PLANNED ACTIONS.....	120
CHAPTER 12: RESOURCE PROTECTION	121
FIRE PROTECTION	121
FOREST PEST MANAGEMENT	129
CHAPTER 13: ARCHAEOLOGY	134
HISTORY OF SDSF AREA	134
ARCHAEOLOGICAL RESEARCH.....	134
PREHISTORIC SITES	135
HISTORIC SITES.....	135
MANAGEMENT GUIDELINES	136
PLANNED ACTIONS.....	136
CHAPTER 14: ROADS AND OTHER IMPROVEMENTS.....	137
SDSF FACILITIES.....	137
PUBLIC ACCESS ROADS	138
ROAD BUILDING AND MAINTENANCE	139
MANAGEMENT GUIDELINES	140
PLANNED ACTIONS.....	141
REFERENCES.....	143
LITERATURE.....	143
PERSONAL COMMUNICATION	154
APPENDICES	155
APPENDIX A: TEXT OF ASSEMBLY BILL 1965.....	155
APPENDIX B: SPECIES LISTS FOR SDSF.....	157
APPENDIX C: SOQUEL DEMONSTRATION STATE FOREST MONITORING PLAN...	172
APPENDIX D: BEST MANAGEMENT PRACTICES	178

LIST OF FIGURES

Figure 1.	Locator Map for SDSF.....	10
Figure 2.	SDSF and Adjacent Ownerships.....	11
Figure 3.	Locations of Soil Series in SDSF.....	13
Figure 4.	Recent Acquisition Map for SDSF	23
Figure 5.	Vegetation Type Map for SDSF	27
Figure 6.	Estimated Juvenile Steelhead Population Comparison Index Reaches, Soquel Demonstration State Forest, 1993-2012	46
Figure 7.	Water Temperature Data for SDSF.....	51
Figure 8.	Soquel Creek Watershed Map	57
Figure 9.	Annual Peak Streamflow for Soquel Creek	58
Figure 10.	Rainfall and Streamflow for East Branch of Soquel Creek	59
Figure 11.	SDSF Timber Harvesting Plans	76
Figure 12.	Timber Site Class Map for SDSF	79
Figure 13.	Conifer Diameter Class Distribution for SDSF	80
Figure 14.	Old-growth Stands and Late-Succession Management Areas	83
Figure 15.	SDSF Projected Inventory by Planning Period.....	90
Figure 16.	SDSF Projected Annual Growth and Harvest.....	90
Figure 17.	Santa Cruz County Redwood Stumpage Values.....	93
Figure 18.	Estimated Basal Area for SDSF's Hardwood Tree Species	94
Figure 19.	Transportation System of SDSF	114
Figure 20.	Recent Fire History	123
Figure 21.	Fire Defense Improvements	124

LIST OF TABLES

Table 1. Soil Type and Associated Erosion Hazard of SDSF.....	14
Table 2. Key to California Wildlife Habitat Relationship Vegetation Type Codes.	26
Table 3. Invasive Exotic Plant Species Occuring at SDSF.....	33
Table 4. Special-status Vertebrate Fish and Wildlife Species Occuring or with Potential to Occur at Soquel Demonstration State Forest	36
Table 5. Habitat Requirements and Vulnerability by Each Salmon Life Stage.....	44
Table 6. SDSF 2006 Timber Inventory Summary	77
Table 7. Individual Redwood and Douglas-fir Tree Characteristics	84
Table 8. SDSF Forest Land Availability.....	89
Table 9. Projected SDSF Forest-wide (full and limited availability) Average Annual Inventory, Harvest and Growth by Planning Period	89
Table 10. Forest Pests of the Soquel Creek Watershed	130
Table 11. Drivable Road Segments of SDSF.....	139

EXECUTIVE SUMMARY

BACKGROUND INFORMATION ON SOQUEL DEMONSTRATION STATE FOREST

Total acres:	2,681
Vegetation types present:	Redwood, Montane Hardwood-Conifer, Coastal Oak Woodland, Douglas-fir, Mixed Chaparral, Annual Grassland, Riparian Community, and Freshwater Marshes
Miles of roads:	19
Miles of trails:	12.2
Primary management goals:	Watershed protection, monitoring, and study; public education about forestry, timber production, and the environment; and protection of old-growth redwoods.

The Soquel Demonstration State Forest (SDSF) was established in 1990 according to Assembly Bill 1965 of 1987 (now Public Resources Code (PRC) Sections 4660-4664). The basic purpose of the enabling legislation is to establish and preserve SDSF (also referred to as the Forest) as an intensively managed, multifaceted research forest. The Public Resources Code Sections 4660-4664 state that SDSF will do all of the following:

- Provide watershed protection for local communities and base-line monitoring and studies of the hazards, risks, and benefits of forest operations and watersheds to urban areas.
- Provide public education and examples illustrating compatible rural land uses, including sustained yield timber production, as well as the historic development of timbering and forestry machinery, within the context of local community protection and nearby pressures.
- Provide a resource for the public, environmental groups, elected officials, environmental planners, the educational community, and the media as an open environment for the inspection and study of environmental education, forestry practices, and effects thereof.
- Protect old-growth redwood trees.

Soquel Demonstration State Forest is located near the center of Santa Cruz County, California, approximately eight miles northeast of the city of Santa Cruz. Positioned in the southern portion of the Santa Cruz Mountains, SDSF is eighteen air miles south of San Jose and within a two-hour drive of the San Francisco and Oakland metropolitan areas.

SDSF is bordered by both state and privately-owned properties. These include The Forest of Nisene Marks State Park, timberland managed by Redwood Empire, the Olive Springs Quarry, and private rural-residential parcels that range in size from 1 to 80 acres.

Public access is currently limited to a road access point off Highland Way or entry through The Forest of Nisene Marks State Park. The inability of the public to drive to the Forest when county roads are closed limits accomplishment of SDSF's mission to provide for forestry education and

demonstration.

Between the late 1920s and early 1940s, the previous owner managed the timber resource on an even-aged basis with clearcutting and natural regeneration. Since 1990, SDSF has been managed on an uneven-aged basis, utilizing either single tree and/or small group selection. Watershed, soil, fisheries, and wildlife resources are monitored and protected before, during, and after all timber harvesting activity. Funds derived from timber harvesting provide for forest staff, facilities, operations and maintenance.

Existing old-growth redwood areas in the Forest have been excluded from timber harvesting, as mandated by SDSF's authorizing legislation. In addition, late-succession areas are managed to promote the development of old-growth habitat. These areas make up 15 percent of the overall forest, and protect 300 feet on each side of the East Branch of Soquel Creek, Amaya and Fern Gulch Creeks. The East Branch of Soquel Creek provides valuable steelhead trout spawning and rearing habitat, and is part of the domestic water supply for the local community.

BACKGROUND INFORMATION ON THE GENERAL FOREST MANAGEMENT PLAN

In 1988, The Nature Conservancy entered into a lease with the State of California to manage the SDSF property. The lease and management responsibilities were transferred to the California Department of Forestry and Fire Protection (CAL FIRE, formerly CDF) in 1990. The first General Forest Management Plan was developed with input from the public, resource management professionals, and members of the SDSF Advisory Committee.

The Board of Forestry and Fire Protection (the Board) approved the original General Forest Management Plan on May 18, 1998. The environmental impacts of the 1998 management plan were investigated in an Environmental Impact Report (EIR), State Clearinghouse Number 94023033. Mitigation measures identified in the EIR have been incorporated into a Monitoring Plan located in Appendix C.

The SDSF General Forest Management Plan remains in effect until it is amended or a new plan is adopted. In 2003 and 2009, the Board reviewed and reauthorized the plan. No revisions were made at those times. This revision of the plan incorporates the results of new studies, monitoring and research pertaining to fisheries, wildlife habitat, watershed issues, archaeology, and timber management.

CAL FIRE plans to work with the Advisory Committee to reexamine the General Forest Management Plan every five years and determine what changes are needed. These will be developed by CAL FIRE and the Advisory Committee in accordance with Public Resources Code Sections 4660 – 4664 and the California Environmental Quality Act.

CHAPTER 1: INTRODUCTION

CALIFORNIA'S DEMONSTRATION STATE FOREST SYSTEM

The demonstration state forest system of the California Department of Forestry and Fire Protection (CAL FIRE) was established in the mid-1940's to meet local needs for research, demonstration, and education related to forest management. Currently, the demonstration state forest (DSF) system encompasses over 71,000 acres of land in the form of eight state forests. DSFs are healthy, living forests which demonstrate conservation and protection of wildlife, fisheries, vegetation, soil, and watershed resources as well as sustained-yield forest management activities.

The Soquel Demonstration State Forest (SDSF), when established in 1990, was the first addition to the DSF system in over 40 years. SDSF contains 2,681 acres including mostly coast redwood and mixed evergreen forest types. Former Assemblyman Sam Farr authored SDSF's enabling legislation, Assembly Bill 1965 of 1987 (now Public Resources Code (PRC) Sections 4660-4664), which provided for the protection and preservation of SDSF as an intensively managed educational and research forest. It also contained special provisions for the use of SDSF, including a limited amount of commercial timber operations on the property within SDSF in order to provide funds for the maintenance and operation of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Section 4660 on SDSF. AB 1965 is reprinted in Appendix A of this plan.

SDSF was formally transferred from its interim managers, The Nature Conservancy (TNC), to CAL FIRE on July 13, 1990. Former CAL FIRE Director Harold Walt, former Assemblyman Sam Farr, a representative of State Controller Gray Davis, and Steve Johnson of TNC dedicated the Forest, emphasizing the nature and purpose of this addition to the DSF system.

SDSF was the first of California's demonstration state forests to have an advisory committee formed to assist the Department in planning future management of the Forest. The Advisory Committee, also required by AB 1965, met monthly during the planning process to facilitate the creation of the original 1998 General Forest Management Plan. The Advisory Committee reconvened in 2011 and plays a vital role in reviewing the revisions to this updated version of the plan.

THE GENERAL FOREST MANAGEMENT PLAN

Following the acquisition of the SDSF property (see the Administration Chapter), TNC created an interim management plan for what they called the Soquel Creek Forest. Recognized as a temporary plan, TNC's document provided direction for current and future management decisions involving SDSF. The 1998 General Forest Management Plan incorporated elements of the TNC plan and information from other sources. It was developed with input from the public and resource professionals. Public workshops were held to obtain feelings, opinions, and factual information about the management of SDSF's forest resources. Furthermore, individuals representing many interests contributed data, publications, and personal knowledge for consideration through conversation with Forest staff. Public comments and concerns relating to

various subjects are summarized in each chapter. A supplemental document, titled Public Input to the Soquel Demonstration State Forest General Management Plan, presents all input received during the development of the draft plan.

Local resource professionals contributed a significant amount to the 1998 plan. Knowledgeable individuals served on the Advisory Committee, provided factual information about various resources, and composed elements of the plan itself. Without their assistance and experience, this management plan could not have been written.

The environmental impacts of the 1998 management plan were thoroughly investigated by Jones and Stokes Associates, Inc. in an Environmental Impact Report (EIR), State Clearinghouse Number 94023033, which was completed in compliance with the California Environmental Quality Act (CEQA). A primary objective of this EIR was to identify mitigation measures to reduce or avoid adverse environmental impacts that could result from implementation of any SDSF projects. As required by CEQA, mitigation measures identified from this EIR have been incorporated into a Monitoring Plan located in Appendix C.

FUTURE PLANNING

The Board of Forestry and Fire Protection (the Board) approved the original General Forest Management Plan on May 18, 1998. In 2003 and 2009, the Board reviewed and reauthorized the plan. No revisions were made at those times. As indicated throughout this plan, SDSF staff continues to formulate more specific management guidelines and planned actions. This revision of the plan includes new studies and the results of monitoring and research regarding the management of components such as fisheries, wildlife, watershed, archaeology, and timber.

This SDSF General Forest Management Plan will be in effect until it is either amended or a new plan is adopted in accordance with the procedures prescribed in PRC Section 4663. Working with the Advisory Committee, CAL FIRE will reexamine the General Forest Management Plan every five years, as per the policy of the Board of Forestry and Fire Protection, and determine whether any changes are necessary or desirable. This plan embodies the legislative intent of PRC Sections 4660-4664, and any subsequent amendments of this plan or any new plan must be consistent with the PRC except to the extent, if any, that subsequent legislation changes that intent. If changes are desired, the changes will be developed by CAL FIRE and presented to the Advisory Committee for consideration at one or more public meetings. The changes shall be approved by the Advisory Committee prior to review and adoption by the Board. (Statutory authority for final adoption of the plan rests with the Board per 0351.10 and PRC 4645.) Any changes will be subject to environmental review as provided by the California Environmental Quality Act.

CHAPTER 2: MANAGEMENT GOALS

SDSF's management goals represent a combination of legislation, policy, and public input. PRC Sections 4660-4664, the enabling legislation for Soquel Demonstration State Forest, is the preeminent authority with regard to the management of SDSF. Consistent with the objectives of that legislation to protect and preserve SDSF as an intensively managed, multifaceted research forest and to the extent not in conflict with that enabling legislation, SDSF will be managed in accordance with the state forest system legislation (PRC Sections 4631-4658) and Board of Forestry and Fire Protection policy.

Public Resources Code Section 4660 states that the intent of the Legislature in establishing the Soquel Demonstration State Forest is to provide an environment that will do all of the following:

- * Provide watershed protection for local communities and base-line monitoring and studies of the hazards, risks, and benefits of forest operations and watersheds to urban areas.
- * Provide public education and examples illustrating compatible rural land uses, including sustained yield timber production, as well as the historic development of timbering and forestry machinery, within the context of local community protection and nearby pressures.
- * Provide a resource for the public, environmental groups, elected officials, environmental planners, the educational community, and the media as an open environment for the inspection and study of environmental education, forestry practices, and effects thereof.
- * Protect old-growth redwood trees.

Section 4661 further states that CAL FIRE may permit a limited amount of commercial timber harvesting in order to provide the funds needed for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF.

Below is a listing of SDSF's general management goals which elaborate on the legislative intent. Other subjects and greater detail relating to the topics listed here can be found throughout the following chapters of this General Forest Management Plan.

RESOURCE PROTECTION AND ENHANCEMENT

1. Protect, restore, and enhance the significant natural values of the Soquel Demonstration State Forest.

2. Provide watershed protection and conduct baseline studies and monitoring of hydrological resources.
3. Demonstrate fire protection using a coordinated fire prevention and control system which includes education and enforcement of fire prevention guidelines, Forest patrol, and vegetation management including prescribed fire, fuelbreak construction, pre-attack strategies, and suppression tactics.
4. Improve fisheries and wildlife habitat to foster healthy populations and promote biodiversity.
5. Monitor, study, and implement controls for various forest pests using Departmental and outside specialists.
6. Monitor, study, and implement controls for invasive plant species.
7. Identify all significant archaeological and historical features and protect them during all management activities.
8. Conserve soil resources by reducing erosion resulting from flooding, earthquakes, logging activities, roads, and trails.

DEMONSTRATION AND EDUCATION

1. Conduct innovative demonstrations and education in forest management including silviculture, habitat diversity, logging methods, hydrology, resource protection, and recreation.
2. Provide forestry education opportunities for the public, forest landowners, the educational community, the media, natural resource professionals, and environmental groups.
3. Develop interpretive resources to help Forest visitors understand the various coast redwood forest communities and the basics of forest land management.
4. Establish a volunteer program to assist Forest staff in providing forestry interpretation for visitors.
5. Provide suitable public access and parking.
6. Plan for a Forestry Education Center to be designed and constructed to serve as the Forest's focal point for demonstration and education activities.

RESEARCH

1. Conduct research in forestry and natural resource management, including the benefits

and risks of forest operations in watersheds close to urban areas.

2. Serve as a laboratory for in-house projects and encourage research by other agencies, interest groups, and educational institutions.
3. Disseminate information obtained from the State Forest to appropriate individuals in an effective and timely manner.

TIMBER MANAGEMENT

1. Demonstrate sustained-yield timber harvesting practices through harvest operations that balance harvest rates with growth over time and are compatible with rural land use in Santa Cruz County, while promoting recreation opportunities, forest health, watershed protection, wildlife, and fisheries values as well as aesthetic enjoyment.
2. Protect old-growth redwood and old-growth Douglas-fir trees and recruit additional late-successional forest stands.
3. Incorporate demonstration, research, and restoration objectives into timber management activities whenever possible.
4. Research and implement hardwood stand management alternatives including modification to enhance wildlife habitat, utilization for various forest products, and conversion to softwood timber stands consistent with the legislative goals of PRC Sections 4660-4664.

RECREATION

1. Provide for recreational opportunities which are oriented toward foot, bicycle, and equestrian traffic and include trails, roads, and picnic areas. Limited camping may be permitted if consistent with Forest objectives.
2. Integrate recreation management, forestry education, resource protection and examples of timber harvesting so as to demonstrate how they can be compatible.
3. Unless specifically authorized, the activities that are prohibited include fishing, the use of motorized vehicles, shooting, hunting, camping, fires, and night time use.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Throughout the planning process, members of the public have indicated that adherence to

SDSF's enabling legislation, AB 1965, is legally required. Neighbors and visitors of the Forest have communicated that management goals and actions should abide by the written legislation. CAL FIRE intends to abide by this plan and to act consistently with the intention of the enabling legislation as expressed in this plan.

CHAPTER 3: PROPERTY DESCRIPTION

LOCATION AND BOUNDARIES

Soquel Demonstration State Forest is located near the center of Santa Cruz County, California, approximately eight miles northeast of the city of Santa Cruz (Figure 1). Positioned in the southern portion of the Santa Cruz Mountains, SDSF is eighteen air miles south of San Jose and within a two-hour drive of the San Francisco and Oakland metropolitan areas. Access to the property is via State Highway 1 or 17 and local county roads. The entrance to the Forest is from Highland Way, a county road in the Santa Cruz Mountains that connects State Highway 17 with Watsonville. Virtually all of the Forest's 2,681 acres are located within the East Branch of Soquel Creek watershed.

SDSF's boundaries were originally established by metes and bounds rather than the more familiar township and range system. Formerly part of the Shoquel Augmentation Rancho (a Mexican land grant that was later more commonly referred to as Soquel), this area has always been defined differently than the land which surrounds it. Some of the corners listed in the survey, prepared by George Dunbar of Dunbar Land Surveys, have been verified and are in place. The Santa Cruz County parcel numbers for the Forest are 098-101-04; 098-161-06; 098-351-01¹; and 099-181-02, 03, 04, 06.

ADJACENT OWNERSHIP

SDSF is bordered by both state and private property (Figure 2). The Forest of Nisene Marks State Park borders the State Forest for three and one-half miles along Santa Rosalia Ridge to the south. Approximately three-hundred-forty acres directly east of the Forest boundary are owned by Roger and Michelle Burch. This land is managed by Redwood Empire and includes the main entrance and parking area for the Forest off Highland Way. To the north and west, the adjacent ownerships are private rural-residential parcels, including the large holding of Spanish Ranch. Most of these parcels range in size from 1 to 80 acres. On the southwest border is the property containing the Olive Springs Quarry, owned by the CHY Company.


With the considerable amount of private property surrounding the Forest, public access is currently limited. The only undisputed public access points into the Forest are from Highland Way and The Forest of Nisene Marks State Park.

HISTORY OF OWNERSHIP

Prior to the arrival of Europeans, the Ohlone Indians inhabited the area. In the mid-1800's, the title to the 32,000-acre Shoquel Augmentation Rancho was awarded to Martina Castro de Depeaux viuda de Lodge viuda de Cota, the daughter of a Spanish Colonial soldier.

¹ Note that historic parcel number 098-161-02 was changed to 098-351-01 by the Santa Cruz County Assessor in 1998.


Figure 1. Locator Map for SDSF.



SDSF was contained within the rancho, and Martina gave this portion to her daughter, Antonia Lodge de Peck. Frederick A. Hihn, a German-born entrepreneur, was able to acquire portions of the Shoquel Augmentation through a discrepancy in legal title. He was particularly interested in Lodge de Peck's parcel and purchased it in 1863.

In the 1880's, Hihn established the Valencia-Hihn Company and began selectively logging the old-growth redwood on his lands to produce shingles, posts, and rails. Upon his death in 1913, Hihn's heirs assumed management of his lands and continued to harvest the area. In 1924, the Valencia-Hihn Company sold their land to the Monterey Bay Redwood Company (MBRC). The MBRC owned the State Forest property for 37 years and performed extensive harvesting in the 1920s and 1930s. They sold their property to the Glenco Forest Products Company of Sacramento in 1961, which later changed its name to the CHY Company. Eighteen years later, in 1979, CHY sold the State Forest portion of their land to the Pelican Timber Company. Additional details about the history of the Forest can be found in Archaeological and Historical Survey of Soquel Demonstration State Forest (Dillon, 1992).

Figure 2. SDSF and Adjacent Ownerships.



In 1988, Pelican was involved in a debt-for-nature land swap with the State of California and the Bank of America (see the Administration Chapter for more details). A result of this land swap was the creation of SDSF as authorized by former Assemblyman Sam Farr's Assembly Bill 1965. The Nature Conservancy acted as the interim managers of the Forest until its transfer to CAL FIRE in 1990.

CLIMATE

The climate of the Santa Cruz Mountains is Mediterranean, characterized by dry, warm summers and wet, cool winters. SDSF is often cool and damp because of the dense canopy of forest vegetation and its location on a north-facing slope. The average minimum January temperature is 38 degrees Fahrenheit, and the average maximum July temperature is 76 degrees Fahrenheit.

Most of the precipitation in the area occurs from November through April. The average annual rainfall for the East Branch of Soquel Creek is 44 inches (Linsley et al., 1992). At elevations above 2,000 feet, snowfall occurs about every other year and averages less than five inches total.

During the late spring and early summer months, Santa Cruz County often has foggy or cloudy skies. In the Forest, this marine layer is generally limited to early morning and late evening hours. Winds generally blow from the west or southwest (onshore) and are mild to moderate throughout the year. Strong winds, however, come in with winter storms and are strongest at higher elevations. Pressure gradients inland may occasionally cause strong northeasterly winds to occur.

SOILS AND GEOLOGY

SOIL TYPES

The parent material of soils found in SDSF is primarily sedimentary and consists of fine and coarse-grained sandstone, consolidated shale, weathered mudstone, and siltstone. Schist and intrusions of granitic rock are also present. There are nine soil series which developed from these parent materials; see Table 1 and Figure 3 (USDA, 2004). They are all deep and well-drained soils except for the Maymen Stony Loam which is a shallow, well-drained soil. According to the Natural Resources Conservation Service, most of the soils support watershed, recreation, and wildlife resources. Five of the soils (Ben Lomond, Felton, Lompico, Nisene, and Aptos) also support timber production, with the primary species being coast redwood and Douglas-fir.


GEOLOGIC ACTIVITY

In 1992, a detailed geologic study was completed by the California Geologic Survey (Manson and Sowma-Bawcom, 1992). This investigation resulted in a report which focuses on the process and degree of instability in both the State Forest and surrounding areas. The report, titled Geology, Slope Stability, and Earthquake Damage in Soquel Demonstration State Forest, includes maps of general geologic and geomorphic characteristics, landslide features (indicating the relative degree of stability), stream orders, roads to be considered for abandonment, and

Alquist-Priolo Special Studies Zones².

SDSF is seismically very active. The San Andreas Fault runs through the northeastern boundary and along the East Branch of Soquel Creek to the mouth of Ashbury Gulch, where it turns north. The Zayante Fault, part of the San Andreas Rift Zone, runs through the southwest edge of the Forest. The epicenter of the 1989 Loma Prieta Earthquake was located approximately two miles south of SDSF, in The Forest of Nisene Marks State Park. Numerous cracks and fissures dating from the 1989 earthquake have been located in the State Forest. Geologic activity, coupled with past fires and severe rain storms, has helped form the steep terrain found throughout the Forest. These events have also contributed to the many landslides present within the inner gorges of streams and along steep roadcuts. The numerous natural springs and sag ponds found throughout the Forest are also the result of past geologic activity.

Figure 3. Location of Soil Series in SDSF.



The elevation of SDSF ranges from 500 feet at the East Branch of Soquel Creek to 2,500 feet at the southeast corner on Santa Rosalia Ridge. The higher elevations occur in the southeast portion of the Forest and decrease along the ridge going southwest.

² Alquist-Priolo Special Studies Zones are areas along traces of the San Andreas Fault where geologic investigations are required prior to development.

Table 1. Soil Type and Associated Erosion Hazard of SDSF³.

SOIL TYPE	PERCENT SLOPE	PERCENT ACRES	EROSION HAZARD
Ben Lomond sandy loam	15-50	1.77	moderate to high
Ben Lomond sandy loam	50-75	11.61	very high
Ben Lomond-Felton complex	30-50	0.66	high
Ben Lomond-Felton complex	50-75	15.81	very high
Hecker gravelly sandy loam	50-75	0.48	very high
Lompico-Felton complex	5-30	7.44	moderate to high
Lompico-Felton complex	30-50	23.95	high
Lompico-Felton complex	50-75	14.71	very high
Madonna loam	15-30	0.88	high
Maymen stony loam	15-30	0.06	high
Maymen stony loam	30-75	1.97	high to very high
Nisene-Aptos complex	15-30	1.90	moderate to high
Nisene-Aptos complex	30-50	6.76	high
Nisene-Aptos complex	50-75	7.62	very high
Riverwash	-	2.94	-
Zayante coarse sand	30-50	1.44	moderate to high

WATER RESOURCES

The East Branch of Soquel Creek is a perennial stream that flows through the entire length of the Forest. It is fed by the perennial streams of Fern Gulch and Amaya Creeks from the north, and numerous unnamed intermittent and ephemeral streams. The total size of the East Branch watershed is approximately 19 square miles or 12,240 acres.

As previously mentioned, natural springs and sag ponds can be found in the Forest. The two largest springs are Sulphur Springs, located near Sulphur Springs Road, and Badger Spring,

³ From Soil Survey of Santa Cruz County, California (USDA, 2004).

located near the main picnic area. Badger Spring was at one time a developed water source as is evident by the remains of a spring box and steel pipes scattered around the area. A third spring, located east of Sulphur Springs along Hihn's Mill Road, was created by the 1989 earthquake. The natural springs of Sulphur and Badger, as well as other unnamed springs, are also special to SDSF. These springs, plus Amaya Pond, enhance particular biotic communities and offer various research opportunities.

Amaya Pond, a seasonal body of water, is located in the northwestern arm of the Forest. Approximately one-half acre in size, it is located on the east side of Amaya Creek Road, approximately one-third of the way down from Comstock Mill Road. (See Preliminary Biological Assessment of Soquel Demonstration State Forest, Santa Cruz County, California, [Holland et al., 1992] for more details about Amaya Pond.)

The portion of the East Branch that runs through the Forest is well known for its steelhead rearing habitat. The California Department of Fish and Wildlife prohibits angling in this part of the Soquel Creek watershed in order to protect this important resource. The Fisheries Chapter of this report contains more information on the creek and its fisheries assets.

Soquel Creek, including the East Branch, is also part of the domestic water supply for the local community. The lower portion of Soquel Creek serves as part of the natural groundwater recharge system for residents' wells and supplies surface water to a number of intakes along the creek. Soquel Creek is within the Central Coast California (CCC) Coho Salmon Evolutionarily Significant Unit (ESU). (See the Fisheries and Watershed Chapters for additional information on watershed condition, use, and management.)

ZONING AND GENERAL PLAN

The Santa Cruz County General Plan (1994) can be described as the county's "blueprint" for future development. It is a long-range policy document that looks at the future of the community and takes into account the types of development that will be allowed, the spatial relationships among land uses, and the general pattern of future development. The stated objective for timber production (Santa Cruz County General Plan, Chapter 5, and Objective 5.12) is to encourage the orderly economic production of forest products on a sustained yield basis under high environmental standards, to protect the scenic and ecological values of forested areas, and to allow orderly timber production consistent with the least possible environmental impacts.

Under the concept of zoning, various kinds of land uses are grouped into general categories or "zones". A zoning ordinance is the local law (refer to County Code) that spells out the immediate, allowable uses for each parcel within the County. Zoning regulates present development through specific standards such as lot size, building setbacks, and a list of allowable uses. Zoning must comply with the general plan. The purpose of zoning is to implement the policies of the general plan. The Santa Cruz County General Plan allows timber harvesting and associated operations, requiring approval of a Timber Harvesting Plan by the California Department of Forestry and Fire Protection for the following zoning designations: Timber Production (TP), Parks, Recreation and Open Space (PR) (except in the

coastal zone), Mineral Extraction Industrial (M-3), and the Commercial Agriculture (CA) (except in the coastal zone).

SDSF is entirely classified as Timber Production Zone (TPZ). This statewide zoning designation was created by the Forest Taxation Reform Act of 1976. Counties throughout the State were required to rezone parcels that met the definition of Timberland as defined in Government Code Section 51104(g) and consistent with Sections 51112 and 51113. TPZ land is devoted to and used for growing and harvesting timber and other compatible uses as defined in Section 51104(h). Compatible uses include but are not limited to: watershed management, fish and wildlife habitat management, outdoor education and recreation activities, and may include a residence or other structure as necessary for management of land zoned as Timberland Production.

Any development must meet not only the specific requirements of the zoning ordinance, but also the broader policies set forth in the local general plan. For the purpose of determining the development potential of rural parcels, the Santa Cruz County General Plan designates SDSF land as Mountain Residential. Objectives of this designation are to provide for very low density residential development in areas which are unsuited to more intensive development. Additional considerations include the presence of physical hazards and development constraints, the lack of public services and facilities to support higher densities, the protection of natural resources, the retention of rural character, and for maintaining sustainable use of natural resources.

CHARACTERISTIC FEATURES

A distinctive feature of SDSF is its proximity to the large urban areas of San Francisco Bay, Monterey Bay, and San Jose. This provides prime opportunities for urban children to experience forestry education on a first-hand basis.

The presence of steelhead trout in a portion of the Soquel Creek watershed also contribute to the special characteristics of SDSF. Once abundant along the entire west coast, steelhead populations have declined due to habitat loss and several other factors. The East Branch of Soquel Creek, the portion of Soquel Creek that flows through the Forest, supports a steelhead population and its required habitat. A very limited number of coho salmon and their habitat are also present in the watershed.

As mentioned above, the San Andreas Fault and Rift Zone are directly associated with SDSF. The effects of both ancient and contemporary seismic activity are apparent throughout the Forest. The history and future of this very active system make for an interesting addition to SDSF's abundant natural features.

Finally, the Forest contains archaeological and historical sites discovered during on-going archaeological surveys. The Archaeology Chapter of this plan describes the sites and their significance in detail. Both prehistoric and historic, these sites will enhance SDSF's demonstration and education programs.

CHAPTER 4: ADMINISTRATION

THE LEASE

On March 7, 1988, State Controller Gray Davis and the Bank of America settled a 13-year long lawsuit over unclaimed bank accounts. The settlement included \$35.7 million in cash and four undeveloped natural parcels in Tehama and Sonoma Counties. The property that is now SDSF was acquired during the settlement process and added to the package.

The settlement properties are held in a trust with the State as the beneficial owner and the Exchange Bank as trustee. The properties can be sold to pay unclaimed funds if they exceed the \$35.7 million in cash set aside for this purpose. It is doubtful, however, that this will ever happen.

At the time of the settlement, The Nature Conservancy (TNC) volunteered to act as steward for these properties. A 25-year lease was developed which stated that TNC would manage these properties and that past land use practices could continue. Any revenues generated from these activities were to pay for property taxes, operations and maintenance, natural resource enhancement, and access improvement projects.

The Nature Conservancy transferred their lease of the Santa Cruz County property (now SDSF) to CAL FIRE on April 18, 1990. CAL FIRE assumed management at that time and a dedication ceremony for SDSF was held on July 13, 1990. In 2013, at the end of the 25-year lease, the property will be transferred permanently to the State, free and clear.

Under the terms of the lease, both the trustee and the Controller have certain rights and responsibilities. The trustee's primary responsibility is to monitor the lessee's performance as managers of the properties. The Controller is responsible for the sale of any or all the properties in the event that cash assets are insufficient to satisfy all claims. As previously mentioned, this is unlikely to occur.

CAL FIRE ADMINISTRATION

Authority to *administer and operate* state forests in California comes from the Legislature and is contained in the Public Resources Code (Sections 4631-4664 and 4701-4703). Rules and regulations governing *use* of state forests are contained in the California Code of Regulations (Title 14, Sections 1400-1439 and 1510-1521). The State Board of Forestry and Fire Protection gives policy direction to the Demonstration State Forest Program, which is administered by the Director of CAL FIRE.

CAL FIRE is administratively broken into two Regions, each with a Region Chief who reports to CAL FIRE's Director. Each region includes units, and state forests are administered by a local Unit Chief. SDSF is in the Northern California Region (with headquarters in Redding) and is within the San Mateo-Santa Cruz Unit (headquartered in Felton). The State Forest office is located next to the CAL FIRE Soquel Forest Fire Station

at 4750 Soquel-San Jose Road in Soquel, California. When fully staffed, SDSF has a staff of five: Forest Manager, Assistant Forest Manager, half-time Office Technician, and two seasonal Forestry Aides. The Forest Manager is supervised by the Unit Chief.

The Forest staff is responsible for the on-site operation of the Forest as provided for by the Public Resources Codes, California Code of Regulations, and the State Board of Forestry and Fire Protection. Forest regulations, policy, and other issues prescribed by the Director of CAL FIRE are used to develop plans and procedures to govern development and perform maintenance of the Forest. The General Forest Management Plan will be reviewed and approved by SDSF's Advisory Committee (described below) and ultimately approved by the Board of Forestry and Fire Protection.

SDSF is an important resource for CAL FIRE training as well as for other agencies and affiliated organizations. CAL FIRE specifically uses SDSF for training of state personnel in chain saw operations, off highway driving for fire apparatus, wilderness first aid, search and rescue operations, swift water rescue, Forest Practice Regulations, Resource Management training for Joint Apprenticeship Committee requirements, archaeology, and erosion control practices.

FUNDING AND TAXES

The Demonstration State Forest Program, including SDSF, is funded through the regular annual state Budget Act. AB 1965 did not establish a separate fund for SDSF and it is not listed as a line item in the state budget. Revenues from all state forests are deposited in a special fund called the Forest Resources Improvement Fund (FRIF), providing money for the annual budgets of state forests.

Expenditures for all state forests are included in a single budget line item in the Department's annual budget. Soquel State Forest was added to the Department's state forest budget in the 1990-91 fiscal year with a minimum of staffing and operating expense. The Budget Change Proposal recognized that there would be little revenue from SDSF for the first few years and that FRIF would need to contribute over one million dollars in operating expenses before the Forest could produce revenue. It was also recognized that many years would pass before revenues would equal expenses.

The Department may permit a limited amount of commercial timber operations on SDSF in order to provide funds on a cumulative basis as necessary for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF. The enabling legislation requires a minimum level of timber harvesting, or "floor", to provide income for all costs of operation and for research and educational purposes of SDSF. The legislation authorizes a higher level of harvesting, or "ceiling", which shall not exceed long-term sustained yield (LTSY) and on a cumulative basis shall not exceed the level of timber harvesting necessary to provide the funds needed for the maintenance and operation expenses, reasonable capital costs, and other expenses incurred in fulfilling all the objectives identified in PRC Sections 4660-4664 on SDSF. These additional objectives include watershed protection and monitoring,

demonstrations of compatible rural land uses, and historic development of timbering and forestry machinery.

As a practical matter, the various objectives overlap and cannot be completely separated. For example, demonstration or experimental timber harvesting could qualify as research and public education as well as being an example of compatible rural land uses. The protection of old-growth redwood trees will occur under normal operations of SDSF and does not need to be identified as a separate purpose with separate funding.

Funding for SDSF needs to be increased over time in order to fulfill the objectives of PRC 4660-4664. In order to adhere to the administrative and budgeting processes currently in place while simultaneously assuring compliance with the limitations placed on the Department by the enabling legislation, the Department will publish accurate annual reports which will compile revenues and expenses itemized by program. The Department will post the Annual Report on the CAL FIRE website. Large capital expenses (e.g., for additional properties to provide proper access to SDSF or for construction of a Forestry Education Center and administrative facility) will be funded through the FRIF fund or any fund source approved by the Legislature. Amortization of these capital outlays may be included in the computation of cumulative expenses in the annual reports.

The State pays property taxes to the County of Santa Cruz on land values within SDSF. Additionally, purchasers of state forest timber are liable for payment of timber yield taxes according to Public Resources Code, Section 4654. SDSF's timber sale purchasers are required to file quarterly tax returns with the California Board of Equalization.

ADVISORY COMMITTEE

The legislation which created SDSF (AB 1965) called for the establishment of an advisory committee to assist with the development of SDSF's General Forest Management Plan. A main function of the committee is to act as a critical link between CAL FIRE and the community in the planning effort. This allows SDSF to learn what the community expects and their opinions regarding relevant issues.

The original Advisory Committee had nine members appointed by the Director of CAL FIRE in August of 1991. Five positions were specified by the legislation and four were added by the Director. The original committee consisted of one representative from each of the following:

- * State Board of Forestry and Fire Protection
- * Santa Cruz County Board of Supervisors
- * California Department of Parks and Recreation
- * California Department of Fish and Wildlife (CDFW - formerly the Department of Fish and Game)
- * The Forest of Nisene Marks State Park Citizens Advisory Committee
- * The Nature Conservancy
- * Soquel Creek Water District

- * Neighborhood Representative
- * Local Registered Professional Forester

The Committee held monthly meetings during the development of the 1998 General Forest Management Plan. The Advisory Committee became inactive following the approval of the 1998 Plan. The Advisory Committee reconvened in 2011 to provide input on updating the Plan. Most of the groups listed above continue to be represented on the Committee. However, the Soquel Creek Water District removed itself from the Committee and the State Park Citizens Advisory Committee has been disbanded. The latter groups have been replaced with representatives from the Stewards of Soquel Forest and from the Resource Conservation District of Santa Cruz County. Each member serves a three-year term or until the General Forest Management Plan updating process is completed (whichever is longer). Once these conditions have been met, one third of the members' terms shall expire on the last day of each year. Following the approval of the updated Plan, the group will meet twice a year or as needed to review progress on plan implementation and contribute to on-going planning activities.

SAFETY

The remote and rustic character of SDSF makes safety an important management consideration. Forest visitors need to be informed of safety issues and hazards inherent to the Forest. Roads, trails, and facilities are maintained in safe condition. The staff coordinates with local CAL FIRE Emergency Command Center, the Loma Prieta Volunteer Fire and Rescue, the county Sheriff's Office, and other agencies and groups for emergency medical response. Forest personnel, including volunteers, are trained in first-aid; permanent Forest personnel maintain certification as Emergency Medical Responders. Additionally, search and rescue organizations are allowed to train in the Forest so as to develop their skills and better acquaint themselves with the terrain.

The following safety protocols are currently in place:

- * Restriction and regulation signs are posted at Forest entrances. Hazards, safety issues, and the primitive nature of the area are stated on signboards and in the SDSF brochure.
- * The staff works with the California Department of Parks and Recreation to provide trail maintenance, safety, and coordinated emergency response along the common boundary.
- * All trails, roads, and emergency helicopter landings are regularly inspected and maintained. Fallen trees and other hazards are removed as needed to maintain safe conditions.
- * Motorized vehicles owned by the public are prohibited beyond designated parking areas. Exceptions are made through special permission and for management, patrol, and emergency purposes.

- * Coordination with CAL FIRE's Emergency Command Center in Felton and the county Sheriff's Office ensures effective emergency response in the Forest. All responses for emergency assistance will be recorded and compiled, including calls for police, fire, medical, or search and rescue services (see Appendix C).
- * Volunteers from the Stewards of Soquel Forest and the SDSF chapter of the International Mountain Bike Association provide trail maintenance and patrols, assistance to forest visitors, and support for emergency response.

LAW ENFORCEMENT

Forest regulations and policies are posted on signs and enforced through patrol and citation of violators (also see the Resource Protection Chapter). CAL FIRE peace officers, authorized under the California Penal Code, will be used to detain violators, with local law enforcement agencies providing backup when necessary. The CDFW wardens will enforce fishing, hunting, and trapping laws. Apprehension and prosecution of violators shall be actively pursued. Violators will generally be cited and expected to appear in court, but may be taken into custody if warranted. Methods to prevent illegal activities and alternatives which curtail unwanted behavior will be explored and developed to reduce law enforcement problems.

State Forest trespass violators will normally be cited under sections of the California Code of Regulation (Title 14). Illegal trespass includes removal of trees without a permit and parking, camping/campers, or building in the Forest.

Marijuana cultivation in the Forest has been relatively minor. Since the dedication of the Forest, the remnants of nine old gardens have been found and four active gardens have been eradicated. The general inactivity of marijuana cultivation is due in part to the majority of the Forest being south of the East Branch of Soquel Creek with a northern exposure. Furthermore, SDSF is inaccessible by vehicles and, therefore, is not readily available for use.

Detection of marijuana gardens will occur during normal patrol activities or as leads are developed. Most detection efforts will depend on flights by the Santa Cruz County Sheriff's Marijuana Eradication Program. Information about gardens found in SDSF will be referred to the Sheriff's Office.

ACCESS NEEDS

As stated in the Soquel Demonstration State Forest Recreation Study Final Report (McNally and Hester, 1993) and the Recreation Chapter, Forest access is a significant problem and complex issue. The inability of the public to drive to the Forest when county roads are closed and trespass across private property to and from the Forest are challenges.

The SDSF property came into state management with two verified legal access routes (see

the Roads and Other Improvements Chapter for more details). The first is an administrative and public right-of-way through the Burch property off Highland Way. The second is across CHY Company property, through the Olive Springs Quarry, which includes a right-of-way for administrative use only.

Many visitors come in via Ridge Trail from The Forest of Nisene Marks State Park. Some recreationists, particularly equestrians, have permission to pass through private property to enter the Forest. The public's use of Comstock Mill Road is prohibited due to neighbors' opposition.

There is an obvious need to develop improved access into the State Forest, particularly along the south and west ends where most trespassing occurs. It is important to provide Forest visitors with additional safe and legal access in order to reduce trespass onto private property. Furthermore, alternative access points are necessary when landslides close Highland Way and/or Eureka Canyon Road.


LAND ACQUISITION PRIORITIES

Land acquisition to improve access to SDSF is a top priority. This has been actively pursued since the dedication of the Forest. Negotiations with various neighboring property owners regarding appropriation and/or easements have been ongoing. Parcels formerly owned by the Noren family were purchased in 2005. These parcels make up about 9.5 acres and are shown on the Acquisition Map (Figure 4). The State acquired the properties for the express purpose of providing an improved access to SDSF in anticipation that successful negotiations with the CHY Company would eventually add the other key piece needed for this access. This parcel is adjacent to the Noren properties and would provide a suitable road alignment into the SDSF ownership, as well as a site for the Forestry Education Center and other facilities (see the Demonstration and Education Chapter for more information).

Another potential acquisition parcel is the Burch property which contains the entrance to the Forest (see Figure 2 in the Property Description Chapter). The benefits of this procurement would include access control, additional areas for resource management and recreational visitor use, and a location for an informational kiosk and restroom. There would also be clear authority and responsibility for maintenance of and improvements to the bridge, roads, and parking area.

Other options for improving public access aside from direct land purchases will be considered. These could include purchase or trade of rights of way or easements, lot line adjustments or land trades. Any land acquisitions funded by timber harvest revenues from SDSF shall be limited to those which improve access to SDSF or otherwise directly enhance SDSF. All acquisitions for SDSF will comply with CEQA.

Figure 4. Recent Acquisition Map for SDSF.



COOPERATION WITH ADJACENT LANDOWNERS

As the previous CAL FIRE Director, Harold Walt, indicated at SDSF's dedication ceremony in 1990, it is important for SDSF to cooperate with their neighbors. Forest staff will continually work with the community regarding local issues. These issues include but are not limited to fire prevention, trespass, watershed impacts, fisheries restoration in the East Branch of Soquel Creek, mitigation of recreational and timber harvest impacts, emergency response and invasive species management. SDSF staff works with local schools in forestry education and also coordinates fire prevention programs with the local CAL FIRE Unit.

COOPERATION WITH OTHER AGENCIES AND INSTITUTIONS

SDSF cooperates with other agencies in resource protection, fire prevention and suppression, law enforcement, and safety. Cooperation is also encouraged for demonstration, forestry education, and university research projects. Other agencies that work with SDSF include the California Department of Parks and Recreation, California Department of Fish and Wildlife, California Geological Survey, National Oceanic and Atmospheric Administration Southwest Fisheries Science Center, National Marine Fisheries Service, US Fish and Wildlife Service, US Geological Service, the County of Santa Cruz, and the Resource Conservation District of

Santa Cruz County. Cooperative institutions of learning and research include UC Santa Cruz, Cabrillo College, California Polytechnic State University at San Luis Obispo, San Jose State University, UC Berkeley, UC Cooperative Extension, Washington State University, West Valley College and local public schools.

*THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE
1998 GENERAL FOREST MANAGEMENT PLAN*

Public comments and concerns involving the administration of SDSF have largely been focused on the issue of safe and legal access but have also touched upon the FRIF program and Advisory Committee composition.

Suitable public access into SDSF is a major concern for neighbors, users, and Forest staff. As previously stated, there is an obvious need to develop adequate entry and exit points and to stop illegal trespass. Several individuals have made specific requests regarding the location of access points, usually focusing on the area from Olive Springs Quarry to Comstock Mill Road. However, neither end is owned or controlled by the state and only administrative access is allowed. SDSF will continue working with its neighbors and on potential access acquisitions to alleviate this complex and pressing issue.

During the process of establishing SDSF and creating the General Forest Management Plan, questions regarding the purpose and use of FRIF monies have been raised. As required by the Public Resources Code, all revenues from SDSF's timber sales must go into FRIF to be managed and allocated. As part of a state agency, SDSF will adhere to the requirements of FRIF as outlined by law.

Finally, neighbors of the State Forest have expressed that they would like greater representation on SDSF's Advisory Committee. Letters were written to former CAL FIRE Director Richard Wilson by both neighbors and the committee chairman requesting a review of the public's concerns. After a careful and lengthy evaluation, the Director determined that the committee composition was adequate due to strong current local representation.

CHAPTER 5: BIOTA

INTRODUCTION

Biota are defined as the flora (vegetation) and fauna (wildlife) which inhabit a particular area. When people envision forests such as SDSF, biotic elements are what generally come to mind. Biota, however, are only one element of an ecosystem, small pieces of the larger puzzle.

An ecosystem has been defined as "the interacting populations of plants, animals, and microorganisms occupying an area, plus their physical environment" (Hunter, 1990). The physical environment consists of abiotic factors such as soil, water, space, and climate. The presence and actions of humans make up a yet another component of ecosystems. Though humans often consider themselves to be separate entities, at a basic level they are part of the biota and are intricately connected to everything in an ecosystem. When biotic, abiotic, and human components of a forest ecosystem are working together in dynamic balance, diverse biota and healthy forests are attained. (Dynamic balance refers to the continual interaction of ecosystem components which leads to a balanced yet constant state of change.) In real life, there are rarely clear boundaries between adjacent habitat communities or even ecosystems. Ecosystems and habitats blend and overlap but can be given a label based on general wildlife, vegetation, and location features. Management planning for ecosystems or habitat communities which looks at landscape patterns caused by this overlap can benefit all inhabitants, whether they be stationary (e.g., plants) or mobile (e.g., animals).

Prescribed fire plays an important role in reducing fuel loads, recycling nutrients and sustaining plant communities. Prescribed burning has been a tradition, ritual and tool since prehistoric times in the Santa Cruz Mountains and has shaped the environment that we see today. Many species and habitats have become rare due to the suppression of fire. By conducting prescribed burns under controlled conditions, hazardous forest fuels are reduced which limits the risk of a catastrophic wildfire while at the same time providing for ecological benefits. Further discussion about prescribed burning is discussed in the Resource Protection Chapter.

In this chapter, the biota are described for each habitat type, or community, found in the Forest. Only the most common plants and animals found in these communities are mentioned, but complete flora and fauna lists are located in Appendix B. For a detailed account of the biota of SDSF, refer to Preliminary Biological Assessment of Soquel Demonstration Forest, Santa Cruz County, California (Holland et al., 1992). For specific information about trees (size, abundance, etc.), see the Timber Management Chapter. Fisheries resources are described in a separate chapter.

HABITAT COMMUNITIES

Communities consist of the living organisms collectively found in an ecosystem (Hunter, 1990). Even though considerable overlap often occurs, communities are individually labeled

and classified for research, inventory, and education purposes. They are generally named for the dominant plant species within each community. The dominant plant species is dependent on specific environmental conditions (e.g., soil, climate, water) that further characterize the community. Because communities overlap, plants indicative of one habitat type may be found in others. Poison oak, for example, can be found growing in virtually all of the communities of SDSF but is most abundant in drier habitat types. Also, some fauna considered to be permanent residents of a particular community actually travel through several communities. These animals, including large mammals (deer, bobcat, gray fox, mountain lion), can be found throughout SDSF as they search for food, water, and shelter.


In 2006, a timber inventory was completed which gathered tree data on plots located on a grid layout. This data was analyzed along with digital image segmentation and aerial imagery to classify the vegetation using the California Wildlife Habitat Relationship (CWHR) habitat classification system. Using this protocol six CWHR habitats were classified on SDSF. These are by order of abundance; Redwood, Montane Hardwood-Conifer, Coastal Oak Woodland, Douglas-fir, Mixed Chaparral and Annual Grassland. The CWHR system further classifies each type by the amount of canopy closure. The amount and extent of tree canopies are used in the CWHR system to help predict which wildlife species may be supported by these ecosystems. On SDSF there are Moderate (M) density (40% to 59% canopy closure) and Dense (D) (60% to 100% canopy closure) density classes. The CWHR system then further categorizes by the average tree size classes. On SDSF the size classes correspond to 3 (pole size, 6 to 11 inches diameter at breast height), 4 (small tree size, 11 to 24 inches diameter breast height) and 5 (medium/large tree size, greater than 24 inches diameter breast height). Figure 5 displays these types with their corresponding locations and acreages.

Table 2. Key to California Wildlife Habitat Relationship Vegetation Type Codes.

SDSF CWHR Vegetation Types		
CWHR Classification	Average Tree Size (inches dbh⁴)	Canopy Closure (%)
Coastal Oak Woodland 3D	pole size (6 - 11)	60 - 100
Coastal Oak Woodland 4M	small tree size (11 - 24)	40 - 59
Coastal Oak Woodland 4D	small tree size (11 - 24)	60 - 100
Douglas-Fir 4D	small tree size (11 - 24)	60 - 100
Montane Hardwood-Conifer 3D	pole size (6 - 11)	60 - 100
Montane Hardwood-Conifer 4D	small tree size (11 - 24)	60 - 100
Redwood 4D	small tree size (11 - 24)	60 - 100
Redwood 5D	Medium/large tree size (>24)	60 - 100
Mixed Chaparral		
Annual Grassland		

⁴ Diameter at breast height

Figure 5. Vegetation Type Map for SDSF.



COAST REDWOOD FOREST COMMUNITY

The coast redwood forest is dependent on areas of high moisture and comprises about 1,229 acres of SDSF. In its northern range in California, large continuous stands of redwood are formed. Southern redwood forests are restricted to moist canyon slopes and riparian zones since more moisture is available in these areas. As its name suggests, the dominant tree of this community is coast redwood, the majority of which in SDSF are second growth (regrowth after original clearcutting). Small groves of old-growth redwoods occur at Badger and Sulphur Springs, and individual old-growth redwood trees are scattered throughout the Forest. Other common trees of this community are tanoak, Douglas-fir, and madrone. Understory species found in the redwood community of SDSF are redwood sorrel, California hazel, wild ginger, and western sword fern. Common wildlife residents include pygmy nuthatch, Steller's jay, and Trowbridge's shrew. The redwood community provides nesting habitat, cover, and food for birds and a variety of small mammals. Redwood forests have comparatively little forage value for deer and other large mammals, however, limiting use of this habitat type.

MONTANE HARDWOOD-CONIFER COMMUNITY

This forest community is dominated by both conifers and hardwoods, often in a closed canopy. About 969 acres of the Forest are categorized as Montane Hardwood-Conifer. Primary tree species include Douglas-fir, Shreve oak, and redwood; and to a lesser extent madrone, California bay-laurel, black oak, and big leaf maple (Holland et al., 1992). Montane Hardwood-Conifer forests are on the drier slopes above the redwood community, though the two overlap considerably. Also included in this community are stands with significantly more Douglas-fir which are classified as CWHR Douglas-fir and compose about 214 acres.

Common understory species include poison oak, California blackberry, vetch, toyon, and yerba buena. Familiar wildlife species are Merriam's chipmunk, dusky-footed woodrat, western gray squirrel, California slender salamander, acorn woodpecker, sharp-shinned hawk, and screech owl. Evidence of feral pig activity is also found throughout the community. Oak trees located in these areas have very high value for wildlife, providing both nesting and roosting substrates and a food source through acorn production.

Some stands in this community historically supported a more significant conifer component than exists today. These stands failed to regain the original species distribution following intensive harvesting during the 1930s and 1940s. These stands are dominated by tanoak and Shreve oak. Understory vegetation is typically minimal except for occasional conifers where canopy openings permit (See the Timber Management Chapter for additional hardwood management discussion).

COASTAL OAK WOODLAND COMMUNITY

The Coastal Oak Woodland community is composed primarily of hardwoods and covers about 262 acres. These hardwood stands appear to be long dominated by a combination of

coast live oak, Shreve oak, and tanoak. They occupy sites less favorable to conifers. As with the Coast Redwood and Montane Hardwood-Conifer communities, other hardwoods such as madrone, bay-laurel and black oak are also present. These stands include large senescent oaks with unique structural features beneficial to wildlife. Acorn production is important to many species, especially as winter range. Understory species are similar to the Montane Hardwood-Conifer community. Any management activities conducted in these stand types will be conducted solely for long term maintenance. Management activities may include selective harvesting of hardwoods for fuelwood as well as tree planting where conditions are favorable for increasing the diversity of species and stocking. Several research opportunities exist in the Coastal Oak Woodland Community especially related to Sudden Oak Death Death (*Phytophthora ramorum*) and other hardwood specific pathogens. See the Research Chapter for information on hardwood research projects.

RIPARIAN COMMUNITY

Riparian communities are named for the intermittent or continual presence of fresh water rather than the vegetation of such areas. Riparian communities are located along the edges and floodplains of streams or surrounding lakes. In SDSF, an abundant riparian community exists along the floodplain of the East Branch of Soquel Creek and to a lesser degree along Amaya Creek. This community is dominated by deciduous hardwoods such as white alder, bigleaf maple, black cottonwood, and California sycamore. Along with these trees, red and yellow willows grow in dense clumps along the banks of the East Branch. Horsetails and hedge nettles are common ground cover along the edges. Wildlife residents include vireos, warblers, Pacific-slope flycatcher, long-tailed weasel, and raccoon. Pacific newts, brown-colored salamanders with bright orange bellies, are abundant in the riparian community and a great delight to young forest visitors. The Pacific tree frog, thought to be common, is only found in a few locations (Holland et al., 1992). Additionally, large colonies of ladybug beetles gather along creeks to overwinter and breed.

Riparian communities are the most productive and resilient terrestrial habitat type for wildlife because of structural diversity and the presence of water. Many migratory songbirds are dependent on riparian habitat for breeding and foraging. Large mammals use the riparian zone as a water supply, and incorporate it into their home ranges. The riparian community is probably the most significant habitat type in the Forest due to its high value to wildlife and limited regional occurrence.

OTHER COMMUNITIES AND ADDITIONAL SPECIES

There are a few other limited communities present in SDSF. Freshwater marshes are areas where the soil stays wet the majority of the year, supporting characteristic vegetation. These marshes usually occur along the perimeter of ponds, at springs, near shallow pools of streams, or in areas of high water tables. The freshwater marsh community in SDSF is scattered, including only Amaya Pond and a few natural springs (Sulphur, Badger, and a couple of small, unnamed springs caused by or increased from the Loma Prieta earthquake). Wildlife residents include migratory waterfowl, great blue heron, black phoebe, belted kingfisher, and garter snakes.

Approximately four acres of SDSF are comprised of grassland and mixed chaparral communities. Some of the grassland areas are natural, due to soil conditions conducive to permanent grassland establishment. Other grassland areas are the result of past disturbance. SDSF's grasslands primarily contain wild oats and annual fescue grasses. Most grasslands are being encroached upon by coyote brush, lupine, poison oak, and Douglas-fir. Common wildlife residents of the grasslands are the gopher snake and Botta pocket gopher.

There is one significant chaparral stand located in the Longridge Road area on the south facing slope above Soquel Creek. Chaparral species are also found mixed in the Montane Hardwood-Conifer and Coastal Oak Woodland vegetation types along the exposed ridge tops and on south-facing slopes at higher elevations. These dry locations support the fire-adapted woody shrubs of manzanita, buck brush, coyote brush, and chamise. Common wildlife residents are Bewick's wren, California towhee, scrub jay, western fence lizard, and brush rabbit.

FUNGAL RESOURCES

A local mycological organization has identified a wide variety of mushrooms in the Forest (refer to Appendix B). Fungi are broken into three categories based on their relationship to the immediate environment: mycorrhizal, saprophytic, and parasitic. Saprophytic fungi occur on wood that is already dead whereas parasitic fungi attack and can kill live trees. The most common fungi found in SDSF are mycorrhizal.

Mycorrhizal species form a symbiotic relationship with the trees they grow under. These organisms grow around the rootlets and collect water and trace nutrients for use by trees. The trees in turn provide carbohydrates to the fungi. Trees and mycorrhizae, therefore, depend on one another for optimum health. According to a representative of the Fungus Federation, SDSF is a reasonably healthy forest because of the wide occurrence of mycorrhizal fungi (Nathan Wilson, personal communication, 1993).

In general, mycological research of California's wildlands has been minimal, particularly regarding conditions conducive to fungi growth. SDSF provides ample opportunities for mycological research including fungi population analysis and succession of fungal species in the Forest. To protect the fungal resources of SDSF, a permit system for mushroom collection is maintained that prohibits any collection for commercial purposes.

FERAL PIGS

Since the establishment of SDSF, feral pigs (*Sus scrofa*) have been observed throughout the Forest. Feral pigs are an introduced species and are present throughout the Santa Cruz Mountains. Their populations shift from year to year based on weather patterns and forage availability, and seem to be transient between the Forest and the surrounding areas. The pigs are a problematic species because they can cause damage and/or alter the Forest's native communities. They frequently wallow in soft soil and wet, marshy areas which can contribute to sediment in watercourses and disturb flora and fauna that use these habitats.

Their rooting/foraging behavior often damages roads and encourages invasive plant species. Feral pigs also aggressively defend themselves and under certain conditions may pose a threat to Forest users. In the past, the Forest has received calls about damage done to neighboring properties because of the belief that the feral pigs reside in SDSF. Requests were made for increased monitoring and management of the feral pig population.

The population is managed through the California Department of Fish and Wildlife's (CDFW) Wild Pig Depredation permitting process. When increased pig activity is noted during patrols (rooting along roads, tree damage, wallowing at sumps and ponds), depredation permits are issued.

BULLFROGS

Bullfrogs are non-native aquatic vertebrates from the genus *Ranidae*. These frogs are found throughout the range of the Santa Cruz Mountains. Perennial ponds and areas of perpetual calm water are capable of supporting populations of bullfrogs, once introduced, or which migrate from another location. Bullfrogs are aggressive feeders, and given their large size are capable of directly consuming or creating intense indirect competition for smaller, native frog species such as the California red-legged frog and the foothill yellow-legged frog, which are both found on SDSF. Bullfrogs are also known to spread the chytrid fungus *batrachochytrium dendrobatidis* (Bd) to healthy populations of native aquatic organisms. Bullfrogs have not been observed at SDSF, although due to the known presence of California red-legged frog and foothill yellow-legged frog at SDSF, any future bullfrog observation is cause for action.

Habitat for bullfrogs within SDSF is very limited. Amaya Pond is an annual pond and calm water areas within the East Branch of Soquel Creek are often flushed out by winter rains. Neighboring ponds are often annual in nature or drained on a regular basis.

Aquatic biological monitoring takes place on SDSF on a regular basis with herpetological specific surveys occurring since 2011. If bullfrogs were to occur within SDSF, there is a high likelihood of detection. If bullfrogs are encountered, control measures could include mechanical culling or removal of habitat for as long as necessary to ensure population control.

CORVIDS

Corvids are birds from the genus *Corvus*, and in the Santa Cruz Mountains include common species such as crows, ravens and jays. Unlike many other bird families, corvid fitness and reproduction increase with human development. Corvids are especially significant in the Santa Cruz Mountains because they are major predators on eggs and chicks of the endangered marbled murrelets. Large populations of corvids are frequently associated with campgrounds in forests where unnatural foodstuffs are found. Big Basin State Park, a nesting site for murrelets, works specifically to contain trash and food at campgrounds as well as

inform the recreating public about the sensitivity of nesting murrelets. Corvid populations are unlikely to be higher in SDSF than in the surrounding privately managed timberlands. Camping is not allowed on a regular basis at SDSF and visitors are expected to pack out all trash. Signs are posted to pack your trash and Forest staff makes a great effort to pick up any trash left behind in the parking area on a regular basis. Marbled murrelets have never been detected at SDSF, although the old-growth stand near Badger Spring is considered suitable habitat for the species. Surveys were conducted in 2003 and 2004 by biologist David L. Suddjian, and no murrelets were detected on any of the surveys.

INVASIVE SPECIES

Invasive species control is an ongoing process at SDSF. Forest staff along with Ben Lomond Conservation Camp Crews and volunteers log hundreds of person hours each year to reduce and control French broom. The primary method to reduce invasive species (predominantly French broom and jubata grass) has been a continuous mechanical removal approach by pulling plants and roots or cutting stems. Other control methods such as herbicide applications and flaming have not been employed extensively, however more efficient and cost effective approaches will be considered. Herbicide use is part of the overall invasive species control program and efforts to control large continuous infestations of French broom with herbicides began in 2013. Combining hand-pulling with herbicide spraying every other year has proven effective for controlling French broom on Santa Cruz Water District properties.

Efforts will be made to control existing invasive plant populations, reduce opportunities for further spread of existing species, and prevent the introduction of other species not currently present on SDSF. During the preparation of the Fern Gulch THP, a botanical survey was conducted where 24 non-native species were identified which are representative of species found throughout SDSF (Table 3). Approximately half of the species identified are considered a high priority for control.

Many non-native plant seeds prefer bare mineral soil to germinate. Preventing the establishment of new or expansion of established populations is emphasized through THP mitigations that minimize soil disturbance and the amount of exposed mineral soil following operations. Additional project mitigations include avoiding the introduction of weedy grasses into project areas, avoiding the use of invasive seeds for erosion control, and using only certified weed-free straw (preferably rice straw) for mulching to prevent erosion. Short-lived cereal crops like barley and rye have been used for erosion control locally in Santa Cruz County and have not been found to be invasive.

Table 3. Invasive Exotic Plant Species Occurring at SDSF.

Common Name	Scientific Name
*French Broom	<i>Genista monspessulana</i>
*Periwinkle	<i>Vinca major</i>
*Poison hemlock	<i>Conium maculatum</i>
*Jubata Grass	<i>Cortaderia jubata</i>
*English Ivy	<i>Hedera helix</i>
*Eupatory	<i>Ageratina adenophora</i>
*Subterranean clover	<i>Triflorium subterraneum</i>
*Italian thistle	<i>Carduus pycnocephalus</i>
*Italian ryegrass	<i>Lolium multiflorum</i>
*Forget me not	<i>Myosotis latiflora</i>
*Bermuda buttercup	<i>Oxalis pes-capre</i>
*Orchard grass	<i>Dactylis glomerata</i>
Bull thistle	<i>Cirsium vulgare</i>
Cutleaf geranium	<i>Geranium dissectum</i>
Rough cat's-ear	<i>Hypochaeris radicata</i>
Common chickweed	<i>Stellaria media</i>
Field bindweed	<i>Convolvulus arvensis</i>
Broadleaf fleabane	<i>Conyza sumatrensis</i>
Spiny sowthistle	<i>Sonchus asper</i>
Soft chess	<i>Bromus hordeaceus</i>
Bur-Chevril	<i>Anthriscus caucalis</i>
Sticky chickweed	<i>Cerastium viscosum</i>
Fiddle dock	<i>Rumex pulcher</i>
Olive	<i>Olea europaea</i>

* The survey botanist recommended these species have the highest priority for control.

PLANT SPECIES OF CONCERN

The California Natural Diversity Data Base (CNDDDB) was queried on May 10, 2010 to collect information on listed species and species of concern known to occur in the Laurel quadrangle that contains the Soquel Demonstration State Forest. A total of eight plant species are state and/or federally listed as threatened or endangered. In addition, six plant species are categorized as CNPS (California Native Plant Society) List 1B. The plants of List 1B are rare throughout their range with the majority of them endemic to California. Most of the plants of List 1B have declined significantly over the last century. List 1B plants constitute the majority of the plants in CNPS' Inventory with more than 1,000 plants assigned to this category.

All of the plants constituting List 1B meet the definitions of Sec. 1901, Chapter 10 (Native Plant Protection Act) or Secs. 2062 and 2067 (California Endangered Species Act) of the California Department of Fish and Wildlife Code, and are eligible for state listing. These species must be fully considered during preparation of environmental documents.

A nine quad search of processed CNDDDB data, centered on the Laurel quadrangle identified 48 plant species. One plant species is CNPS List 1A, 32 are CNPS List 1B, and 15 are federal and/or state listed as threatened or endangered. Although there is no suitable habitat for most of these taxa on Soquel Demonstration State Forest, the number of species listed provides a rough indicator of the extent of plant species of concern in the general vicinity of the State Forest (see Appendix B for a list of plant species known to occur at SDSF).

SPECIAL-STATUS WILDLIFE SPECIES

Although the biological assessment of the Forest conducted in 1991-92 found no threatened or endangered plant or wildlife species, this is not the case today. The Central California Coast (CCC) coho salmon Evolutionarily Significant Unit (ESU) was federally listed as threatened in 1996 and relisted as endangered in 2005. In 2012 the range of CCC coho was extended south to include Soquel and Aptos Creeks. Coho salmon south of the San Francisco Bay were listed as endangered under the California Endangered Species Act in 1995. Steelhead within the CCC Distinct Population Segment (DPS) were federally listed as threatened in 1997. See the Fisheries Chapter for more information about the status and recovery plans for coho and steelhead.

The California red-legged frog has been found on the Forest and is federally listed as threatened. Additionally, a few wildlife species of special concern to the State of California have been seen in SDSF (Holland et al., 1992). Those species observed were the foothill yellow-legged frog, western pond turtle, sharp-shinned hawk, Cooper's hawk, and golden eagle. The long-eared owl and yellow warbler may also occur in the Forest, but they have not been observed. Suitable breeding, nesting, or foraging habitats exist in the Forest for all species observed except the golden eagle.

The Laurel quadrangle CNDDDB query for animal species indicates the occurrence of two federally listed endangered insects and two federal or state listed threatened and endangered fish species. In addition, one amphibian is federally listed as threatened and one amphibian, one reptile, and one mammal species are presently California Department of Fish and Wildlife Species of Special Concern. See Table 4 for further information.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

A major focus in the management of SDSF's biota involves species evaluation and monitoring, particularly for special status species. Comments have largely focused on fauna though concern about edge vegetation and introduced plant species has been expressed. As stated in the Management Guidelines and Planned Actions below, SDSF plans to monitor selected biotic elements of the Forest and evaluate effects of forest management activities on the condition of those resources.

Another primary concern is the restoration of degraded habitats and maintenance of exceptional resource values in SDSF. Emphasizing this concern are comments regarding

management of habitats individually (e.g., manage riparian habitats separately from mixed evergreen habitats) and management activities which allow interior forest species to thrive. Many commenters feel that development and management of SDSF should be performed in such a way that biotic resources are preserved or improved. The Management Guidelines and Planned Actions deal with these issues as well.

Finally, input has been received regarding the desire for details on exactly how SDSF will manage and maintain its biotic resources. Since this management plan is intended to be general, specific information on the what, how, why, and when of SDSF management activities will be outlined in other formats. More specific management strategies will be developed for different areas and habitats based on anticipated management endeavors and research and monitoring results.

Table 4. Special-status Vertebrate Fish and Wildlife Species Occurring or with Potential to Occur at Soquel Demonstration State Forest.

SPECIES	LEGAL STATUS ^a FEDERAL/STATE	HABITAT	OCCURRENCE IN SDSF ^b
American peregrine falcon	/FP	Nests in cliffs, forages in a variety of habitats	4
Merlin	/WL	Wintering only, frequents open habitats	2
Marbled murrelet	T/E	Nests in old-growth conifer forest; forages in pelagic habitats	4
Golden eagle	BCC/FP	Nests in cliffs and trees in forests and woodlands; forages in grasslands, shrublands, and chaparral	2
Osprey	--/WL	Nests in snags and spike-top trees; forages in open water	2
Cooper's hawk	--/WL	Nests and forages in woodlands and forests; also forages in open habitats	2
Sharp-shinned hawk	--/WL	Nests and forages in conifer forest habitats	1
Long-eared owl	--/CSC	Nests and forages in riparian and woodland habitats	2
Olive-sided flycatcher	--/CSC	Tall conifers used for nesting, perching; Forages over open/low vegetation	2
Purple martin	--/CSC	Nests and forages in woodland and forest habitats in tree cavities	4
Vaux's swift	--/CSC	Nests in large tree cavities with a preference for redwood and Douglas-fir habitat	2
Black swift	--/CSC	Nests on cliffs, steep rocky outcrops, canyons near water	2
Yellow warbler	--/CSC	Nests and forages in riparian habitats	2
Foothill yellow-legged frog	--/CSC	Occurs in streams with rocky substrate	2
California red-legged frog	T/CSC	Occurs in slow-moving streams, pools and ponds	2
Western pond turtle	--/CSC	Occurs in pools, ponds, and lakes	2
Steelhead (Central CA Coast ESU)	T/CSC	East Branch Soquel Creek	1
Coho salmon(Central CA Coast ESU)	E/E	East Branch Soquel Creek	3
Pallid Bat	--/CSC	Buildings, rock outcrops	3
San Francisco dusky-footed woodrat	--/CSC	Forested habitats of moderate canopy and moderate to dense understory	2

^a Status codes:

Federal: T = threatened, E = endangered, P = proposed for listing as threatened or endangered, BCC = bird of conservation concern USFWS

State: CSC = species of special concern, T = threatened, E = endangered, WL = watch list, FP = fully protected
^b 1 = confirmed nesting/reproduction, 2 = observed, 3 = not observed, 4 = unlikely to occur

Table 4 Sources:

Jones and Stokes Associates, Inc., 1996

California Department of Fish and Game Special Animals List, July 2009

California Department of Fish and Game State and Federally Listed Endangered & Threatened Animals of California, January 2010

California Department of Fish and Game, California Natural Diversity Database, Laurel Quadrangle, May 10, 2010

MANAGEMENT GUIDELINES

1. Ongoing monitoring will be performed to detect listed and special status species. Monitoring will include keeping current with state and federal lists as well as conducting periodic floral and faunal surveys. Inventories will emphasize special-status species expected to be present but not yet observed in SDSF. New findings will be added to current species lists. Every consideration will be given to protecting these species and their habitat as required by law and determined by a qualified biologist.
2. Old-growth trees will be protected as outlined in SDSF's authorizing legislation, AB 1965. Areas of old-growth redwood have been located and protection will be provided in all phases of forest management. Additionally, areas have been designated to promote late-succession stands of trees (see the Timber Management Chapter for more details).
3. Restore, maintain, or enhance resource values of native habitat communities to promote natural diversity and stability. Measures to achieve this include:
 - * snag recruitment and retention
 - * preservation of appropriate logs and other woody debris
 - * maintenance of natural ponds and springs
 - * protection of riparian zones for use as movement corridors for wildlife
4. Achieve mutual benefit with timber harvesting, demonstration and education, and recreation programs while respecting native biotic elements. Wildlife habitat improvements, such as those mentioned in Management Guideline 3 above, will be considered during the planning and implementation of timber sales, demonstration and education activities, and recreational facilities. Conversely, possible impacts of present or future wildlife and vegetation management projects on other management objectives will be studied prior to project approval and implementation.
5. Control and/or eradication of exotic invasive plant species utilizing Integrated Pest Management techniques will be incorporated into management activities, as appropriate. Ben Lomond Conservation Camp crews are utilized for hand pulling and cutting. Forest volunteers also help with the removal of invasive plants. Additional

efforts including herbicides and/or flaming⁵ will be employed where and when appropriate.

6. Control mushroom collection by issuing permits for scientific, educational, and personal use. Mushroom gathering for commercial purposes is prohibited.

PLANNED ACTIONS

1. Encourage researchers to study wildlife habitats, populations and unique characteristics at SDSF. Continue to support the UC Santa Cruz Puma Project and the newly developed herpafauna survey efforts; both are described further in the Research Chapter.
2. Examine the California Natural Diversity Database (CNDDDB) reports during project planning and incorporate measures into all project development and monitoring processes for all known species as well as special status species that may be present. Submit CNDDDB Field Survey forms to CDFW for any sightings of listed, rare or special status species.
3. Conduct preharvest and post-project surveys to identify active nest sites of all raptors and special-status bird species that may occur in the Forest. Those species that may occur in SDSF are Cooper's hawk, sharp-shinned hawk, long-eared owl, and yellow warbler. In addition, preharvest surveys will be conducted for California red-legged frog, foothill yellow-legged frog, and southwestern pond turtle.
4. Continue to evaluate the feral pig situation, and develop management strategies and actions to diminish existing problems.
5. Continue to build the inventory of old-growth trees across SDSF (further described in the Old Growth section of the Timber Management Chapter).
6. Continue to use mechanical methods for controlling invasive species with Ben Lomond Conservation Camp Crews and volunteers each year. Use additional follow up treatments to improve effectiveness, such as herbicide application or flaming where and when appropriate. Extra effort will be concentrated on new populations prior to them becoming established and producing seed banks.
7. Conduct biological assessments incrementally in new project areas. These will include results of CNDDDB reports, botanical surveys, and other site specific assessments.

⁵ Flaming uses a propane torch to kill plants when they are very small by applying heat. This method is very effective for controlling weeds such as broom, is faster and cheaper than pulling, and is more selective than herbicide use. This method can be safely used without risk of fire hazard during periods with cool temperatures and high vegetation moisture levels in the targeted vegetation.

CHAPTER 6: FISHERIES

INTRODUCTION

Approximately 7.5 miles⁶ of fish-bearing streams flow through SDSF, including the East Branch of Soquel Creek (5 miles), Amaya Creek (2 miles), and Fern Gulch Creek (0.5 miles). The East Branch currently provides valuable steelhead trout spawning and rearing habitat and could provide essential habitat to promote reintroductions and recovery of coho salmon. Based on the amount of stream miles within SDSF and the high potential for ecological restoration of lost or degraded habitat components such as instream complexity and floodplain connectivity, SDSF provides a unique opportunity to support recovery of this invaluable fisheries resource.

RANGE AND LEGAL STATUS: STEELHEAD TROUT AND COHO SALMON

The fishery resources of greatest concern in SDSF are the steelhead trout and coho salmon. Anadromous fish such as steelhead and coho spawn (mate and lay eggs) in freshwater creeks or rivers but spend most of their adult lives in the ocean.

Historically, coho salmon spawned in coastal streams from the Bering Sea of the Arctic and the coast of Japan to the Monterey Bay in California. The steelhead's range extended further to the north coast of Baja California. Steelhead and coho populations have been declining throughout their entire range, both in fresh and salt water, for decades due to a number of factors including habitat loss and fragmentation in freshwater systems. In California, numbers decrease from north to south, with the southernmost population of steelhead and coho at the greatest risk of extinction. This is particularly important since Soquel Creek and Aptos Creek represent the southernmost watershed along the Pacific Coast with recent confirmed observations of coho salmon. While the Soquel Creek watershed is within the range of steelhead and coho, there has been a considerable decline in numbers for both species, with coho observations limited to a few individuals in 2008.

Legal Status: Steelhead

While steelhead were technically removed from the genus *Salmo* nearly 40 years ago and incorporated in the genus *Oncorhynchus*, they are often still referred to by their traditional common name steelhead trout. For the purposes of this document, we will simply refer to the species as "steelhead". The genus and species for steelhead is *Oncorhynchus mykiss*. It should be noted that steelhead are genetically identical to rainbow trout with the fundamental difference between the two fish having to do with life history. While steelhead are anadromous and move between the ocean and freshwater, rainbow trout are year-round residents in freshwater and do not migrate to or from the ocean. Steelhead within Soquel Creek are part of the Central California Coast (CCC) Distinct Population Segment (DPS) and were federally listed as Threatened under the Endangered Species Act in August of 1997.

⁶ According to the Soquel Creek Salmonid Assessment and Enhancement Plan (Alley, 2003) fish migration barriers exist in Fern Gulch up 382 feet from the confluence with Soquel Creek and partial barriers exist on Amaya Creek at 2,091 feet, 2,181 feet and 3,886 feet from the confluence with Soquel Creek.

The CCC DSP stretches from the Russian River in the north to Aptos Creek in the south. As such, Soquel Creek is near the southernmost portion of the range of this DPS. Steelhead from the Pajaro River south to the Santa Maria River are within the South Central California Coast (SCCC) DPS and are considered to be a different genetic population. The National Marine Fisheries Service (NMFS), which is charged with protection of federally listed anadromous fish, is in the process of developing a recovery plan for the CCC steelhead and the draft plan is expect to be released to the public in early 2014. According to Jon Ambrose (pers com) of NMFS, plan recommendations will closely overlap with the recommendations put forth in the recently published CCC Coho Recovery Plan (NMFS, 2012). The steelhead plan will provide additional details and recommendations for recovery of steelhead within this DPS and, in conjunction with the coho plan, can be used to identify and guide recovery actions on SDSF. While steelhead are not technically listed under the California Endangered Species Act, the California Department of Fish and Wildlife (CDFW and formerly the California Department of Fish and Game) issued the 1996 "Steelhead Restoration and Management Plan for California" in an effort to focus conservation actions on the protection of this species. Finally, CDFW also develops an annual "Statewide Steelhead Task List" to support and guide funding actions through the Fisheries Restoration Grants Program (FRGP).

Legal Status: Coho

Coho salmon within Soquel Creek belong to the CCC Evolutionarily Significant Unit (ESU) of the species. This ESU was first listed under the Federal Endangered Species Act as Threatened in October of 1996 (Federal Register, 1996) and then relisted as Endangered in June of 2005 (Federal Register, 2005). At the time of listing, the ESU extended from Punta Gorda in the north to the San Lorenzo River in the south. In March of 2012, the NMFS extended the southern range of the ESU to include Soquel and Aptos Creeks. This decision was based on, "observations of coho salmon in Soquel Creek in 2008, genetic analysis of tissue samples indicating that the fish from Soquel Creek were closely related to nearby coho salmon populations in the ESU, and the ecological similarity of Soquel and Aptos creeks with other nearby creeks that support coho salmon" (Federal Register, 2012). Coho salmon south of the San Francisco Bay were listed as endangered under the California Endangered Species Act in 1995. California Fish and Game Commission extended the range of the listing designation north to Punta Gorda, Humboldt County in 2005.

Both the State and Federal governments have developed recovery actions for this species. The Recovery Strategy for California Coho Salmon was adopted by the California Fish and Game Commission in February 2004. The primary objective of the Recovery Strategy is to return coho salmon to a level of sustained viability, while protecting the genetic integrity of the ESU. For the Big Basin Hydrological Study Unit, of which Soquel is a part, the key recovery recommendations focus on protection of instream flows, upgrading of culverts for fish passage and mobilization of Large Woody Debris (LWD), and implementation of high priority restoration actions for coho from watershed plans. NMFS published the "Final CCC Coho Recovery Plan" in September of 2012 and this plan provides specific data and recovery recommendations for Soquel Creek. The plan identifies SDSF as a recovery partner for restoration actions in the East Branch of Soquel Creek, including Amaya Creek, which are considered core areas for species recovery. The Plan calls for a recovery target of 1,122

returning adult coho and highlights the following five high priority immediate restoration actions:

- Delineate reaches possessing both potential winter rearing and floodplain areas
- Implement a long term study project in Soquel Demonstration State Forest to demonstrate effective LWD projects to citizens of Santa Cruz County
- Promote conjunctive use of water for water projects whenever possible
- Provide incentives to water rights holders willing to convert some or all of their water rights to instream use
- Re-establish a naturally reproducing run of coho salmon

LIFE HISTORY AND HABITAT USE

As anadromous fish species, both steelhead and coho utilize freshwater for mating/spawning, egg development and early maturation and move to the ocean for a period of rapid growth and weight gain prior to returning to freshwater to spawn. The life cycle begins with the development of eggs into young fish in freshwater streams. Once the eggs hatch, young fish develop in the watercourse and gradually make their way to the ocean. Steelhead trout in this area typically spend two years in fresh water, although a few may spend additional years inland before migrating out to sea. The length of time spent in streams depends on environmental and genetic factors, and some individuals never migrate (Barnhart, 1986). Research by Smith (2005) suggests that one of the key environmental factors may be food supply and growth. According to these data, size is a critical factor in determining when a juvenile steelhead will leave freshwater, and once juveniles reach approximately 3.5 inches in forklength by the fall, they tend to out-migrate the following spring. While growth in freshwater habitats in SDSF may require at least two years due to slow growth rates, steelhead growth can increase substantially in food rich lagoon environments like the Soquel Lagoon (Alley 2011).

In order to acclimate to saltwater, both steelhead and coho go through a process of smoltification prior to entering the ocean and juvenile fish leaving freshwater are referred to as smolts. Steelhead and coho along the California coast usually spend two years in salt water, attaining sexual maturity and storing fat for their journey back up their natal streams to spawn and restart the life cycle process. While females of both species and most males spend two years in the ocean, a portion of male coho, called jacks, are known to return to freshwater after one year in the ocean. Due to the abundance of food, anadromous fish species experience most of their growth once they have reached the ocean. Therefore, jacks are generally identified due to their smaller size and weight.

While there are many similarities in the life cycle for these species, there are some key differences that should be highlighted. These include:

- Timing of adult return to freshwater and spawning- Coho are known to return to their natal streams in the southern portion of the ESU between November and January with the height of spawning peaking in February and March (NMFS 2012, from Moyle, 2002). Steelhead spawners generally return to their natal streams later in the winter

- and spawn through April or May depending on climatic conditions.
- Juveniles freshwater rearing- Whereas steelhead often spend multiple years as juveniles in freshwater, the vast majority of juvenile coho salmon spend only one year in freshwater before going to the ocean. As such, coho smolts are generally younger and smaller than most steelhead smolts.
 - Post spawning adults - While coho adults always die following spawning, some steelhead adults can return to the ocean after spawning and may repeat that cycle to spawn up to four times, though most repeat spawners do so only twice.

The basic stream attributes for steelhead and coho spawning, rearing, and migration include cool water temperature, high concentrations of dissolved oxygen, adequate water depth, sufficient pool space, and low sediment levels (Barnhart, 1986 and Anderson, 1995). Riparian habitat also provides a favorable microclimate for amphibians. Riparian vegetation also stabilizes streambanks and can play a major role in either supporting or degrading habitat for these fish. Riparian zones are strips of water-loving vegetation and associated organisms that follow the path of watercourses. Essential to healthy aquatic ecosystems, these zones help maintain favorable water quality and provide important food and habitat conditions. Trees along the water's edge shade the water, maintaining cool temperatures for anadromous fish spawning and rearing, as well as maintaining ground cover that intercepts eroded materials from upslope, minimizing the amount of sediment that enters the stream. Additionally, vegetation adds food and nutrients to the water for use by both fish and aquatic invertebrates. Large woody debris falling into the water provides cover for fish, collects and controls the movement of sediment, and creates deep scour pools favored by rearing juveniles.

Water temperature is a critical habitat component that can have dramatic effects on growth and development of steelhead and coho. A complication to understanding the effect of temperature on salmonids is that food availability is the key variable that governs how water temperatures affect fish. While both salmonid species have mortality thresholds with respect to water temperature, higher water temperatures do not always directly relate to lower growth and productivity. Water temperatures above 21.1°C (70°F) make it difficult for coho salmon and steelhead to extract oxygen from the water. Optimal rearing temperatures for juveniles are 7.22-14.4°C (45-58°F) for steelhead and 11.67-14.4°C (53-58°F) for coho (Ligon et al., 1988). Temperatures between 14°C and 21°C (57°F and 69.8°F) may have a positive impact on growth if there is ample food supply to keep up with the increased metabolic demand of fish caused by higher water temperatures. Conversely, temperatures at and below the lower end of optimal can slow metabolism significantly and result in muted growth rates, translating to lower ocean survival rates.

Table 5 provides details linking fish life stage with habitat requirements. All of these habitat conditions need to be considered when working to restore, maintain, or enhance anadromous populations. Data from the NMFS 2012 CCC Coho Recovery Plan highlight the need to prioritize restoration actions that increase the extent and availability of "off-channel" habitats such as floodplains, backchannels, alcoves and tributaries. The Plan also calls for implementation of projects that increase the amount of LWD in the stream. Both LWD and off-channel habitats are particularly important for coho, but also valuable to steelhead, for

providing refuge to adult and juvenile fish during high flows in the winter and low flows in the summer. LWD aids in the sorting of stream bed materials. In the winter, when flashy flows result in high instream velocities, off-channel habitats and LWD can provide slow water sheltering areas for fish of all sizes. During the summer, deep pools formed through scour downstream of LWD provide salmonids with cool water refuge and cover from predation. Perennial off-channel habitats such as ponds, alcoves and back-channels can provide some of the highest quality summer rearing habitat with high levels of primary productivity and insect production.

LOCAL FISH POPULATIONS

Anadromous fish populations in Soquel Creek and along most of the Central Coast have declined significantly since the late 1960s. While steelhead declines have been significant, the 2012 CCC Coho Recovery Plan sums up the status of CCC coho as, “... *gravely close to extinction. Despite being listed under the Federal and California Endangered Species Acts, populations of CCC coho salmon continue to decline precipitously. Immediate and focused action is essential to increase the survival of, and provide the highest protection for, remaining populations (NMFS, 2012).*”

While there are scant data on coho population numbers in Soquel Creek over the past 50 years, there is a significant body of archaeological data that indicates the historic range of this species extended as far south as the Pajaro River and possibly the Salinas River. In addition to natural runs, we also know that coho salmon were planted into the East Branch of Soquel Creek in the 1930s originating from the Brookdale, Big Creek, Prairie Creek and Fort Seward hatcheries (Anderson, 1995). Coho salmon were thought to be extinct in Soquel Creek in the 1990s and most of the first decade of this century, until a small population of juvenile fish was observed in 2008 near the entrance of Hinckley Creek below SDSF. For the purposes of the Final CCC Coho Recovery Plan (NMFS, 2012), these juvenile coho form the basis of the estimate of two adults in Soquel in 2008 (i.e., at least one spawning pair of coho were in Soquel for these juveniles to exist). Genetic analysis of tissue samples indicates that the 2008 coho salmon in Soquel Creek were closely related to nearby wild coho salmon populations in the ESU.

Table 5. Habitat Requirements and Vulnerability by Each Salmon Life Stage (NMFS, 2012).

<p>Eggs: Incubation requires clean water, free of contamination and siltation. Disturbance of a single “red” (nest of eggs) could result in the death of thousands of salmon embryos.</p>	<p>Freshwater Streams</p>
<p>Alevins: After hatching, alevins remain nestled in the small spaces between the gravels, and feed from their attached yolk sacs. They are highly vulnerable to siltation and scour. Once the yolk is absorbed, the young salmon emerge from the gravels.</p>	<p>Freshwater Streams</p>
<p>Juveniles: Deep cool pools are critical for the summer rearing juvenile’s survival. Riparian vegetation helps support some of the insects consumed by juveniles, provides cover from predators (when recruited to streams can create wood-formed pools), and limits solar radiation to streams keeping water temperatures cool. Tree roots stabilize streambanks and create habitat structure. Large woody debris or downed wood creates cover and refugia for the tiny salmon to reside during high velocity flows. Pools and wetlands provide shelter from high flows, predators, and help filter sediments from the water column.</p>	<p>Freshwater Streams</p>
<p>Smolts: Juvenile salmon undergo a physiological change known as “smoltification” enabling them to transition, in estuaries or lagoons, for a life adapted to saltwater. Smoltification can occur primarily within the freshwater areas, or in the nearshore environment. Smolts need adequate flow from upstream rearing areas to be able to travel downstream to estuaries. Estuaries should provide cover and adequate feeding habitats to facilitate the transition into the ocean. Estuaries should be deep to provide cool temperatures and buffered with freshwater to dilute seawater (Moyle, 2002). The quality of these areas has implications to the survival of smolts entering the ocean environment.</p>	<p>Freshwater Streams, Estuaries, Lagoons, and Ocean</p>
<p>Sub-Adults/Adults: Maturation occurs during ocean residency over a two year period leading up to the adult salmon’s return to streams of their birth. The patterns of migration in the ocean vary, and shifts in ocean conditions affect food, migration patterns and survival. Fish in the ocean need adequate supplies of food to facilitate rapid growth. As the salmon return to their natal stream to reproduce, they once again undergo change from saltwater to freshwater; they depend on the near shore and estuarine environments for this transition.</p>	<p>Ocean</p>
<p>Spawners: Migration begins after heavy late fall or winter rains breach sand bars of coastal streams, allowing fish to move into lagoons (Moyle, 2002). Once the adult spawners arrive at their home river or stream, they need adequate flows, cool water temperatures, deep pools and cover to rest and hide as they migrate upstream. Females seek clean, loose gravel of a certain size in highly oxygenated riffle type flow water for laying their eggs. The site must remain stable throughout egg incubation and emergence, and allow water to percolate through the gravel to supply oxygen to the developing embryo.</p>	<p>Ocean, Estuaries, Freshwater Streams</p>

There are significant amounts of data on the historic steelhead populations in Soquel Creek. Soquel Creek was historically considered one of the most important steelhead spawning and

rearing streams in Santa Cruz County (Titus et al., 2005). Based on data collected during surveys in 1959, the juvenile abundance in this system corresponded to an adult steelhead run of approximately 500-1,000 spawning pairs or 1,000-2,000 adult fish (Becker and Reining 2008). Alley (2006) calculated adult steelhead population indices for Soquel Creek between 1997 and 2005 as an average of 523, with a range between 360-780 adults. These data were calculated based on estimates of juvenile population size and an application of an adult return model and these estimates are considered the best available data for the watershed.


Local fish population inventories along the East Branch of Soquel within SDSF began in 1993 and are conducted annually in cooperation with NOAA's Southwest Fisheries Science Center (SWFSC). This work is conducted by electrofishing at five sites: four on the East Branch of Soquel Creek and one on Amaya Creek (Sogard et al., 2009). The surveys assist with monitoring the fish populations in fish bearing creeks that run through SDSF and also Support research performed by biologists from NOAA's SWFSC to monitor steelhead growth and migration. The graph below (Figure 6) shows a summary of data for the time period 1993 to 2012.

Historic populations of both coho and steelhead throughout most of California were supplemented through much of the 20th century by releases of hatchery fish for commercial or recreation purposes. Principal hatchery production for the Central California Coast steelhead originates from the Warm Springs Hatchery on the Russian River and the Monterey Bay Salmon and Trout Project on a tributary of Scott Creek. The most recent planting of hatchery-raised steelhead in SDSF (i.e., the East Branch of Soquel Creek) occurred in the 1930s when steelhead trout from the Brookdale and Big Creek hatcheries were released in the East Branch of Soquel Creek (M. McCaslin pers. comm.). The Monterey Bay Salmon and Trout Project stocked steelhead annually in the main stem of Soquel Creek (downstream of SDSF) in the 1980s and 1990s. Hatchery steelhead planting was discontinued in Soquel Creek in 2001 because of CDFW's concern regarding genetic integrity of planting fish that originated in the San Lorenzo River stock (Alley, 2001 and 2002).

No production/mitigation hatcheries (hatcheries that produce fish with the goal of increasing recreational and commercial harvest or for mitigation purposes) for CCC coho salmon currently exist. The two hatchery operations in the CCC ESU (noted above) are captive broodstock facilities operated expressly for conservation and recovery purposes with significant oversight by CDFW and NMFS. Coho salmon reared at these two facilities are listed under the Endangered Species Act. In Santa Cruz County the goals of the captive broodstock program include increasing population size, maintaining genetic diversity, and producing sufficient numbers of fish to promote straying into neighboring streams (Sturm et al., 2009).

In order to help protect remaining steelhead trout populations in Soquel Creek, CDFW prohibits angling in the East Branch of Soquel Creek (Fish and Wildlife South Central District Regulation, Title 14, Section 7.00 [e] [4]). This regulation was adopted by the

Figure 6. Estimated Juvenile Steelhead Population Comparison Index Reaches Soquel Demonstration State Forest, 1993-2012



California Fish and Game Commission in December of 1981 and became effective in March, 1982. In spite of this regulation, poaching of adult steelhead during winter spawning is prevalent and continues to affect fish populations. To avoid adverse impacts of public use on fish in SDSF, Forest staff and CDFW will conduct ongoing patrols to enforce prohibitions on fishing and fish harassment.

Other species of fish within the boundaries of SDSF include the Pacific lamprey, resident rainbow trout (population above Ashbury Gulch), Sacramento sucker, prickly sculpin, and northern threespined stickleback. Additional fish species that could be found in the East Branch include coast range sculpin and California roach.

HABITAT AND RESOURCE CONSIDERATIONS

HABITAT LOSS AND DEGRADATION

The loss of habitat for both winter and summer rearing is thought to have had a significant impact on the anadromous fish populations of the Soquel Creek drainage. Habitat availability and quality has declined due to diversion and overdrafting of water by residents, past logging practices, flood control measures (including LWD removal), increased development along the creek (including vegetation removal), pollution, and naturally unstable hillslopes. These impacts have synergistically increased water temperature and sediment inputs, altered stream flow patterns, reduced habitat complexity, and decreased the amount of surface water during the critical summer months and periods of drought. (See the Watershed Assessment Chapter for details on SDSF's watercourse conditions.)

SDSF comprises about 10% of the watershed area of the Soquel Creek basin. Over the years, the combination of increased sediment and lowered water levels has resulted in a considerable loss of pool habitat and a simplification of the channel throughout the Soquel Creek watershed. Making matters worse, wholesale removal of LWD from the 1950s to the 1990s further exacerbated the loss of pools as well as the disconnection of the channel from many "off-channel" habitats. With systematic removal of LWD, channels tend to incise with the removal of the natural grade control that LWD can create, leading to decreased bank stability/increased bank erosion as well as disconnection from side channels, alcoves, and floodplains that have not experienced the same rate of incision. Since formation of habitat complexity elements like sheltered pools for winter refuge and summer rearing or gravel bars for spawning are directly linked to changes in streamflow velocity (i.e., sediment deposition where water is slow and scour where water is fast), loss of LWD and its natural ability to affect water velocity can result in simplification of the stream, loss of pools and more uniform water velocities. Loss of covered pool habitat and disconnection of off-channel habitats, along with general fresh and saltwater habitat degradation (producing low marine survival rates), are believed to have directly contributed to the nearly complete disappearance of the coho salmon along California's central coast. Extremely low population numbers make it clear that human intervention and cooperation with recovery actions are essential for coho to become viable again and for existing steelhead populations to rebound (NMFS, 2012). Restoration of overwintering refuge from high water velocities both in the channel and "off-channel", as well as summer rearing habitat, will benefit recovery of both steelhead

and coho populations.

The diversion and overdrafting of water are significant problems in the Soquel Creek drainage, especially along its lower reaches. Near the town of Soquel, a portion of the creek has dried up on various occasions, including the summers of 1991, 1992, and 1994. Some residents along the creek use the water for agricultural as well as domestic needs. The Soquel Creek Stream System was formally adjudicated by Decree No. 57081, Superior Court for Santa Cruz County. The Decree was entered March 14, 1977 in Book 2731, page 581 of Official Records. There are water allotment requirements, but the requirements of fish were not considered when maximum amounts for residences and businesses were allocated. Although Soquel Creek was adjudicated, no water master was appointed and no diversion rates were independently measured to confirm that the adjudication is being followed. Because young steelhead (and coho) will move both upstream and downstream as upper portions of streams dry out in the summer, drying of downstream reaches limits available space and foraging areas for the entire population and could limit the ability of juvenile salmonids to gain access to wetted reaches and habitat in SDSF.

Greater public awareness and response regarding value and current status of the aquatic resources of the Soquel drainage are essential for garnering support for implementation of fish-friendly management actions and implementation of ecological restoration projects. Information, education and programs to help residents conserve water, allow the creek to flow in its natural channel, preserve riparian corridors, understand the value of LWD, and prevent accelerated erosion should be pursued. While improving the anadromous fish resource within SDSF is a start, it is not enough because Soquel Creek's fisheries require enhanced habitat conditions along every reach of the watercourse and throughout the watershed.

HABITAT RESTORATION AND ENHANCEMENT

Fish habitat at SDSF will be enhanced through implementation of habitat enhancement projects in conjunction with ongoing timber operations. The NMFS 2012 Recovery Plan for Central California Coast Coho Salmon (NMFS, 2012) recommends increasing the quantity of large wood in the channel as one of the highest priorities for Soquel Creek. In response, a Large Woody Debris (LWD) and Habitat Complexity Project has been designed for the East Branch of Soquel Creek. The proposed project is also part of the Integrated Watershed Restoration Program (IWRP) for Santa Cruz County and designed collaboratively with NMFS, CDFW, CAL FIRE, the California Geological Survey, NOAA's Southwest Fisheries Science Center (SWFSC), *Alnus* Ecological and the Resource Conservation District of Santa Cruz County. In an effort to collaboratively identify the best opportunities for fisheries habitat restoration along the creeks of SDSF, a group of fisheries scientists, resource specialists, and ecosystem restoration experts from the agencies list above, walked nearly two miles of East Branch in November of 2010 and identified restoration opportunities. The group agreed that these reaches were lacking complexity, that channel incision had left large areas of floodplain disconnected from the channel, and that SDSF had a unique opportunity to implement an array of different type projects to benefit fisheries. The group identified eight potential project sites and then refined the project list down to five sites. The final

project, as designed, entails placing LWD along a 0.7 mile section of the creek in four, 200-foot reaches. Three of the four reaches each contain three LWD elements, and the fourth has one LWD structure. Each element includes one to four pieces of LWD. In order to address the potential of LWD to mobilize and move far downstream, beyond the boundaries of SDSF, the project was designed to use a significant number of trees that are between 1.5 times and two times the width of the streambank and have their rootwads still attached. Between the heavy rootwads and long trunks, restored structures will closely mimic trees naturally falling in the stream due to landslide or bank undercuts and the structures will be unlikely to move a significant distance downstream, even in major storm events. Due to the size of the LWD required for each site (60 feet long or longer and up to approximately 50 inches diameter breast height) and the desire for the project to be a demonstration for private landowners, SDSF excavated 12 riparian redwood trees and dropped the entire tree and roots into the stream. A pilot project to determine the feasibility of this type of activity was successfully implemented in 2012 at one of the four sites. The other three sites were constructed in the late summer/early fall of 2013. Two additional rootwads with 25 foot logs attached were imported from the Fern Gulch timber sale for use in the project area.

Preliminary data from California Geological Survey indicates that the LWD projects at the pilot site have already had a significant effect on channel complexity with pools developing, the channel becoming more sinuous, and gravel bars forming between the wood structures (Reynolds, 2013). A fifth site for this LWD project is a stream bank repair site just east of Hihn's bridge where a fish-friendly revetment system was installed with rootwads and other components that will serve the dual purpose of repairing the road and providing shelter and pools for fish. This road repair project was completed in August 2014. A long term monitoring plan has been developed by NOAA's Southwest Fisheries Science Center (see Monitoring Section below for more details). Note that the baseline data collection for the monitoring work was funded by the State Coastal Conservancy and will aggregate biological and physical effectiveness data from LWD restoration sites on Soquel Creek at SDSF with those from a site on San Vicente Creek in northern Santa Cruz County.

While these first five sites represent an important start to the process of reversing historic trends in loss of habitat and significant impacts to salmonid populations, a number of other fisheries restoration opportunities have been identified along the East Branch of Soquel Creek in SDSF including two potential sites to reconnect historic backchannels/alcoves and a number of opportunities to enhance and reconnect floodplains. A complete halt to removal and/or cutting of LWD within SDSF is perhaps the lowest cost and most effective action to take to improve conditions for listed salmonids within the Forest. CAL FIRE has the unique opportunity to take the lessons learned from these projects and the related changes in management and work across the landscape with private forest landowners to use THPs and other means to implement additional stream and floodplain restoration projects.

MONITORING

A fisheries resources assessment was completed and used to develop a draft Fisheries Management Plan in 1995 (Berlekamp and Sutfin, 1995). The assessment inventoried the

condition of the fisheries population as well as habitat status within SDSF. The management plan includes prioritized improvements, monitoring systems, research opportunities, and funding sources (see Appendix C). Subsequent fisheries assessments in the watershed have been completed by the Santa Cruz County Resource Conservation District (Alley, 2003) and by NMFS (NMFS, 2012).

Fish population inventories began in 1993 and are conducted annually in cooperation with NMFS. This work is conducted by electrofishing at five sites: four on the East Branch of Soquel Creek and one on Amaya Creek. The surveys assist with monitoring the fish populations in fish bearing creeks that run through SDSF and also support research performed by biologists from NOAA's SWFSC to monitor steelhead growth and migration (Sogard, 2009).


In-stream temperature loggers are installed at seven locations in SDSF to measure and record stream water temperatures throughout the dry season. An additional temperature logger is also installed at one site to measure and record air temperature during the same time period. In 2006, the California Regional Water Quality Control Board for the Central Coast Region released a set of protocols for continuous water temperature monitoring for their timber harvesting water quality waiver program. SDSF's methods are consistent with these protocols. An analysis of the number of days with temperature readings exceeding 21.1°C and 14.4°C provides valuable information for fisheries management in SDSF. No temperatures greater than 21.1 °C were recorded in 2009 - 2011. In 2008, Longridge Crossing had two days over 21.1°C. Temperatures over 21.1°C were recorded on 11 days in 2007. Detailed reports and temperature data analysis are compiled annually. The graph below (Figure 7) is a summary of the high temperatures recorded from 2005 to 2012. In 2011, the Amaya Creek logger was defective and in 2012 the Southwest Boundary logger was defective.

AQUATIC INVERTEBRATES

Aquatic invertebrates are organisms that lack an internal skeleton and live in water for at least part of their life cycle. They include insects (e.g., mayflies, stoneflies, caddisflies), crustaceans (e.g., crayfish), mollusks (e.g., snails), and freshwater earthworms. An important component of aquatic ecosystems, aquatic invertebrates are an essential part of the food web. They typically act as indicator species of fishery habitat quality and water pollution.

When tree litter falls into the water, microorganisms such as bacteria and fungi colonize and decompose the coarse particulate organic matter, creating a much more nutritious material. This substance, algae, and other invertebrates become food for invertebrates, which in turn become food for fish. In this way, aquatic invertebrates provide a link in the food chain between decomposers and fish, a connection crucial to fish survival.

Figure 7. Water Temperature Data for SDSF.



In fresh water such as the East Branch, juvenile steelhead feed primarily on immature, aquatic stages of insects, but will also feed on adult terrestrial insects (Barnhart, 1986). Steelhead prefer the larvae of mayflies, true flies (e.g., midges), and caddisflies. They are often opportunistic, however, and may feed on any available insect. Adult steelhead typically do not eat during migration and spawning, instead utilizing energy from fat accumulated while living in the ocean.

Since they function as food, aquatic invertebrates, particularly insects, are key indicators of good fish habitat. In most cases, abundant and diverse species of aquatic invertebrates signify an adequate food supply, increasing the ability of the stream to support larger and healthier populations of fish. Similarly, invertebrates need adequate amounts of leaf litter falling into the stream (therefore, adequate riparian vegetation) and appropriate conditions to support the bacteria and fungi which convert the litter to a usable form. Because of their specific roles and different, species-specific habitat requirements, invertebrates are useful indicators of stream conditions and changes. Through inventory and monitoring the composition of aquatic invertebrate communities, the health of aquatic ecosystems can be carefully evaluated. Regular inventory and monitoring has not occurred at SDSF. Some aquatic invertebrate data have been collected through various CDFW studies, but the data have not been made available. Future plans for studies of aquatic invertebrates are included in the monitoring plan by NMFS for the Large Woody Debris and Habitat Complexity Project. This work would monitor the habitat, including aquatic invertebrates, for the restored sites as well as control sites along the East Branch Soquel Creek.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Comments and concerns regarding the aquatic resources of Soquel Creek were numerous and often passionate. Given the condition of California's CCC coho salmon and steelhead resource and the significance of the Soquel Creek watershed, one would expect a great concern for its future. The comments, concerns, and suggestions received as a result of this concern are represented by two major categories.

The first and probably most important concern involves the maintenance and enhancement of fisheries and other aquatic resources within SDSF. Individuals have stated that, at the very least, SDSF should maintain the resources as they currently exist. Additionally, suggestions for improving the habitat and, therefore, the steelhead fishery touched on in-stream improvements and careful logging methods. SDSF will assuredly work to enhance all wildlife resources and habitats, including those of steelhead. Moreover, SDSF is already working with NMFS and intends to continue to be an active partner in recovery of both steelhead and coho in the Soquel Creek Watershed.

The second category of concern is really an extension of the first. Comments related that, in order to maintain and enhance SDSF's aquatic resources, inventories and monitoring of these resources must be performed. Specifically, commentators expressed that biologists familiar with the Soquel Creek watershed should set up and maintain a monitoring system which tracks the effects of SDSF's management activities on the resource. Strategies to ensure

maintenance and enhancement of aquatic organisms could in turn be developed from the results of inventory and monitoring activities. SDSF plans to conduct the activities mentioned above, as well as research and habitat improvement projects, in order to satisfy these concerns (see Planned Actions below).

Consistent with the Anadromous Salmonid Protection Forest Practice Rules approved in 2009, conifers will be planted in riparian zones, in areas where none exist, to promote long-term recruitment of large instream woody debris. These conifers will eventually provide small intertwined pieces of debris, such as branches and twigs, that act as collectors of leaf litter and provide more food for invertebrates.

MANAGEMENT GUIDELINES

1. Protect stream channels, streambanks, and riparian zones during all management activities. Late-succession management areas (see the Timber Management Chapter) have been established along all fish-bearing streams and receive specialized management designed to enhance the riparian zone. This will ensure protection of stream integrity, including the channel, bank, and vegetation as well as fisheries resources.
2. Increase the fisheries potential by improving the spawning and rearing conditions of the East Branch of Soquel Creek within SDSF by implementing projects to increase winter high flow refuge, summer rearing, and increase general habitat complexity as well as projects that reduce production of fine sediments.
3. Demonstrate that other forest management activities are compatible with the maintenance of healthy fisheries populations and habitats through educational programs and tours of harvested areas and stream enhancement projects.
4. Help residents of the East Branch watershed learn about the components of a healthy watershed and the importance of stewardship as it relates to LWD management, management of rural roads, and water conservation. Educational programs, either formal or informal, will focus on stream health, riparian zones, and the impacts of human use. Once established, the Forestry Education Center (refer to the Demonstration and Education Chapter) will likely be the setting for these programs, allowing residents and other interested individuals to learn about the significance of aquatic ecosystems.
5. Continue to work with NMFS and CDFW on means and methods to help provide incentives and to facilitate private landowners playing an active role in recovery of threatened and endangered salmonids (e.g., assistance with permitting, including LWD projects as part of THPs, etc.)

PLANNED ACTIONS

1. Implement mitigations through the timber harvesting plan process that benefit anadromous fish.
2. Monitor the Large Woody Debris and Habitat Complexity Project in cooperation with the California Department of Fish and Wildlife, National Marine Fisheries Service, California Geological Survey, Resource Conservation District of Santa Cruz County, SWFSC and local qualified biologists.
3. Continue to support aquatic ecosystem research opportunities in the East Branch and its tributaries.
4. Coordinate with the County of Santa Cruz, the Resource Conservation District of Santa Cruz County, National Marine Fisheries Service, California Department of Fish and Wildlife, and other groups to complete habitat restoration and maintenance projects, including installing large wood in Soquel Creek and control of invasive plants. Private groups such as the Monterey Bay Salmon and Trout Project or crews from the Ben Lomond Conservation Camp may help with projects such as channel stabilization, channel complexity and pool creation, retention of existing instream cover via LWD, riparian vegetation maintenance and enhancement, and reconnection of floodplains and off-channel habitat.
5. Restoration and enhancement projects for the Forest will be prioritized based on recommendations in the Soquel Creek Watershed Assessment and Enhancement Project Plan (Alley, 2003), the DFG Recovery Strategy for Coho (DFG, 2004), the National Marine Fisheries Service (NMFS) in their Final Recovery Plan for the Evolutionarily Significant Unit of Central California Coast Coho Salmon (NMFS, 2012), and the forthcoming NMFS Recovery Plan for the Distinct Population Segment of Central California Steelhead. In addition, the Forest will continue dialogue with biologists from NMFS as well as NOAA's Southwest Fisheries Science Center, CDFW and private biologists to identify critical opportunities for restoration and enhancement of fisheries resources. Projects will be implemented over time as funding, equipment, and/or personnel become available.
6. Continue fish sampling in Amaya Creek and the East Branch of Soquel Creek in cooperation with NOAA's Southwest Fisheries Science Center. Electrofishing will be the primary method but other procedures such as underwater observation may be incorporated. Fish will be evaluated for species, size, health, and location.
7. Continue to monitor in-stream temperatures in the East Branch of Soquel Creek and Amaya Creek at the seven sites that have been established.
8. Build on existing funding from the Integrated Watershed Restoration Program to develop baseline fisheries, macroinvertebrate, water quality, and habitat data prior to implementation of fisheries restoration projects to assess effectiveness. Aquatic habitat surveys will be conducted in accordance with CDFW methodologies as

funding allows.

9. Monitor projects that are implemented in accordance with the California Department of Fish and Wildlife methodologies.
10. Explore opportunities for working with the Monterey Bay Salmon and Trout Project, NMFS, NOAA, and CDFW on the potential to add sites within SDSF to the introduction list for coho from the broodstock hatchery program.
11. Conduct ongoing patrols with the assistance of the California Department of Fish and Wildlife to enforce prohibitions on fish poaching and harassment. Incidents will be recorded, compiled, and evaluated annually to determine significant adverse effects on SDSF fisheries.

CHAPTER 7: WATERSHED ASSESSMENT

PHYSIOGRAPHY OF THE WATERSHED

SDSF is located almost entirely within the drainage of the East Branch of Soquel Creek, its landbase covering approximately 21 percent of the basin. For the most part, the Forest is underlain by fine grained sedimentary rocks that are highly weathered and easily eroded. The headwater channel of the East Branch follows the San Andreas Fault Rift Zone and is heavily disturbed from landslides triggered by earthquake activity. Hillslopes throughout the Forest are commonly steep and prone to large, deep-seated landslides. Even greater instability occurs in active inner gorge zones near stream channels. Mass erosion events (e.g., large landslides) comprise the major component of sediment sources in the watershed.

As stated in the Property Description Chapter, the Soquel Creek watershed (see Figure 8 below) has a Mediterranean climate with cool, wet winters and warm, dry summers. Mean annual precipitation in the East Branch watershed varies from 30 to 46 inches and takes place primarily between November and April. Discharge records from the USGS stream gauging station located immediately upstream from the bridge in the town of Soquel show that the average discharge is 1.05 cubic feet per second per square mile. High intensity, long duration winter storms, however, produce extreme levels of runoff. Major floods occurred during December, 1955 and January, 1982, producing log jams and flooding in the town of Soquel (Lassette and Kondolf, 2003; see Figure 9). Singer and Swanson (1983) state that Soquel has a chance of being flooded once every seven to ten years.

Since 1996 streamflow has been measured at a gauge on the East Branch of Soquel Creek just below the Olive Springs Quarry. This gauge and the data are maintained by the Soquel Creek Water District. The watershed area above the gauge is 13.9 square miles and includes essentially the entire SDSF area (a very small percentage of SDSF drains to Hester Creek). Data from this gauge have been coupled with data from a permanent rainfall gauge located just outside the Forest on Longridge Road and are represented on Figure 10 below.

BENEFICIAL USES

The two main beneficial uses of the East Branch of Soquel Creek are cold water fisheries and water supplies for various purposes. The East Branch supports spawning and summer rearing habitat for coho salmon and steelhead trout, but the number of returning fish has declined significantly in the past 40 years (refer to the Fisheries Chapter). For example, in 1982 steelhead populations were only one-third to one-quarter of what they were thought to have been in the 1960s (Singer and Swanson, 1983). Coho salmon were thought to be extirpated from the Soquel Creek basin since 1968 (SCCRCD 2003, NMFS 2012), but 170 young-of-the-year fish were documented during surveys conducted in a stream reach below SDSF in August 2008. The major factors limiting anadromous fish populations are believed to be low summer flows, and limited rearing habitat due to siltation of pools and removal of woody debris. Large wood loading, floodplain connectivity, and estuary function were listed

Figure 8. Soquel Creek Watershed Map.





Figure 9. Annual Peak Streamflow for Soquel Creek.




as poor for the Soquel Creek watershed in the Central California Coast (CCC) coho salmon recovery plan (NMFS, 2012). Soquel Creek and Aptos Creek are considered to be the southernmost extent of coho salmon in California.

The other primary beneficial use in the main stem of Soquel Creek is water supply. There are eight dams in the Soquel Creek watershed that impede or block anadromous salmonid migration (NMFS, 2012). Additionally, numerous small diversions exist that provide water for domestic, agricultural, and industrial purposes. Two permitted domestic water supply systems exist in the main Soquel Creek watershed that utilizes surface flow, but there are no permitted systems utilizing surface water in the East Branch basin. The East Branch Soquel Creek watershed assessment conducted in 1993 reported that there were six surface water diversions in the East Branch basin (Cafferata and Poole, 1993). The largest diversion in the East Branch is operated by the Olive Springs Quarry.

Diversions and overdrafting of groundwater in low discharge summer months have dewatered portions of the East Branch stream channel during drought years. This is likely to be a critical limiting factor for the fisheries resource in the lower part of the basin. Soquel Creek is a fully adjudicated stream under Decree No. 57081. After complaints by the city of Capitola in 1988, the State Water Resources Control Board concluded that a watermaster was needed to effectively regulate water use under low flow conditions, particularly during

Figure 10. – Rainfall and Streamflow for East Branch of Soquel Creek.



drought years, due to the complicated interrelationships of the water rights on Soquel Creek (Cafferata and Poole, 1993). To date, however, no watermaster has been appointed (Alley, 2004).

LAND USE HISTORY

The major types of land use in the East Branch watershed are timber management, recreation on public land, vineyards, and residential development. The East Branch watershed's long history of timber harvesting began with clearcutting in the 1870s and continued into the 1940s. Timber was originally removed by oxen (1870-1895), then by steam donkey (1895-1930), and finally by crawler tractor. Following World War II, selective harvesting replaced clearcutting, continuing to the present day. Currently, nearly all of the old-growth timber has been harvested except for minor reserved groves and widely scattered suppressed trees. Second-growth harvesting has taken place on approximately 961 acres over the past 10 years (2003-2013) within the 9068 acre Soquel Creek Watershed. In recent years, cable yarding has been used along with tractor logging to selectively harvest timber in the basin. Approximately twenty percent of the Soquel Basin has never been harvested as it is in chaparral.

Recreation in the form of hiking and mountain biking takes place primarily in The Forest of Nisene Marks State Park and SDSF (SDSF also allows horseback riding). Additionally, adjacent property owners often allow friends and neighbors to utilize their lands as an access route to the park or SDSF. The Olive Springs Quarry, located near the East Branch channel, has produced sand and gravel from granitic rock exposed along the Zayante Fault since the 1950s. Residential development has occurred in the chaparral communities and has steadily expanded over the past 40 years. Over the whole Soquel Creek watershed, housing development is rated as moderate to high, with approximately 7,000 housing units present in the basin. Residential and commercial development is considered a very high threat to coho salmon (NMFS, 2012).

CUMULATIVE WATERSHED EFFECTS ASSESSMENT

Cumulative watershed effects (CWE) can be defined as the physical and biological impacts that result from multiple land use disturbances over space and time. These impacts occur within and away from the locations of actual land use and are transmitted through the fluvial system. When considering CWE, it is appropriate to estimate how current and future projects, when combined with impacts from past activities, will influence beneficial uses present in the basin under review. Techniques to determine whether CWE are significantly adversely impacting beneficial uses have been developed, but are generally considered to be inadequate for varying reasons. Existing CWE assessment approaches mostly range from checklists or indices that are subjective but inexpensive and simple, to complex physically based models that have large data needs and are difficult to apply (Litschert, 2009).

Watershed analyses and assessments are often used to evaluate cumulative watershed effects.

For example, Berg et al. (1996) found that watershed analysis was the most suitable approach for assessing cumulative watershed effects in the Sierra Nevada. While a formal watershed analysis has yet to be completed for the Soquel Creek watershed, several watershed assessments and studies have been completed over the past 30 years that contribute a considerable amount of information regarding cumulative watershed effects. These studies include: Singer and Swanson (1983), Cafferata and Poole (1993), Santa Cruz County Resource Conservation District (2003), Alley and Associates (2003, 2004), Balance Hydrologics (2003), Greening Associates (2003), Pacific Watershed Associates (2003), and Lassetre and Kondolf (2003). Cafferata and Poole's rapid landscape-level watershed assessment for the East Branch of Soquel Creek was conducted in 1992 and 1993. It is described below in considerable detail. Since specific harvest units had not been defined, the entire Forest served as the project area for this assessment. CWE assessment guidelines suggest that assessment areas must be large enough to detect past impacts and small enough to determine what the impacts of the proposed projects will be on the area. This assessment completed for the East Branch watershed evaluated the current condition of stream channels in the basin and determined how they would likely be altered with future timber operations.

The watershed assessment for SDSF was broken down into two main tasks: a stream channel inventory and a hillslope erosion/sedimentation evaluation. The stream channel inventory was completed in 1992 and illustrated the current condition of the channel network within the assessment area. Channel stability was rated with the U.S. Forest Service's Pfankuch Method, while channel condition was evaluated using guidelines for assessment of cumulative impacts currently found in Technical Rule Addendum No. 2 of the California Forest Practice Rules (CAL FIRE, 2014). Due to limited personnel and time constraints, stream channels within SDSF were analyzed in greater detail than channels located in other areas of the basin (Poole, 1992).

The hillslope component of the CWE evaluation was completed in the summer of 1993. Quantitative estimates of erosion risk and erosion volumes associated with anticipated activities (e.g., timber harvesting and road building) were made based on the results of the Critical Sites Erosion Study (CSES; Lewis and Rice, 1989; Rice and Lewis, 1991). A selected number of random sample locations were installed on existing, planned, and abandoned roads, as well as past harvest areas. The resulting data were input into equations to estimate the risk of generating critical, or large, erosion sites (i.e., a large erosion event producing more than 100 cubic yards per acre). These equations are based on measurements made for slope, amount of hillslope or road curvature, soil color, and rock hardness.

On-site measurements were made to assess other sources of erosion not addressed by the CSES. Estimates were then made of the amount of sediment that could result from the approximated erosion. The significance of the estimated sedimentation was studied by comparing it with expectations of sediment yield for Soquel Creek. Since long-term sediment data from the Soquel Creek basin were not available, information from the San Lorenzo River, a similar, neighboring basin, was used. The methodology utilized for the hillslope erosion and sedimentation analysis was designed by Rice (1993) and was used in several parts of the state in the 1990's.

Locations in the East Branch assessment area that were found to have poor stability and channel conditions were Amaya Creek, Hinckley Creek, and the East Branch between Ashbury Gulch and the upper SDSF boundary. The reaches with the best ratings for stability (high-fair) and channel conditions were the upper part of the East Branch above SDSF and the main stem of Soquel Creek below the junction with the West Branch. The reaches of the East Branch between Ashbury Gulch and the junction with the West Branch were reported as having intermediate stability and channel conditions.

Summarizing the results for the 10 miles of the East Branch surveyed, fair channel stability was the most frequent finding (10% good, 67% fair, and 23% poor). This compares reasonably well to the fair/poor relative overall rating for sediment contribution assigned to the East Branch by Singer and Swanson (1983). The exception was the highly impacted reach along the San Andreas Rift Zone. Additionally, the major tributaries were severely degraded by heavy landsliding activity. Very large quantities of sediment are stored in these headwater tributaries, as well as in wide, vegetated terraces along the lower reaches of the East Branch. Observation of the East Branch channel revealed that riffle stretches generally had cobble and gravel sized particles protecting the stream bottom surface. Inspection of stable pools below Ashbury Gulch, however, showed that this habitat type contained significant percentages of fine sediment.

Large woody debris was not found to be blocking anadromous fish passage in the East Branch, but was limiting habitat use in the major tributaries. Large wood was generally lacking along most of the East Branch of Soquel Creek below Ashbury Falls, except where it had accumulated in a few fairly stable locations. Large woody debris is valuable because of the pool habitat for young salmonids that forms around it in gravel dominated stream systems.

Results of the hillslope evaluation indicate that the risk of generating critical erosion sites is not significantly different from the average for the rain-dominated portions of the northern Coast Ranges in California (Cafferata and Poole, 1993). Very large amounts of existing erosion were measured on a few of the plots, however, and existing erosion was found to be considerably higher than the amount estimated from the critical site equations. For example, one of the road plots had a culvert that had carved an exceptionally large gully below its outlet, while another had large amounts of cut and fill slope erosion 50 feet from a Class II stream (as defined by the California Forest Practice Rules). Therefore, the potential for producing large erosion events is clearly evident in this terrain, even though the critical site equations did not generate exceptionally large estimates.

The terrain slope and distance to a Class II stream were used as an index of the proportion of erosion which would become sediment. The average sediment delivery was found to be much higher than is commonly assumed in forested watersheds. This is due to very high delivery from relatively few plots with extensive amounts of existing erosion. Long-term sediment data do not exist for Soquel Creek at the USGS gaging station; the nearest station with long-term discharge records and a reasonable sediment record is found on the San Lorenzo River at Big Trees State Park. A statistical analysis indicated that peak storm discharges on Soquel Creek were significantly related to those of the San Lorenzo River.

The peak storm discharge/annual sediment yield relationship generated for the San Lorenzo River was then used to estimate the annual sediment yield at Soquel Creek.

Extreme natural variability in sediment yield for Soquel Creek was found for 1952 to 1990, with the range spanning several orders of magnitude. The estimate of mean annual sediment yield based on the stream discharge record for Soquel Creek (60,392 yd³ or 2.35 yd³/ac/yr) was somewhat higher than that calculated based on hillslope erosion plot measurements. Due to the assumptions involved in making these estimates, however, it is unlikely that these estimated sediment yields are significantly different. Both estimates of sediment discharge for Soquel Creek are high for forested areas and illustrate the erosive nature of the basin. Cafferata and Poole (1993) concluded that it was unlikely, however, that the limited timber harvesting and rebuilding of abandoned roads planned for SDSF would significantly elevate the erosion rate and have a significant adverse impact on the aquatic system. The anticipated sediment production from SDSF's timber production was found to be small compared to the range of variability in sediment flux observed in this basin.

More recent watershed assessment work in the Soquel Creek watershed can be compared to the results of the Cafferata and Poole (1993) rapid assessment. For example, Lassette and Kondolf's (2003) much more detailed large wood study included the lower part of East Branch of Soquel Creek. They reported an average large wood loading for this reach of 0.005 m³/m², which they stated lies at the low end of the range of observed values for North American streams (e.g., approximately 36 times lower than for old growth coast redwood forests in Humboldt County). Similarly, Alley and Associates (2003) reported that large wood was extremely scarce in Soquel Creek compared to other coastal streams recently surveyed.

Large wood loading in the East Branch of Soquel Creek is low due to extensive log removal efforts by Santa Cruz County from the 1950s to the 1990s. The Santa Cruz County RCD watershed assessment (SCCRCD, 2003) concluded that the scarcity of large wood limits juvenile salmonid production throughout the Soquel Creek watershed. Similar to other studies, their assessment states that large wood is scarce in the middle part of the East Branch and recruitment is low for this channel reach.

Balance Hydrologics (2003) reported that Soquel Creek has experienced prolonged periods (up to 25 to 30 years) of disturbed watershed conditions over the past 150 years. These conditions, they found, were caused by both natural processes and anthropogenic activities, with the later likely resulting in an increase in the natural, base rate of sediment production in the watershed, although small relative to natural rates. Landslides along Highland Way as well as recent wildfires (Summit Fire and Loma Fire) have also contributed sediment to Soquel Creek.

The Santa Cruz County Resource Conservation District (2003) watershed assessment project concluded that overall lateral channel stability of Soquel Creek was relatively high during the previous 54 years with minor adjustment at several points of meander along the East Branch and mainstem. Major sediment production from natural sources was found to occur in both the East and West Branches and is most pronounced following earthquakes, large magnitude

floods or forest fires, and is usually related to landsliding. However, SDSF will continue to remediate existing sediment sources both during timber harvests and after, working towards a net decrease in sediment inputs from timber harvest operations.

The SCCRCD (2003) assessment states that several factors appear to limit distribution and abundance of steelhead. These factors include passage impediments, poor spawning habitat quality (high proportion of fine sediment, number of constricting, steep riffles below spawning glides), low spring and summer baseflows, limited amount of escape cover (provided by instream wood, undercut banks, unembedded boulders, and water depth), elevated water temperature, and limited water depth. Throughout the watershed, low baseflows and sedimentation limit the amount and quality of rearing habitat.

Pacific Watershed Associates (2003) inventoried the SDSF road network (18.2 miles) and found 82 sites with significant sediment delivery potential. Fifty-seven of these sites were at stream crossings. Three crossing sites were identified as having high treatment immediacy, with a potential sediment delivery of approximately 1,631 yds³. Sites requiring significant analysis, design, and heavy equipment investment, are upgraded and/or mitigated, as CEQA projects are planned and implemented across SDSF. For example, one high priority culvert replacement as well as three other identified sites, were mitigated through implementation of the Rim THP and accompanying CDFW streambed alteration permit in 2011. Additional rocking of the main roads in SDSF has also been implemented annually. Upgrades and mitigations along Longridge Road and Hihn's Mill Road were implemented as part of the Fern Gulch timber sale.

Recent impacts in the Soquel Creek watershed have occurred that have not been considered in past watershed assessments. These include the 2008 Summit Fire, which burned approximately 4,270 acres in portions of the Soquel Creek, Corralitos Creeks, Brown's Creek, and Uvas Creek watersheds (SEAT, 2008). Soquel Creek suffered the greatest amount of high burn severity, with 382 acres. In total, 1130 acres burned in the Soquel Creek watershed (34% high burn intensity, 43% moderate, 20% low, and 3% unchanged.)

Additionally, a large landslide event delivered an extensive amount of fine sediment into the East Branch of Soquel Creek in 2011. Highland Way was closed by a re-activated landslide on approximately March 26, 2011. The slide contributed a substantial amount of soil, rock, and organic debris into the East Branch of Soquel Creek that has had short and long term impacts.

CONCLUSIONS

In summary, it can be stated that the East Branch of Soquel Creek watershed is a highly sensitive basin. This is due to its inherent natural hillslope instability related to seismic forces, steep slopes, weak bedrock, and frequent high intensity rainfall events. Impacts from past management activities, prior to implementation of the modern California Forest Practice Rules in 1975, have contributed greatly to degraded conditions in the basin, as has residential development. In spite of these problems, the East Branch below Ashbury Gulch has not been overwhelmed with sediment, and it is clear from abundant observations of steelhead redds

(gravel beds where female fish lay eggs) and young fish that spawning and rearing habitat remains.

The East Branch system is stressed by fine sediment moving downstream from tributaries, partially due to its close proximity to the San Andreas Fault Zone. This material is filling pools required for rearing habitat for listed anadromous salmonids. Lack of sufficient volumes of large wood and overdrafting of water in low flow summer months compounds this problem. As a result, the aquatic environment is sensitive to further degradation and timber operations must be carried out with extra caution.

Due to the sensitivity of the hillslopes and current channel conditions, special considerations are needed when planning forest management activities in the East Branch. Appropriate mitigations must continue to be utilized and remedial improvements implemented to repair existing problem areas, such as those suggested in the PWA (2003) report. If these considerations are employed, future timber sales can occur without significant adverse impacts to the beneficial uses of the basin.

MONITORING AND ENHANCEMENT

Ongoing management of SDSF will involve maintaining proper drainage along roads and trails by repairing culverts, water bars, and other drainage structures to reduce or prevent soil erosion and stream sedimentation. SDSF is required to monitor all timber operations (including all harvesting areas and new roads, skid trails, and landings) annually for 5-7 years following completion of operations. Increased monitoring occurs the first winter after operations above what is required by regulations. An active watershed remediation program will continue to be used that includes monitoring watershed conditions and implementing enhancement projects. Monitoring has included a forestwide inventory of stored sediments and active landslides, and mapping mass wasting hazards and surface erosion potential. Sediment risk-reduction projects have been rated for cost-effectiveness (PWA, 2003) and will be implemented in conjunction with timber operations according to their priority as available funding permits.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Public comments regarding the East Branch watershed have focused on its distinction and sensitivity. Individuals are concerned that forest management activities will result in general watershed degradation, including sedimentation of streams and excessive hillslope erosion. This is particularly important to them in regard to steelhead numbers and habitat, and possible impacts upon their land during high intensity rainfall. As stated in this and other chapters, careful pre-project evaluation and measures to ensure the maintenance of watershed integrity are a priority for SDSF. The CWE assessment work discussed above and other studies assist the Forest staff to understand and manage for the sensitive nature of the basin.

MANAGEMENT GUIDELINES

1. Design streamside management zones that properly address the inherently unstable nature of the East Branch watershed in SDSF. This includes extending zone widths beyond the standards set by the California Forest Practice Rules, for Class I fish-bearing watercourses, as is appropriate in sensitive areas. These zones must provide for the long-term recruitment of large wood, protection of the stream channels and banks, stream shading, sediment filtration, nutrient input, microclimate control, floodplain function, and prevention of significant ground disturbance.
2. Place heavy emphasis on road design and maintenance, since roads generally produce the largest percentage of management-related sediment in forested watersheds. New seasonal and temporary roads should be outsloped to avoid concentrating water that could trigger landslides or transport sediment directly into flowing streams. Of foremost concern for new road construction is the avoidance of localized unstable areas. For mainline roads, road drainage structures and watercourse crossings must be adequately sized to ensure that the risk of failure is minimized. Roads no longer needed must be properly abandoned. Wet weather use for roads impacting flowing streams should not occur and an active winter maintenance program is needed to ensure that drainage structures are adequately functioning. (Refer to the Roads and Other Improvements Chapter for more information.)
3. No tractor operations will be permitted on slopes which average more than 35 percent without site-specific evaluations of slope stability and erosion potential. This will depend on the ability to mitigate such operations to levels of insignificance.

PLANNED ACTIONS

1. Continue to record data on all timber harvesting operations done in the Forest that could influence soil and water resources. Link THP road mitigation sites, information on road construction and harvesting, and monitoring results through our GIS database.
2. Continue to implement the treatment priorities for high and moderate risk inventoried sediment sources in the Soquel Demonstration State Forest watershed assessment area developed by PWA (2003).
3. Evaluate the performance of each previously-implemented remediation project to determine the success in reducing the risk of large-scale sedimentation. Redesign and modify any project not meeting its intended objective (see to Appendix C).

CHAPTER 8: DEMONSTRATION AND EDUCATION

INTRODUCTION

As stated in the Timber Management Chapter, a goal of SDSF is to demonstrate sustained-yield timber production with innovative forest management practices within the context of local community protection, and subject to the limitation of commercial timber harvesting provided in the legislation (PRC 4660-4664). The intent of this goal is to encourage conscientious forestry practices on private lands and demonstrate these practices to the surrounding urban populations. This can lead to improved attitudes toward our natural resources and forestry in general, enhancing responsible stewardship of our forest lands.

Another goal is to show Forest visitors that timber management, forestry education, public recreation, and environmental protection are interrelated and compatible. This will be accomplished through combinations of these programs whenever possible. SDSF's location is well suited for the development of forestry education programs because it is close to schools in both the Monterey Bay and south San Francisco Bay areas. This proximity is ideal for groups to learn the importance of forest ecology and management. Similarly, it offers local landowners and the general public an opportunity to view the protection, management, and utilization of renewable natural resources. Field trips involve public motor vehicle use.

The Demonstration and Education Programs of SDSF rely on interpretation, volunteer participation, and the future creation of a Forestry Education Center. Each of these elements, as well as the Demonstration and Education Programs themselves, are outlined in the following paragraphs, and further described in the Education Study (Blazej, 1997).

DEMONSTRATION PROGRAM

OVERVIEW

The Demonstration Program is a major focus of SDSF for a number of reasons. Most importantly, it provides an opportunity for the general public and educational groups to observe an active working forest. Forestry demonstration can assist the public in understanding forest management and its role in resource conservation and local economics. Additionally, it reminds us of the many wood products we use daily and the importance of keeping forests healthy and productive. Knowledge such as this can help change the negative public image generally associated with forest management practices. Also, the demonstration of various forest activities can potentially benefit small private landowners in the management, protection, and enhancement of their forest lands.

The opportunity for demonstration projects has increased as SDSF has become better established. The Forest staff remains aware of the demonstration potential at the onset of new activities. Demonstration projects may be short term, with several topics addressed each

year, or long term, extending over many years.

The primary consideration of the Demonstration Program is to enhance the public's understanding and awareness of forest management principles and techniques consistent with environmental protection. Throughout the process of establishing projects, a strong emphasis will be placed on environmental protection. The following are demonstration opportunities at SDSF:

Silvicultural Systems	Erosion Control
Tractor/Cable Harvesting Operations	Hardwood Management
Disease and Insect Management	Growth and Yield
Recreation Management	Riparian Management
Reforestation Methods	Cumulative Watershed Effects
Fisheries Protection/Enhancement	Fuelwood Management
Road Construction/Maintenance	Prescribed Burning
Old-Growth Redwood Protection	Habitat Enhancement(Aquatic and Terrestrial)
Watershed Restoration	Fire Protection
Exotic Species Control	Rare Plant Management

Part of the demonstration that is done at SDSF involves public field trips of active timber harvests. Four to five field trips were held each year for the 1995 Longridge harvest, the 1997 Amaya harvest and the 2012 Fern Gulch harvest. The field trips were attended by 30 to 40 people per trip. Each timber harvest has also held a specific field trip for the SDSF Advisory Committee to provide an opportunity to review current operations. During the tours over the years the advisory committee, media, and public have had the opportunity to observe operations including rubber-tired skidding, tree falling, horse skidding, cable yarding, a portable mill demonstration, the Badger Spring's old growth grove, the Olive Springs Quarry operation, log truck loading, erosion control techniques, and watercourse crossing installations. Topics discussed have included the history and goals of SDSF, watershed protection and enhancement, and timber management. Questionnaires were filled out by field trip participants to get feedback about their experience and recommendations for future field trips. Almost without exception, the field trip participants enjoyed the opportunity and stated that they would be interested to come back in the future. These field trips will continue to be a part of future harvests at SDSF.

PLANNED ACTIONS

1. Demonstrate forest management practices and ecosystem enhancement techniques. To accomplish this, incorporate an identifiable demonstration feature in timber sale planning and implementation, recreation designs and development, and other forest management activities, as appropriate.
2. Develop and implement outreach programs to contact the general public, school groups, and private landowners for demonstrational opportunities. Encourage visits and tours by interested public groups, individuals, schools, and professional organizations.

3. Establish safe and efficient methods of displaying information from demonstration projects and conducting tours of these areas. Displays should be made available and tours held at times that encourage Forest user group's attendance and participation (e.g., summer weekends, evenings).

FORESTRY EDUCATION

OVERVIEW

SDSF has proven to be a valuable and attractive venue for forestry education, just as the Recreation Study (McNally and Hester, 1993) indicated it would. SDSF provides informative and fun educational opportunities and as many hands-on experiences as possible. The program is applied in conjunction with demonstration objectives and occurs in many different forms. Forestry education introduces various target groups to progressive forest management practices, resource protection, logging history, forest ecology, and research. This is accomplished through literature provided, indoor presentations, and outdoor programs.

Instructional organizations can learn about SDSF's resources by sponsoring classroom presentations or bringing groups to the Forest. School programs may represent a theme, such as Arbor Day or Earth Day, and include activities both at school and SDSF. Classroom and outdoor presentations might also be subject oriented, with topics such as watershed management, forest ecology, soil biology, fisheries habitat restoration, or wildlife found in the Forest.

Educator training programs, like the one at Elkhorn Slough Reserve or the Forestry Institute for Teachers, provide a means for teachers and their students to experience forestry education without relying on Forest staff. After completing training, educators may bring their students to the Forest for non-staffed instructional field trips. This removes the time burden from Forest staff members while allowing instructors to absorb new information and reinforce their own training.

SDSF staff has assisted with Forest Conservation Days, the web of life based educational program for fifth graders, since its inception in 1992. SDSF staff is also involved with the California Forestry Challenge, which is a competitive event for high school students in technical forestry and current forestry issues. Since SDSF cannot currently provide a suitable venue for hosting these events on the Forest due to lack of staff, infrastructure and services on site, SDSF staff participates in these programs that occur locally each year.

Information for the general public, professional organizations and small private landowners is disseminated through tours, newsletters, brochures, workshops, seminars, and the CAL FIRE State Forest website. Newsletters and brochures cover information such as current events, research projects, or compatible rural land uses. Workshops provide opportunities to learn about forest management techniques and how to solve problems in a group setting. Seminars focus on specific topics as well as the presentation of ideas, research results, and

group discussion.

PLANNED ACTIONS

1. Continue to provide educational information and programs related to forest management and ecosystem processes as described above and in the SDSF Education Study (Blazej, 1997).
2. Utilize various methods to distribute information about the forest resources of SDSF to different user groups. Methods to present facts about SDSF may include oral presentations, written information, interpretive facilities, and the internet.
3. Continue to encourage educational organizations, such as local public schools, to use SDSF for forestry education programs and field trips. Stimulate participation by offering quality educational experiences that are both informative and enjoyable. Maintain old and establish new working relationships with educators and their students.
4. Encourage teachers to attend and provide information about teacher training programs that will allow educators to lead their own forestry education programs. Offer workshops on topics such as forest ecology, wildlife habitat requirements, and watershed dynamics which apply directly to SDSF.

INTERPRETATION

OVERVIEW

The public can gather information about SDSF in a variety of ways while visiting the Forest. Interpretive facilities such as self-guided trails, information boards, and hands-on activities offer the Forest visitor an opportunity to learn the many features of SDSF. Since groups who utilize the Forest differ in their use of its many resources, information that applies to and interests many different individuals is essential. Interpretive facilities are a great way for the staff to meet the important goals of public demonstration and education without being relied upon for instruction.

INTERPRETIVE OPPORTUNITIES

Self-guided walking tours incorporate designated stops that are marked and indicated in a guide book. Stops on trails in SDSF might focus on unique sites and vegetation of the Forest or provide activities that facilitate learning. Interpretive trails will work well in SDSF because hiking is a common activity. The Forest provides many possibilities for educational trails, including a number of old logging trails and roads.

Information boards are effective tools that offer educational and operational information. Boards located at parking areas, popular recreation sites, trail heads, and along trails provide

information in the form of displays and brochures. Facts that relate to general Forest news are best suited to this type of interpretive facility, and include subjects like riparian protection, current logging practices, and forest ecology.

Hands-on activities with touchable items and sensory-oriented exercises for adults and children are another form of forestry education. These activities either stand alone or are coordinated with self-guided and staff-led tours. For example, a touchable tree round on a self-guided trail may help visitors understand how tree age is determined or how human history compares to that of a tree. Forestry tools such as clinometers or diameter tapes are used to show visitors and school groups how trees are measured. Areas containing hands-on activities are located near popular recreation sites, picnic areas, and parking areas.

Staff-lead tours will travel over expansive areas not covered by self-guided trails and contain stops that are not visibly marked. Guidebooks and hands-on activities may accompany tours, with staff members offering supplemental information.

On the internet – see <http://www.icogitate.com/~tree/SDSF.htm>.

PLANNED ACTIONS

1. Coordinate planning of interpretive facilities with all Forest activities, including recreational use, demonstration projects, resource protection, and timber harvesting. Install interpretive facilities near popular recreation sites, parking locations, and areas receiving silvicultural, enhancement, and restoration treatments.

VOLUNTEER PROGRAM

OVERVIEW

A volunteer program is used to assist the small Forest staff in several ways, including efforts to provide a complete and comprehensive forestry interpretive program. The major volunteer group associated with SDSF is the Stewards of Soquel Forest. The Stewards of Soquel Forest is a non-profit group that was established in 2000. The goals of the Stewards of Soquel Forest are to enhance the public's experience on the Forest, foster educational programs and assist with trail maintenance and infrastructure improvements. The Stewards have held education and demonstration events, raised funds for SDSF programs and facilities and have been instrumental in maintaining SDSF recreational trails (additional information is provided in the Recreation Chapter). SDSF's volunteers have an interest in both SDSF and natural resources in general, and enjoy sharing their experiences with others.

SDSF's volunteers can be available for public contact and interpretation in both the Forest and educational settings (such as classrooms). In the field, they benefit the Forest by providing visitors with information on current events, facilities, vegetation and wildlife, cultural history, and rules and regulations. Ideally, volunteers will have training in first aid and be able to radio for help during emergencies.

Volunteers also assist in research and monitoring programs, and maintenance and construction projects. The individual skills and talents of SDSF's volunteers, such as knowledge of local flora and fauna, leadership and interpretive skills, and experience working with people, are utilized to support the Forest. Through their work, volunteers benefit the operation of SDSF and acquire a better appreciation of state forests and natural resources in general.

PLANNED ACTIONS

1. Continue to develop volunteer programs which include docent recruitment. Volunteers will lead tours, patrol the Forest, and assist with education, research, and monitoring programs.
2. Continue to provide training to familiarize volunteers with the history of the state forest system, SDSF's history and objectives, rules and regulations, patrol procedures, and interpretive skills. General training will also include more specific information relating to SDSF's access routes, vegetation, wildlife, and research and monitoring objectives.

FORESTRY EDUCATION CENTER

OVERVIEW

If created, the Forestry Education Center (FEC) will be the focal point of SDSF. It could house the Forest headquarters, staff offices, an interpretive center for Forest visitors, and a meeting place for demonstration and educational tours. The FEC may also house a resource library, research laboratory, overnight accommodations, and an auditorium. The auditorium, useful for both administrative and educational purposes, could serve as a location for meetings, classes and workshops, seminars, informational slide shows, and videos.

The FEC would be used to present a range of information about SDSF's resources including maps, self-guided tour booklets, announcements of current events and management activities, signups for demonstration tours, and fire prevention information. Examples of interpretive exhibits and displays that could be made available through the FEC include:

Logging History/Equipment	Herbarium
Silvicultural Systems	Tree Physiology
Wood Products	Wildlife
Watershed Protection	Habitat Restoration
Fire Safe Programs	Fisheries

Plans have been approved for a replacement facility at the current SDSF office site in conjunction with the replacement of the CAL FIRE Soquel Forest Fire Station. The new SDSF office would replace the trailer acquired in 1992 and would include offices and a classroom. The project has not been funded and there is no estimate of when CAL FIRE

might eventually fund implementation.

PLANNED ACTIONS

1. Continue to progress on plans for the FEC based on expected use, cost, building size, and exhibit development. Select and acquire, if possible, an appropriate location for the FEC based on the above criteria as well as proximity to the State Forest and accessibility by the general public. For a detailed analysis of these topics, see the Forestry Education Center Study, Soquel Demonstration State Forest Proposed Program (Butler and Hester, 2001) and companion reports, Forestry Education Center Study, Soquel Demonstration State Forest Listening Report (Butler et al., 2001) and Forestry Education Center Study, Soquel Demonstration State Forest, An Analysis of California Environmental Education Centers (Butler et al., 2001).
2. Determine specifically how the FEC will be used. Conduct subsequent environmental documentation under CEQA. Design and build the facility, including grounds, so that expected use can be accommodated. Expansion and improvement of the Center in the future should also be taken into consideration during initial planning.
3. Plan and develop public education exhibits that meet forestry demonstration and educational objectives yet remain interesting and fun for all age groups. Encourage rotating presentations from local schools or special interest groups as well as permanent displays.
4. Seek alternative funding sources as contributions toward development and operation of the FEC. Supplemental funding from grants and private donations may be used for interpretive presentations, display development, and facility improvements.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Public comments and concerns relating to forestry demonstration and education have conveyed much enthusiasm. Respondents to the recreation study (see the Recreation Chapter) indicated excitement about the potential opportunities to view and learn about the workings of a forest such as SDSF. Additionally, comments about the FEC and demonstration goals express an interest in the facilities and methods through which demonstration and education will take place.

Forest neighbors and users have indicated curiosity in the future of the FEC, specifically its location and proximity to the Forest. Since the actual development of the FEC has yet to be determined, its location and accessibility remain undecided. Time and finances permitting, SDSF plans to carefully select and acquire a piece of property well-suited to everyone's needs.

CHAPTER 9: TIMBER MANAGEMENT

INTRODUCTION

The main purpose of SDSF's timber management and harvest program is to conduct demonstrations, education, and research in forest management. Sustained yield management is being demonstrated through the use of timber harvesting that ensures proper land stewardship, improves forest health, and protects and preserves SDSF as an intensively managed, multifaceted research forest. This is consistent with the legislative goals of PRC Sections 4660-4664.

As stated above, a major purpose of the SDSF timber management program is to demonstrate sustained yield management with examples of timber harvesting. In simple terms, sustained yield is the yield of commercial wood that an ownership can produce continuously at a given intensity of management consistent with required environmental protection and which is professionally planned to achieve, over time, a balance between growth and harvest. SDSF is used to demonstrate examples of timber harvesting under sustained yield management, while also sustaining or improving air, fish and wildlife, water resources, watersheds, aesthetic values, and recreation. Silvicultural methods and harvest techniques that can be applied under the California Forest Practice Rules for the Southern Subdistrict of the Coast Forest District are and will continue to be researched and utilized to demonstrate sustained yield management with an uneven-aged forest structure.

The harvesting of timber will comply with the discussion found in the Funding and Taxes section of the Administration Chapter. A secondary goal of harvesting timber is to generate revenue to cover maintenance, operation, and other costs of SDSF. This includes funds needed for research, inventory, monitoring, and rehabilitation projects of the various resources in SDSF. Revenues did not meet expenditures, even at the current minimum levels, during the initial nor the second ten-year period of this plan. It is anticipated that revenues and expenditures will converge in approximately the fifth decade of operations. Specific revenue projections are not made because of the inherent uncertainty of timber values and markets. Annual harvest levels may need to be adjusted from time to time to reflect physical conditions in SDSF, such as catastrophic events. If a significant drop in timber prices occurs, timber harvesting will not exceed the limits discussed in the Harvest Planning section of this chapter. CAL FIRE will attempt to maintain basic operation and maintenance services from the FRIF fund in line with other State Forests. For SDSF, this plan provides for harvest levels well below the level of growth. If future harvest continues at the current level, the forest will develop denser stands with bigger trees and more closed canopy over time.

HARVEST HISTORY

The first timber harvesting in SDSF occurred during the late 1800's when the Forest was part of land owned by F. A. Hihn (see the Property Description Chapter for a more detailed account of the history of ownership). Hihn logged old-growth redwood along portions of

Amaya Creek and his private road, which is now called Hihn's Mill Road. Upon his death in 1913, Hihn's heirs assumed ownership of his lands and continued limited old-growth redwood and tanoak removal.

In 1924, the property was sold to the Monterey Bay Redwood Company (MBRC). The MBRC harvested approximately 100 million board feet⁷ of old-growth redwood from 1926 to 1942. Between 1926 and 1934, logs were yarded to roads using ground lead cable systems powered by steam donkeys. Logs were taken to the millpond, located south of Olive Springs, on trailers towed by tractors. From 1934 to 1942, logs were yarded by tractor and transferred to the mill by log trucks.

The MBRC sold their property to Glenco Forest Products, later known as the CHY Company, in 1961. During their ownership of the SDSF property, CHY performed a small amount of selective timber harvesting between the Sulphur Springs area and the eastern boundary.

In 1979, the Pelican Timber Company purchased a portion of CHY's land, including what is now SDSF, and prepared extensive harvest plans. Pelicans's Timber Harvesting Plans were strongly opposed and, after court battles, expired before large amounts of timber were removed. Pelican did harvest a small amount, however, prior to state acquisition of the property. Approximately 15 acres of hardwoods in the Sulphur Springs area and 230 acres of second-growth redwood and hardwoods along the western boundary were selectively harvested under timber harvesting plans (THPs) 1-80-328 SCR, 1-81-25 SCR, and 1-83-56 SCR, operated on between 1979 and 1984.

Since the dedication of SDSF in 1990, four THPs have been completed (Figure 11). The first, Longridge THP (1-94-307 SCR) harvested in 1995, was a 64-acre single tree selection harvest and yielded 556,000 board feet. The Amaya Creek THP (1-98-027 SCR) harvested in 1998, was also a selection harvest, but included some small (1/4-1/2 acre) harvest groups. This THP harvested 138 acres and yielded 1.6 million board feet. Operations on the Rim THP (1-09-107 SCR) occurred over two years and were completed in 2012. The Rim THP was 158 acres and yielded 1.6 million board feet. The fourth plan, the Fern Gulch THP (1-09-096 SCR) was also completed in 2012 and yielded 2.2 million board feet on 201 acres. All four harvests combined represent about 6 million board feet over a period of 22 years, roughly equal to the Forest-wide growth in two and one-half years. The Comstock Mill THP (1-13-027 SCR) is planned for operations in 2014 on 228 acres for approximately 1.8 million board feet.


CONIFER VOLUME INVENTORY

There have been four known timber inventories completed for the SDSF property since the turn of the 20th century. The first inventory was completed in 1916 preceding significant harvest of the area. Timber cruisers from San Francisco were hired to cruise the Valencia-Hihn ownership comprising 7,071 acres including the SDSF acreage. The purpose of the

⁷ Conifer volume is measured in board feet; a board foot is a unit of measure one foot by one foot by one inch.

cruise was to estimate the standing timber volume and to establish the value prior to selling the land. When sold to the Monterey Redwood Company, 4,488 acres were estimated to contain 172 million board feet of redwood and Douglas-fir (Powell, n.d.). This yields an average per acre volume for this land, which includes SDSF property as well as portions of adjacent watersheds, of 38,300 board feet per acre. However, evidence from other similar tracts and logging history in the area suggest that this figure likely underestimates the amount of timber volume present in the redwood stands at that time.

Figure 11. SDSF Timber Harvesting Plans



The second inventory was completed in 1978 for Harwood Products, a potential buyer of the property, to ascertain timber growth and volume. Average stand age was approximated to be anywhere from 30 to 90 years old. The average volume per acre for the SDSF portion of the property was estimated at 13,600 board feet, 90 percent of which was redwood. The third inventory was conducted by SDSF staff in the summer of 1991. The goals of this variable-radius plot inventory were to establish current volume and basal area by species and to acquire information on species distribution. Based on harvest history, the average age of most Forest stands was estimated to be 65 years old. The average volume for SDSF was estimated to be 28,850 board feet per acre, with a forest-wide total conifer volume of approximately 77.3 million board feet.

The most recent inventory was completed in the spring of 2006. This was a variable-radius plot inventory with plots placed every five acres on a systematic grid. Data measured for all live trees included species, size and unique features such as goose pens, fire scars, and broken tops with potential wildlife habitat value. Table 6 contains the results for trees per acre, volume, and basal area of commercial conifers and hardwoods. The average age of most forest stands was estimated to be 80 years old based on harvest history. The average conifer volume for SDSF was 42,441 board feet per acre with a forest-wide total conifer volume of approximately 117.4 million board feet. This represents a 52 percent increase in Forest-wide volume in 15 years.

Table 6. SDSF 2006 Timber Inventory Summary.

	Redwood	Douglas-fir	Total Conifer	Hardwood	Total
Trees per acre	180	33	213	315	529
Basal Area⁸ per acre	178	35	212	116	328
Volume per acre (Board feet for Conifers) (Cubic Feet for HW >=7.0")	34,445	7,996	42,441	2,648	NA
Forest Wide Total Volume	95,261,398	22,114,678	117,376,076	7,323,309	NA

GROWTH

A conifer growth study was conducted at SDSF in 1993. This study produced per acre values for redwood and Douglas-fir on a forest-wide basis. The 1993 survey consisted of visits to 30 of the plots installed in the 1991 inventory to assess growth. Volume growth was estimated based on measurements of tree height and five- and ten-year radial increment growth. The results of this study indicated that forest-wide annual conifer growth averaged 972 board feet per acre with a standard error of 7.2 percent.

Growth based on the 2006 inventory data was estimated using the Forest and Stand Evaluation Environment (FORSEE) growth and yield model. Growth varied by vegetation type and stand density. The percent growth ranged from up to three percent in the lower density oak woodland stands to one percent in the highest density redwood stands. The average overall conifer growth for the Forest was estimated to be 975 board feet per acre per year, which represents 2.1 percent forest-wide growth per year. The growth across the Forest is estimated to be 2,615,360 board feet per year.

⁸ Basal area is the total cross sectional area of trees measured at four-and-one-half feet above ground and expressed as per unit area of land (typically square feet per acre). It is a measure of forest stocking or density.

In the future, a continuous forest inventory will be utilized at SDSF by establishing permanent plots systematically located across the Forest. Through this method, sample inventories can be completed periodically to monitor changes in Forest structure and growth rates over time.

VEGETATION TYPE

A new vegetation type map for SDSF was developed by staff foresters based on their on-the-ground experience, historical vegetation type maps, a remotely sensed imagery vegetation classification project (Clinton, 2009), and the 2006 inventory plot data. The vegetation map uses a standard California Wildlife Habitat Relationships classification system (Mayer and Laudenslayer, 1988) that includes tree species, structure, and density. Figure 5 (see Chapter 5) shows the vegetation type map for the Forest.

TIMBER SITE CLASS

In 2010, a timber site class map for SDSF was developed by staff foresters to update the timber type map for the Forest that was created in 1979 by Hammon, Jensen, Wallen, and Associates. The site polygon boundaries were based on their experience and knowledge of the productive potential of different areas on the Forest, guided by site tree data gathered in the 2006 inventory. Additional site tree data was also gathered to fill in any gaps in the site tree coverage. A total of 186 site trees were measured. Site class I is the most productive while site class IV is the least productive timberland based on dominant tree height at a given age. Figure 12 shows the timber site class map for the Forest. The most prevalent site class is III (1473 acres), followed by site II (1172 acres), and site IV (36 acres). There is no site I timberland in the Forest.

SILVICULTURE

MANAGEMENT GOALS


The overarching management goal for SDSF is to maintain the Forest as a managed working demonstration forest typical of managed forests in the area with a range of successional stages across the Forest. Early to late successional forest stages are managed across the landscape over time to meet research and demonstration mandates in the context of a changing climate, continually changing societal preferences and research needs.

FOREST DESCRIPTION

SDSF is well suited for harvesting with adequate stocking composed of coast redwood and Douglas-fir. Hardwoods, especially tanoak, are a large component Forest-wide but have been declining due to Sudden Oak Death. The normal process in forest succession is for conifers to overtop and suppress hardwoods, as is occurring in SDSF. This process will ultimately move parts of the Forest through its current mid-successional stage into a

late-successional condition. The 80-year-old conifer forest is reasonably healthy though overly dense in some areas (i.e., too many trees in any one location). SDSF will research and demonstrate ways to improve forest structure and vigor through timber harvesting.

Figure 12. Timber Site Class Map for SDSF.




The diameter distribution (tree diameter class plotted against the number of trees per acre) of SDSF resembles a traditional uneven-aged forest even though the forest is really a young, even-aged stand (Figure 13). Uneven-aged forests contain many diameter classes and at least three age classes. These forests are typically managed on a size and structure basis. Even-aged forests contain one to two age classes of trees and are managed according to age.

The diameter class distribution of a truly uneven-aged forest forms an inverse J-shaped curve, indicating decreasing numbers of trees as diameter increases. Smaller diameter classes commonly represent younger trees; the number of those smaller trees which mature into larger and, theoretically, older trees then decreases through time, as the curve shape implies. Having a greater number of trees in smaller diameter classes is preferable in order to ensure that an adequate amount of regeneration is present and can keep the forest viable through periods of natural mortality.

Redwood trees are shade tolerant and many will persist in the understory as small trees over many years. By harvesting competing trees and providing more growing space, the remaining

smaller trees in the understory will respond by increasing growth. Redwood trees have a tremendous ability to be released through removal of competition, which is a unique characteristic of the species. Harvesting also stimulates sprouting which introduces yet another age class to the Forest, but adequate sprout growth requires relatively high levels of light (Berril and O’Hara, 2009).

Figure 13. Conifer Diameter Class Distribution for SDSF.



SILVICULTURAL TREATMENTS

Silvicultural treatments involve management decisions and actions which direct forest growth, harvest, and regeneration. Between the late 1920s and early 1940s, SDSF was managed on an even-aged basis with clearcutting and natural regeneration. Currently, forests in Santa Cruz County, including SDSF, are managed on an uneven-aged basis as required by the Coast Forest District's Southern Subdistrict rules specified by the California Forest Practice Rules. Two ways to establish and maintain uneven-aged conditions are through single tree or small group selection harvest; both of these methods will continue to be demonstrated by SDSF. Although timber harvesting focuses on the removal of conifers, some hardwoods will also be removed to prevent hardwoods from dominating the residual stands. Sudden Oak Death is also taken into consideration for management decisions and silvicultural prescriptions as new information is learned and forest health conditions change across SDSF.

The silvicultural objectives are to achieve maximum sustained production, a broad range of diameter classes, multi-aged stands, promotion of growth, and encouragement of natural regeneration. Selective harvesting can improve stagnant forest conditions and enhance overall health. For example, removal of some trees from overcrowded stands will enable remaining trees to grow faster because of reduced competition for light, water, and nutrients. Sprouts and seedlings can also become established easier for the same reasons, thereby increasing both size and age diversity.

All silvicultural methods permitted under the California Forest Practice Rules may be used to varying degrees for research, demonstration, and creation of a range of seral stages with diverse forest structure. This protects the Forest's ability to adapt and change as silvicultural paradigms continue to change. Selection of a silvicultural prescription and cutting cycle for any given stand will depend on the specific conditions present, including vegetation types, site classes, historic management and the coordination required for resource protection, recreation and neighborhood concerns.

For most trees species the literature suggests that as trees mature, growth rates level off and slowly decline until natural mortality occurs. However, recent work on large, unsuppressed coast redwood trees in Humboldt County provides information pertinent to SDSF stands. This research suggests that as these trees age, ground-level measurements of annual growth (including height, ring width, and basal area increment) exhibited decreasing growth (or no change in growth) with age. Wood production of the entire main trunk and whole crown, however, increased with size and age up to and including the largest and oldest trees measured (Sillett et al., 2010). Trees similar to those that were measured in the study would be comparable to the protected old-growth trees at SDSF. Old-growth trees will not be included in uneven-aged management as they will be considered separately from the rest of the Forest (see the following Old-Growth and Late-Successional Management section.)

In most cases, forest regeneration will occur naturally after harvesting. This is the best means for SDSF since the dominant tree species, coast redwood, sprouts quickly and vigorously from remaining stumps given sufficient light. Research will be undertaken to study harvest levels required for successful regeneration and sprout growth. Berrill and O'Hara (2009) recommend reducing basal area by at least 45% at each cutting cycle in multiaged stands to provide sufficient light and sustain sprout growth and vigor. In areas that do not regenerate sufficiently (e.g., areas of Douglas-fir, heavy competition, or group openings), supplemental tree planting will take place. Tree planting will also take place for research and demonstration purposes, experimenting with alternative means of forest regeneration. Redwood or Douglas-fir nursery stock will be favored over hardwood seedlings except in areas where hardwood regeneration is insufficient.

SILVICULTURAL PRESCRIPTIONS AND LANDSCAPES

Silvicultural activities such as harvest and regeneration require planning which is typically outlined in silvicultural prescriptions. Silvicultural prescriptions for SDSF will encourage attention to landscape perspectives and patterns that benefit vegetation, wildlife, and other

values, as the Forest is managed for a range of successional stages across the landscape.

Management from a landscape perspective focuses on landscape patterns and the responses of biotic (e.g., plants, animals) and abiotic (e.g., soil and water) factors to changes in those patterns. A large part of this type of management involves the response of biota to both natural and unnatural disturbances (such as burned areas or opening stands through logging) in the landscape, also known as edge effect. Edge effect, considered to be both beneficial and detrimental, can have impacts on species that are adapted to interior forested habitats. The use of selection methods for harvest, however, can help minimize negative impacts by eliminating sharp contrasts between areas of harvest and adjacent forested areas.

As part of demonstration and research objectives, SDSF will experiment with different sizes of group selection openings as well as feathering the edges of openings. The main purpose of experiments is to determine the extent of edge effect on biota associated with different harvest levels and sizes of group selection openings. Results of demonstration and research projects will be used in the development of future silvicultural prescriptions and activities.

CUTTING CYCLE

Designated forest stands will be available for selective harvest every 10 years and successive entries may be up to 20 years or more in some stands. The length of cutting cycles for individual stands will depend on tree densities, diameter distributions, growth rates, stand objectives and other parameters.

Traditionally, harvesting removes the amount of growth that has occurred since the previous harvest. However, over the next decade in most stands within SDSF not all of the growth between cutting cycles will be removed (see the Harvest Planning section). By not removing all of the growth, the amount of volume remaining following each stand entry will increase over time. This reflects SDSF's mandate of remaining a demonstration of a financially viable working forest while at the same time enhancing public trust resources.

SILVICULTURAL TREATMENTS AND NON-TIMBER RESOURCES

One of SDSF's demonstration goals is to display that silvicultural treatments are compatible with aesthetic, biological, and recreation values. Ways to display this compatibility may include harvest methods which are aesthetically acceptable both from a distance and close-up, marking treatment boundaries so that they have minimal visual impact from roads or trails, and providing for habitat improvement through silvicultural prescriptions. These and other examples may be elements of demonstration programs directly related to timber harvest.


OLD-GROWTH AND LATE-SUCCESSION MANAGEMENT

OLD GROWTH

Existing old-growth redwood trees in the Forest have been excluded from all future timber

harvesting, as mandated by SDSF's authorizing legislation AB 1965. Figure 14 shows that the only stand of old-growth trees is located in the Badger Spring area. There is also a small group of old-growth trees located near Sulphur Springs. Individual old-growth conifer trees are identified, marked for preservation and protection during timber harvest preparations and entered into the Forest-wide old-growth tree database. Individual tree characteristics, listed in Table 7, helps foresters to identify these old-growth trees in the Forest. (Table 7 compares tree characteristics for old growth against those of young growth.) No individual characteristic, including age, is used to make a classification. Instead, combinations of these characteristics are used to determine if trees are classified as old growth or young growth.

Figure 14. Old-growth Stand and Late-Succession Management Areas.



LATE SUCCESSION

Late-succession management areas will be administered to promote the development of functional old-growth habitat characteristics through infrequent, low-intensity timber management activities. These characteristics include multi-level canopy structure which provides vertical and horizontal diversity, stand-age diversity, large trees, snags, down logs, and other woody debris. Recruitment and retention of these characteristics, which provide important habitat elements for fish and wildlife species, will be attained through silvicultural prescriptions. For example, trees with crowns that extend over stream channels will not be harvested unless a qualified hydrologist or wildlife biologist recommends removal of such trees to enhance riparian habitat or hydrologic function. These trees may also be removed when recommended by a Certified Engineering Geologist to help stabilize active slides. CWHR classifications will be used to assess and maintain the specific wildlife habitat characteristics of these areas in SDSF.

Table 7. Individual Redwood and Douglas-fir Tree Characteristics.

<u>YOUNG GROWTH</u>	<u>OLD GROWTH</u>
Abundant branches or knots in the lower 1/3 of the bole (tree trunk).	Lower third of the bole is relatively free of branches or knot indicators.
Branches are small, 4 inches and under.	Large branches, many greater than 4 inches in diameter
The tops of the crown are usually pointed and the branches are usually upturned.	Tops of crown are rounded or flat. Branches may be downturned.
Bark is typical of younger trees as described in dendrology books (shallow bark furrows).	Bark is typical of older mature trees as described in dendrology books (deep bark furrows, bark between furrows often plate-like).
Growth is generally fast, as indicated by large annual rings - usually less than 10 annual rings per inch.	Slow current growth as indicated by a long period of narrow annual rings - usually 15 or more annual rings per inch.
Excessive taper in open grown trees.	Very little taper, even in open grown trees.
Few fire scars or superficial fire scars and burned bark.	Large old fire scars, burned bark and goose pens common.
Tree age less than 175 years.	Tree age 175 years or greater.

Like areas of old growth, sections in the Forest designated for late-succession management have been identified and mapped (Figure 14). These areas make up 15 percent of the overall Forest, and correspond to a width of 300 feet on each side of the East Branch of Soquel Creek, Amaya Creek, and Fern Gulch within SDSF boundaries (all Class I watercourses). Timber harvesting in these areas conforms to the following guidelines, as well as the Anadromous Salmonid Protection (ASP) Rules approved by the State Board of Forestry and Fire Protection in 2009:

- * At least 75% total canopy in multiple layers retained with 80% post - harvest overstory canopy from 30 to 100 feet from the watercourse transition line (WTL).
- * At least 25% canopy in trees at least 24 inches diameter at breast height (dbh) retained.

- * All woody riparian (i.e., hydrophytic) vegetation retained except where riparian function would be enhanced by removing such vegetation.
- * Large snags (i.e., standing dead trees at least 20 inches dbh and 15 feet tall) or live wildlife trees (i.e., trees that support bird nests or have cavities or large limbs that make them valuable for nesting birds) will be recruited (created from existing healthy trees) or retained with the goal of reaching an average density of at least five per acre.
- * At selected locations where conifers are lacking and would not compete with riparian vegetation, Douglas-fir and redwood may be planted to promote long-term recruitment of large woody debris in streams.
- * Downed logs at least 24 inches in diameter and 30 feet long will be retained with the goal of reaching an average density of at least two per acre, and total coarse, woody debris will be retained at an average density of at least 10 tons per acre.
- * No harvest within the Core Zone or first 30 feet from the watercourse transition line (WTL).
- * A post-harvest stand that retains the 13 largest conifer trees (live or dead) on each acre that encompasses the Core and Inner Zones.
- * Large trees retained are to be those most conducive to recruitment to the watercourse channel.
- * Additionally, harvesting is prohibited in channel migration zones and additional protection measures are mandated where there are flood prone areas (see 14 CCR 916.9 (f) in the California Forest Practice Rules).

These guidelines will be updated in the future as needed to reflect rule changes for the protection of anadromous salmonid species.

Site disturbance during harvesting operations in late-succession management areas will be kept to a minimum by restricting tractor use and cable-yarding corridors to predesignated trails. Furthermore, all trees to be harvested and all wildlife trees and downed logs to be retained will be marked. A Professional Geologist and a qualified hydrologist will also be utilized to review operations during timber harvest planning.

Areas along the creeks were chosen as late-succession areas for a number of reasons, including protection of sensitive riparian zones, fisheries resources, and existing Watercourse and Lake Protection Zones (WLPZ) rule requirements. Furthermore, these areas provide habitat corridors and buffers between stream channels and nearby harvested areas. Corridors are important for wildlife movement and survival, providing a valuable element to the landscape pattern.

HARVEST SYSTEMS

The terrain where logging activities are to occur will be the key factor in determining whether ground skidding, cable logging, or aerial harvest systems are used. The erosion hazard, slope angle, slope stability, and distance to drainages will also be carefully evaluated in the selection of harvest systems. Additional factors include access, public safety, aesthetics, timing, noise, environmental mitigation, economic conditions, research, and demonstration aspects. The appropriate harvest system for each timber sale will be identified and utilized based on the above considerations. Refinements on the harvest systems used will be made as necessary to accomplish harvesting in an economical, efficient, and environmentally sound manner.

SYSTEMS FOR USE IN SDSF

Ground skidding logging systems, methods which pull logs on the ground to landings (loading areas), consist of tractor, rubber tire skidder, and horse logging. Horse logging, limited to gentle slopes and modest sized trees, can be used to minimize soil disturbance, damage to residual trees in dense stands, or for demonstrative purposes. Tractors and rubber tire skidders are generally used where slopes average less than 35 percent and on steeper slopes after evaluations of slope stability and erosion hazard potential are completed. Operations are contingent upon the ability to mitigate any impacts to negligible proportions. Tracked equipment with an attached, open ended holding container, called a forwarder, is utilized for moving split firewood to a road or landing for loading into a truck.

In SDSF, ground skidding equipment will be restricted to the minimum size capable of moving the harvested timber. Skid trail systems (temporary trails used by ground skidding equipment) will be designed so that existing skid trails are utilized where appropriate. All new and rehabilitated trails will be predesignated. A Certified Engineering Geologist (CEG) will participate in the preparation of all timber harvesting plans. CEGs will review the locations of all proposed roads, skid trails, and landings on unstable soils, unstable areas, and areas where the California Forest Practice Rules require that skid trails be designated before the start of timber operations.

Skyline logging, commonly referred to as cable logging, uses cables to fully or partially suspend logs off the ground while transporting them to designated landings. The cables are attached to a machine, called a yarder, which is equipped with multiple winches and a tower for pulling the cables. A wide range of cable systems are available for logging both large and small timber. In SDSF, cable yarding will generally be used on steep slopes, near drainages, or where road construction is difficult.

Aerial harvest systems incorporate the use of helicopters to move logs from one area to another. These systems are beneficial when the erosion hazard or slope instability are high or when access to an area is limited. In SDSF, aerial systems will be considered primarily for demonstrative purposes and where cable yarding is not possible. Helicopter timber drop-off and landing areas will be located at least 0.5 mile from the nearest occupied dwellings.

A preliminary logging plan for SDSF designated proposed tractor and cable yarding areas,

existing and planned roads, and old growth areas. Based on this plan, approximately 1,700 acres are suitable for yarding by tractors and 900 acres are suitable for yarding by cable yarders or helicopters.

HARVEST SYSTEMS AND COMPATIBILITY

Harvesting operations will be planned and implemented to minimize conflicts with neighbors and visitors where possible. To accomplish this, road and skid trail construction will be designed with potential users in mind. For example, specific logging trails can be designed to serve as future recreation trails where appropriate. Certain logging trails may be closed, however, following harvesting activities where public access is inappropriate (e.g., research or sensitive areas, or where trespass across private property would occur). New roads and skid trails will be carefully located to minimize visual impacts. Additionally, the size and duration of area closures and traffic diversions will be evaluated for each timber sale.

HARVEST PLANNING

HARVEST LEVEL

Short Term

SDSF's annual harvest level for the planning period from 2010 to 2020 will average between 800,000 and 900,000 board feet (approximately 30 to 35 percent of annual Forest-wide growth). This harvest level is equal to 8.6 million board feet per decade and is consistent with AB 1965 in that it provides for timber management demonstrations and promises an intensively managed research forest.

So far during planning period one (2010 to 2020) in year 2013, 3.8 million board feet have been harvested on SDSF. This harvest rate is substantially lower than SDSF's current conifer growth rate of 2.6 million board feet per year and is less than one percent of the standing conifer inventory. It is expected that a timber harvest of 1.5 to 2 million board feet will occur every other year starting in 2014.

Long Term

The projected harvest levels for the first decade (2010 – 2020) are approximately one-third of current growth. If future harvest continues at this level, the forest will develop denser stands with bigger trees and more closed canopy over time. The harvest level is projected to gradually increase as the Forest inventory and growth increases. It is anticipated that the funds needed for the maintenance and operation of SDSF, reasonable capital costs and other expenses incurred in fulfilling the objectives of PRC sections 4660 – 4664 on SDSF will only be fulfilled once long term harvest levels are achieved.

Forest stocking and growth will be reevaluated in the second decade (2020 to 2030) when a new Forest inventory is conducted and updated data are used to project future conditions once again. The harvest level on SDSF will be consistent with the Management Goal in Chapter 2 to “Demonstrate sustained-yield timber harvest practices through harvest operations that balance harvest rates with growth over time and are compatible with rural land use in Santa Cruz County, while promoting recreation opportunities, forest health,

watershed protection, wildlife, and fisheries values as well as aesthetic enjoyment.”

Sustained yield production is achieved by balancing growth and harvest over time, maintaining a timber inventory capable of producing the long-term sustained yield (LTSY) (i.e., the harvest level projected to be sustainable after 100 years), and having the projected annual harvest level for all future rolling ten-year periods not exceed the LTSY. These results were established by projecting stocking conditions, growth, and harvest levels for the planning area over a 100-year period based on the best available information and accepted principles of forest management and statistics. During the development of the 1998 General Forest Management Plan for SDSF, a compromise was made to limit harvest levels at SDSF to no more than 50-60 percent of forest-wide growth regardless of LTSY or revenue needs. There was a fear that SDSF would be used to generate much needed funds for California and that the revenue from timber harvesting would not be used consistent with fulfilling the objectives of PRC Sections 4660-4664. Since SDSF was established, the total harvested volume has been 5,986,000 board feet over a period of 23 years. This amount of volume is accumulated in growth at SDSF in just over two years.

To evaluate timber production capabilities the land area within SDSF was grouped according to its availability for timber harvesting: unavailable (e.g., inoperable areas, old-growth protection areas, and predominantly hardwood areas); limited availability (e.g., late-succession management areas, recreation sites, and unstable areas); and fully available. As a result, 60 percent of SDSF (1,609 acres) is fully available for timber production, 30 percent (804 acres) has limited availability, and 10 percent (268 acres) is unavailable (Table 8). All long-term sustained yield values are based on the full and limited availability acres only. Unavailable acres are not included in the analysis. Figures 15 and 16 illustrate the inventory, growth and harvest rates projected over ten planning periods generated from this analysis based on the 2006 timber inventory data and current constraints on the harvest level.

In future analysis and planning efforts, the short and long term harvest levels will be reevaluated based on the best available information at that time. The level of timber harvesting shall not exceed LTSY and on a cumulative basis shall be limited to the level of timber harvesting necessary to provide the funds needed for the maintenance and operation expenses of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF.

Table 8. SDSF Forest Land Availability.

Timber Capable	Condition Class	Acres	Percent of Ownership
	Full Availability	1,609	60
	Limited Availability	804	30
	Unavailable	268	9.9
	Subtotal	2,677	99.9
Non-Timber Capable	Non-timber	4	0.1
	Ownership Total	2,681	100

Table 9. Projected SDSF Forest-wide (full and limited availability) Average Annual Inventory, Harvest and Growth by Planning Period.

Period	Year	Inventory MMBF ⁽¹⁾	Inventory per Acre MBF ⁽²⁾	Harvest per Year MMBF ⁽¹⁾	Growth per Year MMBF ⁽¹⁾
1	2010-2020	112	46.2	0.866	2.25
2	2020-2030	125	51.9	1.03	2.37
3	2030-2040	139	57.5	1.18	2.65
4	2040-2050	153	63.6	1.45	2.91
5	2050-2060	168	69.6	1.49	3.00
6	2060-2070	183	75.9	1.50	3.10
7	2070-2080	199	82.5	1.55	3.15
8	2080-2090	215	89.0	1.57	3.23
9	2090-2100	232	96.0	1.60	3.28
10	2100-2110	248	102.9	1.60	3.34

1) MMBF = million board feet

2) MBF = thousand board feet

Figure 15. SDSF Projected Inventory by Planning Period.





Figure 16. SDSF Projected Annual Growth and Harvest.



TIMBER SALE PROCESS

An important demonstration for SDSF is to support a viable local forest products industry. A local forest products industry is essential to maintain managed, working forests in the region. Financially viable managed forests in turn are an important tool to maintain large contiguous areas of wildlands in the Santa Cruz Mountains and to avoid land conversion and development pressures.

SDSF staff will implement a timber sale every year or every other year. The scarcity of sawmills in the region reflects the tenuous and sometimes unpredictable nature of timber supply in the Santa Cruz Mountains. A predictable and consistent supply of raw materials from SDSF is important in order to contribute to the viability of these sawmills and to ensure their continued existence.

Each plan will specify, at a minimum, the area and volume to be harvested, logging and silvicultural methods to be used, and restoration and protection measures necessary to address environmental concerns. All potential harvests will be evaluated by CAL FIRE Forest Practice staff and will be available for public review.

Once a timber sale has been approved by the CAL FIRE Director, a minimum bid will be established for the sale. The timber designated for harvest will then be sold to the highest bidder. The timber purchaser will enter into a contract with the State which specifies all requirements in detail, including the operation and payment schedule, adherence to applicable laws, and any additional improvements or demonstrations. SDSF staff will administer the sale, assuring that the contractor abides by the contract. Furthermore, CAL FIRE Forest Practice Inspectors from the San Mateo-Santa Cruz Unit and resource professionals from the other Review Team agencies will conduct a Pre-Harvest Inspection and periodically inspect the operation during and after logging. Additionally, a Waiver from a Waste Discharge Requirement will be obtained from the Central Coast Regional Water Quality Control Board prior to the commencement of timber operations as required.

The timber sale process will be administered through the CAL FIRE Class III permit system for major timber sales. Periodically, small timber sales (less than 100,000 board feet) may be offered for demonstration or research purposes. These harvests would require a Class I minor timber sale permit. Any conifers removed in minor sales or lost through catastrophe (earthquake, wind, fire, flood, or pest infestation) must also be applied to the designated harvest level. In the event of a major catastrophe, salvage logging would occur and the annual harvest level would be recalculated.

It is generally accepted that winter timber harvesting will not occur at SDSF. This means that no heavy equipment operations or log trucking will occur during the winter period, beginning October 15 and ending May 1. Certain activities will require a winter operations plan to be included in a THP. The activities that may occur in the winter period are lopping of slash, falling of timber outside of the stream protection zones during dry periods, tree planting, erosion control maintenance, and firewood cutting and splitting.

TIMBER OPERATIONS MONITORING

To avoid adverse impacts on water quality and fisheries resulting from the discharge of sediment to watercourses attributable to timber operations, SDSF staff will monitor all timber operations (including all harvesting areas and new roads, skid trails, and landings) annually for five to seven years following completion of the operations. Occurrences of substantial surface erosion (e.g., gullies) or mass wasting (e.g., landslides or slumps) resulting from the operations will be identified and described by a Registered Professional Forester. Each substantial gully or landslide will be evaluated to determine its cause, and stabilization measures will be identified that will be most feasible and effective. Such measures will be implemented within 90 days from the date when the subject site is identified, unless due cause for delay is explained and a reasonable alternative schedule for implementation is proposed by the Forest Manager.

Appropriate mitigation measures to be incorporated into future timber operations will be described and specified in future timber harvesting plans to avoid a recurrence of the observed erosion or mass wasting events (i.e., adaptive management).

TIMBER MARKETS


The local timber market is largely influenced by the proximity of sawmills and economic conditions. The closest large sawmill to SDSF is the Big Creek Lumber Company mill, located 17 miles north of Santa Cruz near Davenport. The next closest sawmill is Redwood Empire Sawmills in Cloverdale, 165 miles north of Santa Cruz on Highway 101.

The timber market has historically undergone fluctuations as a consequence of economic conditions. Figure 17 indicates dramatic changes in redwood stumpage prices during the time period 1994 to 2012 (California State Board of Equalization, 2012).

Stumpage prices reflect the value of logs delivered to the mill less the costs of logging, hauling, and cleanup. In the 1990s, stumpage prices increased significantly and have widely fluctuated since then. Factors contributing to the rise in the 1990s included a decline in federal timber sales, a reduction of available timber elsewhere, increased forest regulations, and the protection of threatened and endangered species. Starting in 2008, the demand for forest products declined with the economic recession and decreased markets for lumber related to the housing market. These factors, along with the cyclical nature of the timber market, are expected to continue to influence market activity.

Timber values for SDSF are not expected to be equal to general redwood/Douglas-fir prices. Bidding competition for SDSF sales will generally be lower because of the scarcity of sawmills in the region. Harvesting operations will usually be more expensive because of the amount of rehabilitation and restoration work planned. For example, existing undersized road culverts will be replaced, existing roads will be reshaped and surfaced to reduce erosion and additional fire hazard reduction work will be performed. These revenue-reducing activities will be added to the cost of harvesting.

Figure 17. Santa Cruz County Redwood Stumpage Values.



HARDWOOD MANAGEMENT


The 2006 timber inventory estimated hardwood basal area to be 116 square feet per acre. Figure 18 represents the hardwood basal area by the three most common species and a grouping of lesser occurring species (live oak, willow, alder, and bigleaf maple). As the graph indicates, tanoak is the most common hardwood species present. Hardwoods are a significant component of SDSF and, on average, comprise 40 percent of the basal area. Since the 2006 inventory however, dramatic visual indications of a shift in this composition are evident. Many areas where madrone and tanoak occurred in greater abundance than bay-laurel have seen significant mortality in the tanoak and madrone from several pathogens that affect these species and not the bay-laurel. Various experiments and demonstrations will focus on improving the health of these stands and reestablishing a diversity of species.

Hardwood stands will be evaluated during wildlife and timber management planning. The Forest type map discussed in the Conifer Volume Inventory section classifies hardwoods as well as conifers in SDSF. Areas that have always been hardwood will remain so and will be examined for wildlife habitat enhancement and research opportunities. Approximately 100 acres of SDSF timberlands that naturally support conifers but, as a result of past logging and other management, currently support pure or predominantly hardwood stands will gradually be harvested and reforested where appropriate by planting redwood and Douglas-fir seedlings. These stands are widely distributed throughout the Forest in units one-quarter to one acre in size. An analysis of potential biotic and wildlife habitat impacts will occur prior to all hardwood conversion. Any individual openings will not exceed 0.5 acres. These areas

will be treated by planting group openings or by selectively removing a portion of the canopy and interplanting with conifer. Focusing on areas along the edges of conifer stands has proven the most successful in other areas of the Santa Cruz Mountains. Experimentation with differing levels of canopy reduction as well as prescribed burning in these areas may be pursued as a demonstration.

Another aspect of hardwood management is to examine and experiment with alternative uses for hardwoods. The goal is to efficiently utilize more raw material, targeting portable sawmill owners, timber operators, and private property owners in the process. Some items that can be made from hardwood lumber include furniture, paneling, flooring, and decking. Hardwoods can also be utilized as fuelwood and chips. SDSF will research and demonstrate the management and use of hardwoods as a timber resource.

Figure 18. Estimated Basal Area for SDSF's Hardwood Tree Species.



DEMONSTRATION AND RESEARCH PROGRAMS

SDSF's demonstration and research programs will integrate all facets of timber management consistent with PRC Sections 4660-4664 and subject to the limitations of this management plan. Each timber sale will serve as a demonstration but also may contain a research component. Research projects will evaluate the applicability, practicality, and effectiveness of various strategies of forest and watershed management. Timber management demonstrations will include planning and operational projects as well as tours showing the risks and benefits of timber harvesting. Additional information on projects can be found in the Demonstration and Education and Research Chapters.

Findings of research and demonstration projects will be available to the general public, small private landowners, researchers, and the forest products industry. Information will be distributed through research publications, direct mailing, libraries, and posting on the internet (see http://www.fire.ca.gov/resource_mgt_stateforests_publications.php).

SDSF can best serve the general public by providing information and experiences which enhance understanding of forest management principles and good land stewardship. The best way to provide learning opportunities is through exposure to forest management activities. Demonstration subjects might include the complexities of land management, trade-offs involved with timber harvesting, and the extent of wood products utilized by consumers. With this information, the public can make their own informed decisions regarding the effects of forest management on the environment.

Small private landowners can gather useful information and learn to plan more efficiently by viewing land management alternatives. Visiting project sites and observing results is an effective way to understand management techniques and how these might apply to private land. Of most benefit will be practical, directed projects that clearly demonstrate timber management, natural resource enhancement techniques, and solutions to specific problems.

Timber operators and industrial land managers will have the opportunity to view various harvesting and environmental protection techniques. Demonstration and research project ideas will be solicited from this group. Possible projects may include evaluating methods to enhance forest health and regeneration, testing new harvest and management techniques, and solving operational problems.

MINOR FOREST PRODUCTS

The primary minor forest product that will be sold from the Forest is fuelwood. Fuelwood, primarily from hardwood tree species, will be available on an intermittent basis, usually as the result of other management activities. These activities may include road work, fuel break construction or maintenance, by-products of timber sales, and wildlife habitat improvement projects.

Additional forest products may be sold as they become available. These include salvage sawlogs, redwood split salvage, cull logs, roots and stumps, posts, boughs and other greens. As with the hardwood management program, the use of these products promotes utilization of raw materials that might otherwise accumulate and present safety or fire hazards.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Comments and concerns regarding SDSF's timber management program have been very detailed and encompass a variety of subjects. Since it is impossible to address every comment directly, five broad topics of concern which cover most comments have been

identified.

The first and probably most important concern to neighbors and other users of SDSF involves disturbance caused by logging activities. Logging truck traffic on mountain roads, reduced access to the Forest, alteration of the forest viewshed, and noise are sources of apprehension for many individuals. As stated in the management guidelines below, SDSF will strive to reduce possible impacts to forest neighbors and users during all forest management activities.

Another concern that has generated many comments deals with the possibility of negative impacts on Forest wildlife. Comments generally focus on the effects of timber harvesting on wildlife habitat, including concern for streams and adjacent riparian areas. SDSF plans to evaluate and monitor the response of various plant and animal species to forest management activities. Results of studies may include strategies to improve adverse conditions, enhance mediocre areas, or maintain exceptional situations.

The third area of attention deals with revenue from State Forest harvests. Interested individuals have expressed an opinion that SDSF should only harvest enough to cover basic expenses and that revenue should only be used for these expenses. CAL FIRE recognizes that PRC Sections 4660-4664 limits the level of harvests on a cumulative basis to that needed for the maintenance and operation of SDSF, reasonable capital costs, and other expenses incurred in fulfilling the objectives of PRC Sections 4660-4664 on SDSF. CAL FIRE must also comply with California Forest Practice Act and Rules and Board of Forestry and Fire Protection policy. The Department will not attempt to accurately estimate revenues over any specific time period. As revenues increase over time, the Department will request expenditure augmentations through the normal budget process. Final state forest allocations ultimately rest with the Legislature.

Concern for the natural instability of the Soquel Creek watershed and excessive soil loss is also prevalent. Logging activities in the Forest will adhere to California Forest Practice Act and Rules which limit road and skid trail construction. These regulations require site-specific mitigation as necessary to reduce erosion to minimum levels. Additionally, adherence to the Anadromous Salmonid Protection Rules, environmentally sound logging practices, input from Professional Geologists for all timber sales, and experimental research will be used to minimize damage to this sensitive area.

Finally, neighbors of SDSF would like to see local loggers, trucking companies, and mills perform the felling, hauling, and milling of products from the Forest's timber sales. While this may be ideal, the bid process cannot be limited to local businesses. In some cases, the highest bidder will likely be local, keeping the work and revenue in Santa Cruz County.

MANAGEMENT GUIDELINES

1. Create and maintain a mosaic with a range of seral stages across the Forest that changes over time. Optimize the Forest's attraction as a working forest and research destination to best meet the multiple use mandate.

2. Maintain and enhance a healthy forest ecosystem. This includes the monitoring of basic resources and requires management activities that ensure forest vigor. The Forest will be periodically surveyed on an informal basis for general health, with emphasis on disease and insect activity, tree growth and vigor, and soil stability. Other monitoring activities, such as those developed for fisheries and wildlife, will contribute additional information on the health of individual resources within the ecosystem.
3. Protect and monitor the watershed, soil, fisheries, and wildlife resources during all timber harvesting activity. The fundamental goal is to minimize impacts to these resources through planning and mitigation developed on a site specific basis. Harvesting operations will be planned to occur in an orderly fashion across the plan area and will be completed from the back to the front to minimize the need to re-open landings, skid trails and roads one year to the next. Protection measures may include selecting low impact harvest methods, avoiding sensitive areas, monitoring for the introduction of invasive exotic plants and conserving or improving resource integrity. Timber harvesting will not occur during the period of winter operations (October 15 through May 1), the period prescribed in the ASP Forest Practice Rules, except as noted in the Timber Sale Process section.
4. Design timber management activities based on landscape perspectives. Components to consider will include horizontal and vertical forest structure, vegetation density, edge effect, corridor size, and biological diversity.
5. Timber sales will have demonstrational value and include experimental and educational aspects whenever possible. This may include pre-harvest and post-harvest activities as well as actual harvest procedures.
6. Consider neighbor and visitor concerns during all timber harvesting activities. SDSF will strive to reduce excessive noise, visual impacts, and transport activity. Logging methods and haul routes that facilitate reduced disturbance will be evaluated on a site-specific basis and used as appropriate.
7. Demonstrate timber management compatibility and integration with recreation. Whenever possible, design timber sales to minimize conflicts with recreational use and improve recreation facilities such as roads and trails. Additionally, safe recreational behavior during logging activities will be encouraged through signs, direction from SDSF staff, and alternative routes.
8. Monitor all timber operations annually for five to seven years following completion of the logging operations. Any substantial surface erosion or mass wasting found will be identified and described by a Registered Professional Forester and evaluated to determine its cause. Stabilization measures that will be most feasible and cost effective will be identified and implemented within 90 days (see Appendix C). Each timber harvesting operation will include sediment source remediation. High-priority

remediation sites will be considered when selecting areas for upcoming harvests. In some cases, remediation at locations other than timber harvest areas could constitute offsite mitigation for the watershed impacts of harvesting.

PLANNED ACTIONS

1. Harvest between 800,000 and 900,000 board feet per year for the period from 2010 to 2020. This is estimated to be approximately 30-35 percent of Forest-wide growth.
2. Protect all old-growth redwood and old-growth Douglas-fir trees in the Forest. Maintain and update the old-growth tree data base.
3. Promote the development of functional old-growth habitat characteristics in late-succession management areas within 300 feet of the East Branch of Soquel Creek, Amaya Creek, and Fern Gulch. Follow the Anadromous Salmonid Protection Rules for protection of Class I, II, and III watercourses to enhance riparian functions and to help recover state and federally listed fish species.
4. Identify anticipated harvest areas for the planning period up to 2020 and develop a Management Unit Map. Evaluate all possible harvesting and silvicultural methods, new road construction needs, and compatibility with other forest uses for each area under consideration. Planning for future harvests will include consideration of potential impacts to smaller watersheds and their future management to allow for well-designed monitoring of potential impacts on water quality.
5. Refine a forest type map which includes tree species, CWHR classifications, forest structure, and vegetation density. As resources permit, continue to develop and maintain a Geographic Information System (GIS) database which will eventually include information on soil characteristics, streams, topography, research sites, roads, trails, facilities, and other improvements.
6. Reinventory the Forest on a regular schedule, using either a temporary plot system or by installing a continuous forest inventory system of permanently monumented monitoring plots, to be remeasured at regular intervals. The next inventory will occur between 2016 and 2021.
7. Within each new project area, whether it be a THP or other experiment or demonstration, hardwood stands will be considered for management. Some areas of hardwood will be managed by single tree selection harvesting or group selections. These areas will be planted with conifers as appropriate. Hardwood trees with particular wildlife value will be retained. Experimental treatments will also be demonstrated (i.e., reduction of bay-laurel in stands with extreme oak and madrone mortality).
8. Monitor timber operations areas for infestations of invasive, exotic species. Eradicate new populations prior to them becoming established and producing a seed bank.

CHAPTER 10: RESEARCH

INTRODUCTION

It is the State Board of Forestry and Fire Protection's policy for all State Forests to conduct innovative research in forest management. The purpose of such research is to provide resource management information to the general public, small forest landowners, and the forest products industry. Research opportunities in SDSF are numerous, offering small or large, general or specific, and experimental subjects. Individuals conducting research may represent private organizations, including consulting firms and environmental protection groups, or public agencies, such as institutions of learning or resource-based departments.

A plan to identify specific subjects of concern and research priorities for SDSF needs to be formulated so that continuous and long-range studies can commence. Forest staff will investigate possibilities and rely heavily on what the public, CAL FIRE officials, and resource professionals would like to see analyzed.

COMPLETED PROJECTS

Several research projects and surveys, summarized below, have been completed for resources within SDSF. Procedures and results of each study are outlined in their respective chapters with the exception of the Geologic Survey (Manson and Sowma-Bawcom, 1992), which is covered in the Property Description Chapter.

ARCHAEOLOGICAL STUDIES

A complete surface field survey for archaeological and historical sites was conducted in 1991 by Dr. Brian Dillon. Dr. Dillon is a consulting archaeologist affiliated with California State University at Northridge. Sites found during this study will be protected for cultural, research, and educational purposes.

Additional surveys for archaeological and historical sites have been completed for timber harvest plans, road and trail construction sites, and other projects. Several historic artifacts and archaeological sites have been recorded by CAL FIRE staff and protected.

A cultural resources study of the six-acre Badger Spring Picnic Area was completed in 2011 by Patricia Paramoure, a Master's student at Sonoma State University.

These confidential archaeological reports are filed with the Office of Planning and Research- State Clearinghouse.

BIOLOGICAL ASSESSMENT

An inventory and assessment of SDSF's biota was completed by biologists from California Polytechnic State University, San Luis Obispo, in 1992. The survey resulted in lists of all plants and animals seen, heard, or tracked in the Forest (refer to Appendix B for species lists). The study, which concentrated on special status species, was under the direction of Professors V.L. Holland and Mike Hanson (Holland et al., 1992).

Several bird surveys were conducted by wildlife biologist David Suddjian, including raptor surveys for the Fern Gulch THP in 2001, 2005 and 2006, and a Marbled Murrelet survey in 2003 and 2004. After ten snags were created in Douglas-fir trees as part of the Long Ridge THP in 1995, Mr. Suddjian published "A Tale of Ten Snags" article in the September/October 2001 issue of the Santa Cruz Bird Club Newsletter and also presented a poster, along with Forest Manager Thomas Sutfin, at the Redwood Science Symposium, March 15-17, 2004 in Rohnert Park. After monitoring the post-harvest bird occupancy in these snags, he found that increased bird populations, new tree cavities, active nests and foraging evidence suggested that the snag management program was successful.

Raptor surveys of the Fern Gulch and Rim THP areas were completed by RPF Matt Greene in 2011 and 2012 (Greene, 2011, 2012). Two new species were observed (wild turkey and pileated woodpecker); they have been added to the end of the Birds list under "Fauna of SDSF" in Appendix B.

A botanical survey of the Fern Gulch area was completed in 2002 by Tim Hyland (Hyland, 2002) and updated in 2010 by Tim Hyland and Dylan Neubauer (Hyland and Neubauer, 2010). Ms. Neubauer completed a botanical survey of the Comstock Mill THP area in 2013 (Neubauer, 2013). Lists of the species they observed have been added to the end of the "Flora of SDSF" list in Appendix B.

A study of blood-borne disease in SDSF was started in 2009 under Professor Janet Foley at the University of California Davis School of Veterinary Medicine, and continues currently. The study investigates blood-borne diseases including Lyme disease and anaplasmosis by surveying their hosts, which include ticks and small mammals (Foley, 2011).

A herpetological survey, started in 2010, continues currently with the help of volunteers under the North American Field Herping Association (NAFHA). The purpose of the survey is to verify and photo voucher all varieties of reptiles and amphibians existing in SDSF (Erickson, 2011, 2014). The list of amphibians and reptiles that have been observed are in "Fauna of SDSF" in Appendix B.

Researchers from the Santa Cruz Mountain Puma Project through the University of California at Santa Cruz have been studying mountain lions in SDSF since 2009 using telemetry collars to collect continuous movement and location data. This study,

led by ecologists Dr. Chris Wilmers and Paul Houghtaling, seeks to determine the success rate, effort and community-level consequences of predation, to understand cougar habitat requirements, and to provide guidance on movement corridors for lions within and between the mountain ranges on the central California coast. No formal reports have been provided to SDSF staff, but more information may be found at <http://wildlife.ucsc.edu/>.

GEOLOGIC SURVEYS

An extensive geologic survey of SDSF involved detailed mapping of geologic features and areas damaged during the Loma Prieta earthquake. This analysis was conducted in 1992 by Michael Manson and Julie Sowma-Bawcom of the California Division of Mines and Geology (renamed California Geological Survey) (Manson and Sowma-Bawcom, 1992). Mapped information includes geologic characteristics, landslides, stream orders, and areas where mitigation work can be done. Additional geologic surveys have been completed for timber harvest plans, road and trail construction sites, and other projects.

RECREATION STUDY AND PLAN

A recreation survey to determine Forest recreational users' views and use patterns was sent to the neighbors of the Forest and the local mountain community in 1992 (Hester et al., 1992). Copies of the survey were also distributed in the State Forest and in The Forest of Nisene Marks State Park. Results of the study have been a primary source of public input into the forest management planning process. The principal investigators for the study were Randolph Hester and Marcia McNally from the University of California, Berkeley. The Recreation Study Final Report by McNally and Hester was completed in 1993 (Hester et al., 1993). The Draft Education and Recreation Master Plan was completed in 1996 (University of California, 1996).

EDUCATION

Researchers from the University of California at Berkeley, led by Nova Blazej, completed the Education Study in 1997 (Blazej, 1997). It was intended to inform the development of the SDSF Education and Recreation Plan and to be used by SDSF education coordinators and volunteer coordinators as a planning tool for developing forest education programs. Five basic themes were identified to shape the framework of SDSF education programs: forest history, ecological processes, forest management, regional connections and the spirit of the Forest.

Three significant and detailed studies regarding the proposed SDSF Forestry Education Center (FEC) were completed under the leadership of Barbara Butler from the University of California at Berkeley in 2001. The first consisted of a series of interviews with decision makers in the Monterey Bay Area regarding the possible role and educational niche of the SDSF FEC (Butler et al., 2001).

The second was an in-depth analysis of ten environmental education centers in California to determine what kinds of facilities exist locally and beyond, and to identify a range of potential facilities and their function for the proposed FEC (Butler et al., 2001).

The third and final study incorporated the findings of the two prior studies to design possible facilities and create a site plan for the proposed SDSF FEC (Butler and Hester, 2001). See the FEC section of the Demonstration and Education Chapter for additional information.

TIMBER INVENTORY AND GROWTH/SUSTAINED-YIELD ANALYSIS

The first forest-wide timber inventory and supplemental growth analysis were conducted to determine the current conditions of forest volume and vigor. The timber inventory, completed in 1991 by SDSF staff, surveyed both conifers and hardwoods on a ten percent sample of the Forest (Lee, 1991). The growth study used a portion of the plots established by the timber inventory but assessed the current growth rates of conifers only.

The second forest-wide forest resources inventory was completed in 2006. This forest inventory surveyed all tree species, conifers and hardwoods on a two percent sample of the Forest. The inventory data were used as input to the FORSEE growth model, which along with harvest scheduling software, projected the growth and development of the Forest over a 100-year planning interval. This analysis formed the basis for determining sustainable harvest levels in the next five to ten years that are also achievable in the long term without degrading the productivity of soils and the ecosystem. The results of this analysis are documented in the Option A Plan (CAL FIRE, 2010).

In 2011 Mike Papa, a Forestry Science master's student from California Polytechnic State University at San Luis Obispo, completed his thesis titled "Effects of Silviculture Management on Coast Redwood Forest Composition, Density, and Structure in Santa Cruz and San Mateo Counties". Papa's research included data from the Amaya and Longridge THPs at SDSF to examine forest restoration management (Papa, 2011).

WATERSHED ASSESSMENT

Dr. Brook Kraeger, a consulting hydrologist and neighbor of SDSF, has recorded rainfall and stream flow data from gauges in a 159-acre tributary to Soquel Creek in the Forest since the late 1990s. The intent of this long-term monitoring is to evaluate hydrologic modeling and to examine the impact of timber harvesting on the hydrologic process. Kraeger provides raw data to SDSF that is used in annual reports, the Steelhead Trout Population Survey Reports and the Instream Temperature Monitoring reports (see Figure 6, page 47). This work is also supported by a Campbell Scientific CR1000 weather station that was established at the lower helipad

in 2013.

In 1993, a comprehensive cumulative watershed effects analysis for the East Branch of Soquel Creek watershed was completed by CAL FIRE Hydrologist Pete Cafferata (Cafferata and Poole, 1993). Chris Poole, a student intern from the University of California, Santa Cruz, and Forest staff assisted with data collection. Several subsequent Soquel Creek Watershed studies and surveys have been done by CDFW, the Resource Conservation District of Santa Cruz County, the National Marine Fisheries Service, Natural Resources Conservation Service, and others. Information from this research guides management activities to maintain or enhance the watershed integrity. See the Watershed Assessment Chapter for additional information.

A study on the movement of large woody debris in Soquel Creek was completed in 2003 (Lassettre and Kondolf, 2003). Steve Reynolds, California Geological Survey, completed the 2013 Soquel Creek Large Woody Debris As-Built Report for Site 1 of the LWD project (Reynolds, 2013). Reynolds also provided a report after completion of the four sites in 2012 and 2013, and offers a comparison of large wood loading rates in enhanced stream reaches to those in other minimally disturbed watersheds (Reynolds, 2013). Based on the work done in 2012 and 2013, this report contains findings and recommendations for future restoration work.

FISHERIES ASSESSMENTS

One formal and comprehensive aquatic habitat survey was conducted on SDSF in 1994 (Berlekamp, 1994). A draft fisheries management plan was developed by Forestry Aide Bronwen Berlekamp and former Forest Manager Thomas Sutfin in 1995 (Berlekamp and Sutfin, 1995).

A macroinvertebrate sampling study performed by SDSF staff was conducted in October of 1995. DFG analyzed the collected samples and provided results of their findings in May 1996 (CDFG, 1996).

Annual fish population surveys were conducted on SDSF in cooperation with the California Department of Fish and Wildlife from 1993 to 2001 at four separate sites. From 2002 until the present, SDSF has continued these annual surveys in cooperation with the NOAA Southwest Fisheries Science Center (Sogard et al., 2009). NOAA Fisheries added a fifth site, increased the monitoring frequency, and incorporated additional research components. These Steelhead Trout Population Survey Reports have been completed every year to the present, with the exception of the year 2000 (CAL FIRE, 1993-1999, 2001-2012).

In-stream temperature monitoring has been conducted most years since 1997 using HOBO data loggers at seven sites in the forest (CAL FIRE, 1997-1999, 2001-2003, 2005-2012). Results of this monitoring have been compiled including seven-day moving averages which can provide significant information about the chances for fish

success because these calculations reflect the duration of high water temperatures.

FOREST HEALTH

Since the beginning of the Sudden Oak Death (*Phytophthora ramorum*) outbreak in the Santa Cruz Mountains in the late 1990s, SDSF staff has worked closely with U.C. Cooperative Extension specialists and researchers from the U.C. Berkeley Forest Pathology and Mycology Laboratory lead by Dr. Matteo Garbelotto (Garbelotto, 2013). Several Sudden Oak Death (SOD) studies have been completed over the years on SDSF including early work on various treatment options using direct chemical control with phosphonates and evaluations of the role of bay-laurel as a SOD vector (Hayden et al., 2011). In 2001, a transect/plot system was established for long-term monitoring of the occurrence and spread of SOD over time (McPherson, no date).

A biological control study of California bay-laurel resprouts began in 2013 by Dr. Marianne Elliott from Washington State University to test new strategies for managing the spread of SOD (Elliott et al., 2012). The aim of the study is to develop and evaluate the effectiveness of treatment and eradication strategies to suppress bay-laurel resprouts which can harbor SOD (Elliott, 2013).

A graduate student from the University of North Texas, Alicia Gray, completed a Master's thesis on Sudden Oak Death host species in SDSF in May 2014 (Gray, 2014).

ATMOSPHERIC STUDY

Researchers from the University of North Texas, Department of Geography, led by Dr. Alexandra Ponette-Gonzalez, began a study in 2012 to estimate atmospheric deposition into the Santa Cruz mountains using through-fall measurements and airborne LiDAR. Rainwater was collected in a rain gauge and throughfall was collected in open funnel collectors in grasslands and forest stands of Douglas-fir to measure variation of chemical levels for future deposition modeling. Results from this study were provided in a Master's thesis by Kereen Griffith (Griffith, 2014).

RESEARCH PRIORITIES

Subjects for research will include all ecosystems in the Forest. Studies will concentrate on all components of resource management, including protection, management, and public use. Current research priorities are listed below; as forest management techniques evolve, precedence will adjust accordingly.

WATERSHED MANAGEMENT

- * Monitor the effects of forest management activities on the resources of the East Branch of Soquel Creek watershed within SDSF.

- * Investigate stream enhancement and rehabilitation techniques.
- * Investigate and document effective techniques to minimize erosion and stream sedimentation caused by logging, road building, and recreational use.
- * Monitor stream discharge and precipitation in small headwater basins.

TIMBER MANAGEMENT AND FOREST HEALTH

- * Investigate optimal spacing requirements for growth and regeneration.
- * Study hardwood management and Sudden Oak Death.
- * Study old-growth redwood forest communities and ways to achieve late-succession stand characteristics over time.
- * Investigate logging techniques which minimize visual, auditory, and environmental impacts.
- * Investigate silvicultural methods with varied harvest levels and cutting cycles.

FISHERIES AND WILDLIFE

- * Assess the current condition of the fisheries resources within SDSF and document long-term trends.
- * Study methods to improve steelhead and coho rearing habitat conditions.
- * Investigate methods to improve wildlife habitat and provide for healthy biodiversity.

RECREATIONAL USE

- * Monitor environmental impacts of visitors to the Forest.
- * Study the reactions and responses of recreational users and neighbors to all forest management activities.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

The general public's wishes regarding research include desire to be kept informed of SDSF's research plans and actions, both presently and in the future. Specifically, curiosity relating to research subjects, objectives, and investigators (and their affiliation) has been prevalent. To satisfy this concern, SDSF plans to announce current research projects through publications such as the Mountain Network News and other local newspapers. Additionally, newsletters containing information on present activities will be posted on signboards throughout the Forest.

Other public comments have suggested that information derived from research studies should be used to formulate forest management policy and actions. Gaining information which will aid in the management of SDSF is essentially the purpose of research projects; the Forest staff intends to put to use relevant information obtained through research.

A final category of public interest involves the availability of study results. Individuals have

expressed that research findings be made available as conclusions that may be applicable to their own lands. As stated in the management guidelines below, results will be available through public libraries, natural resource journals, CAL FIRE publications, the CAL FIRE web site, and direct mailing.

MANAGEMENT GUIDELINES

1. Actively design and carry out continuing scientific studies which refine and improve upon existing state of the art forest land management techniques.
2. Coordinate research projects with other State Forests and local, state, and federal public agencies. Additionally, research opportunities have been and will continue to be provided for universities, industry professionals, and private interest groups. Research may be formal or informal, depending on the party conducting the study and available funding.
3. Assure dissemination of research results in a timely and professional manner. Information gained from studies will be made available to local, state, and federal public agencies as well as resource professionals, forest neighbors, and other interested individuals. Reports may be made available through direct mailing, newsletter articles, public libraries, professional natural resource journals, and the CAL FIRE web site.

PLANNED ACTIONS

1. Create a list of priority research needs, identify proper audiences, and define techniques to distribute information effectively and in a timely manner. Encourage innovative research in forest management, resource protection, and recreation. Investigate previous research to determine the extent of what has already been completed in the area as well as topics lacking in information.
2. Serve as an outdoor laboratory for CAL FIRE research projects and encourage investigations by other agencies and educational institutions. Use study results to improve forest practices both in the Forest and statewide.

CHAPTER 11: RECREATION

INTRODUCTION

The legislation creating State Forests (Public Resources Code 4631-4664) and Board of Forestry and Fire Protection policy both state that recreation is to be an integral part of the Demonstration State Forest Program. In addition, SDSF's enabling legislation, AB 1965, states that public enjoyment and open access are to be provided. SDSF's recreational management goal is to provide for uses that are compatible and integrated with resource protection, public education, and forest management while following the guidelines of AB 1965 and Board policy.

Prior to state acquisition of SDSF in 1988, the property was privately owned and used primarily by the owners, their guests, neighbors bordering the property, and frequent trespassers. Users consisted of equestrians, motorcycle riders, four-wheel drive enthusiasts, hunters, target shooters, and a limited number of hikers, mountain bikers, and campers.

Since the establishment of SDSF, recreational use of the Forest has changed. The primary groups who utilize the Forest are (in order of use) mountain bikers, hikers and walkers, and equestrians. The majority of recreational users at SDSF come for the mountain biking. SDSF offers a unique experience with several miles of single-track trails and a lengthy downhill decent from the ridge to Hihn's Mill Road. Other recreational users include picnickers, mushroom gatherers, bird watchers, trail runners, dog walkers, environmental organization members, dog search and rescue training personnel, geo-caching individuals, and people coming to enjoy the forest environment.

RULES, REGULATIONS AND POLICIES

Public access is allowed during daylight hours. Campfires, fishing, hunting, commercial events, and the use of firearms and motor vehicles are generally not allowed.

Current forest regulations prohibit all recreational motor vehicle use. The primary grounds for this regulation are lack of infrastructure (parking areas, barricades to prohibit vehicles driving onto sensitive areas), lack of SDSF personnel to provide patrol, and security and maintenance costs associated with increased traffic on unimproved roads. Other reasons include resource protection, safety, and fire prevention.

ILLEGAL ACTIVITIES

The enforcement of regulations has decreased the number of motorcyclists to a few violators, and four-wheel drive use is now almost non-existent.

In the past illegal shooting at Rattlesnake Gulch, private property across Highland Way from

the Forest, was a detriment to Forest visitors. Gunfire could be heard throughout the Forest and many users found this unnerving. Additionally, the noise and danger of gunfire reduced the quality of forest visitor experiences. Since then, this problem has been solved and is no longer an issue.

Vehicles parked along Highland Way and occasionally in the parking area have been burglarized. Recreation users are warned to not leave valuable in their vehicle and especially not in plain sight. Users are encouraged to use the 911 system to report illegal activities and activate law enforcement response as needed.

Illegal parties at the Highland Way parking lot have been an ongoing problem. CAL FIRE law enforcement, Santa Cruz County Sheriff's deputies and State Parks Rangers have worked together to respond to these parties and have been effective in deterring them through a coordinated effort. Unfortunately, the effects of law enforcement action on illegal party organizers do not deter them indefinitely. Illegal parties pop up in new locations and back in old locations after a period of time. Additional security measures have been employed to prevent these illegal parties and are coordinated through law enforcement officers. Neighbors are encouraged to use the 911 system to report this activity and to activate a law enforcement response.

Other illegal activities that have been ongoing involve illegal harvesting of trees (particularly along property boundaries), illegal trail building, and dumping of trash on the Forest. All of these activities are investigated and responded to by SDSF staff in coordination with law enforcement.

RECREATION STUDY AND PLAN

To gain a better idea of current and future recreational users, their views, and related issues, SDSF implemented a recreation study which obtained information from the community and current users. The study, titled Soquel Demonstration State Forest Recreation Study Final Report (McNally and Hester, 1993), was intended to be the primary source of citizen involvement in the recreational planning process and designed to develop recreational and forest management strategies for SDSF. The results provided baseline information about existing and potential recreational users that will be utilized as more detailed recreation planning occurs.

The recreation study involved mailing questionnaires to all individuals who share a boundary with the Forest or live nearby. Additionally, different user groups who frequent the Forest were asked to fill out surveys while visiting SDSF. Information about the Recreation Study and the questionnaire were also published in the Mountain Network News, the Summit community's local newspaper. Of the 6,600 individual questionnaires distributed, 800 were returned. Several common themes or issues of concern were determined through this process and are discussed in this chapter.

The draft Education and Recreation Master Plan was completed in 1996. Consistent with this plan, several actions have been completed. A campground has been developed and is used

on a limited basis by researchers, educational groups and personnel working on projects in the Forest. Interpretive sign boards have been placed in project areas, new trails have been developed, trail maintenance occurs on a regular basis, a portable toilet has been made available, picnic tables have been added and additional signage has been placed.

The study and plan that were prepared in the 1990s were predicated on securing new and better access on the west end of the Forest. Both plans describe the lack of adequate public access as the biggest challenge to overcome in managing recreation and education programs. As new access continues to be evaluated and negotiated, several new issues related to recreation have developed. In the 20 years since the Recreation Study was completed, new partnerships have developed, recreational activities have changed considerably, and best management practices have evolved. For example, advancements in battery technology has led to the development of electric-assist mountain bikes which look and perform similarly to non-motorized mountain bikes, blurring the lines of what is defined as a motorcycle. Also, the development of high-powered LED light systems has brought illegal night-time recreational use to many nearby trail systems. Management strategies will be reviewed in light of these changes.

PARTNERSHIPS

SDSF staff has fostered the development of groups to support recreation and education goals and to conduct trail patrols. Three groups have partnered with CALFIRE in this way and dedicated volunteers donate hundreds of hours of labor annually.

The Stewards of Soquel Forest is a non-profit organization founded in 2000 that assists with the recreation and education goals of SDSF. The Stewards coordinate volunteers for trail maintenance work and lead trail work days every year. They also assist with other projects such as steelhead trout monitoring, and public demonstration and education events.

A local chapter of the International Mountain Bicycling Association (IMBA), Silicon Valley Mountain Bikers (formerly named Responsible Organized Mountain Pedalers), has sponsored educational events and organized volunteers to do trail patrol at SDSF through the National Mountain Bike Patrol program.

Another local IMBA chapter, Mountain Bikers of Santa Cruz, has taken a leadership role in developing a plan for a new trail on the Forest. They have partnered with the Stewards of Soquel Forest and developed a trail crew leader training program to build a cadre of volunteer crew leaders to help build the new trail. They are leading a fundraising effort and donating their staff time to train, plan and implement the trail building effort.

The relationships with these organizations have been valuable to SDSF, allowing growth and improvements in the trail system and facilitating recreational use while minimizing CALFIRE staff time and reducing costs of managing the trail system. Through improved communication and by working together, trail issues and recreational user issues are continually being resolved.

REGIONAL CONTEXT

SDSF's location is centered between the population centers of the Silicon Valley and the greater Santa Cruz metropolitan area, which provides for day use by many outdoor enthusiasts. SDSF is also a key link in a contiguous greenbelt between Los Gatos and Soquel which has been identified as an important region within the Land Trust of Santa Cruz County's Conservation Blueprint Initiative (2011).

SDSF's recreational opportunities exist alongside and are connected with those provided by other nearby trail systems and facilities, some of which are described below.

THE FOREST OF NISENE MARKS STATE PARK (TFNMSP)

Sharing a common boundary with TFNMSP is positive and offers prime recreational opportunities for both facilities. Recreationists, particularly mountain bikers and hikers, regularly use both facilities in one outing. Reaching the summit of Santa Rosalia Mountain and the entrance to SDSF by way of Aptos Creek Fire Road is a significant athletic feat requiring over 2500 feet of elevation gain.

Some people think the Forest is part of TFNMSP. Management staffs of both facilities have a good working relationship with one another; they meet regularly to share information and solve common problems, strengthening their relationship. As a result of this alliance, the new State Park map has been updated to include SDSF and to reflect the connections of the Ridge Trail in SDSF with the Aptos Creek Fire Road located on TFNMSP.

SIERRA AZUL & UVAS OPEN SPACE PRESERVES

The Midpeninsula Regional Open Space District is planning public access to the Rattlesnake Gulch parcel of the Sierra Azul Open Space Preserve, potentially with a hike-in campground on a knoll overlooking SDSF. The Open Space Master Plan is still being developed and revised with ongoing public comment.

The Santa Clara County Open Space Authority has acquired land to the east of Rattlesnake Gulch which connects to Santa Clara County's Uvas Canyon County Park. Additional acquisitions are being pursued by the OSA in this region.

BAY AREA RIDGE TRAIL

Through the development of additional public access in Sierra Azul there is a potential for a link from SDSF to the Bay Area Ridge Trail, which is the longest recreational trail in the region. When completed, the Bay Area Ridge Trail will connect Sierra Azul Open Space Preserve to Mt. Madonna County Park. A potential trail connection from SDSF to the Bay Area Ridge Trail would likely bring additional visitors to the Forest, who may make use of the Hihn's Mill Road parking area or roadside parking along Highland Way.

BAY TO BAY TRAIL

Public support for a long distance regional trail connecting the San Francisco Bay to the Monterey Bay began to develop in 2001. This trail concept is a route from the Bay Trail in Alviso, up along the Guadalupe River Trail in San Jose, through Sierra Azul Open Space Preserve, crossing the Bay Area Ridge Trail near its highest point, then down into SDSF to TFMSP, and ending at New Brighton State Beach.

CAMP LOMA

Camp Loma is a recreational facility operated by Santa Cruz County Youth Activities. Facilities include a large covered dining area, commercial-style kitchen, lawns, pool, showers, campsites and other amenities. It is available for rental and several groups have made use of the facilities to support events on SDSF.

ADAMS RANCH CONSERVATION EASEMENT

The Land Trust of Santa Cruz County owns a conservation easement over a parcel between Longridge Road and Adams Road, which includes an existing trail and language which provides for its eventual connection to the Fern Gulch area of SDSF. Other landowners in this neighborhood have expressed interest in providing easements to improve connectivity to SDSF.

PEDESTRIAN & CYCLING ROUTES

A roadside pathway along Summit Road between Highway 17 and Soquel-San Jose Road provides a separated environment for pedestrians and cyclists of the surrounding community. Soquel-San Jose Road has recently been designated as a cycling route by Santa Cruz County. “Sharrows” have been painted on the roadway and signs installed to encourage safe sharing of the road. Both of these facilities have the potential to bring recreational users to SDSF in an auto-free manner should trail connections to the Forest be completed.

NEIGHBORING COMMUNITIES

Neighbors of the Forest have a range of opinions and concerns regarding recreational use. Some neighbors who live nearby want to assure they can continue to access, or to gain access to the Forest directly from their neighborhood without encouraging general public use of narrow local roads. Alternative access points would require careful planning and management to address concerns.

Some neighbors are concerned that recreation and timber harvesting will disrupt their quality of life in the mountain community. They are worried that too many people will come from the larger metropolitan areas and impact their roads, increase fire danger, and vandalize property. Some individuals have expressed concern regarding excessive noise and their views being destroyed by timber harvesting.

Some trespassing through private property to enter and leave the Forest has been reported. Forest patrols continue and citations to trespassers have been issued. Publishers of trail guides have been contacted to correct errors and to promote an accurate depiction of public access routes. SDSF maps provided to the public do not show private roads adjacent to the Forest.

Every effort is made to consider the impact of proposed management activities on surrounding neighbors. SDSF works with the community to aid in their understanding of SDSF as public land with certain legislated mandates that include education, public use, and timber management.

Public meetings and tours are conducted to discuss SDSF mandates and hear public concerns. Access issues are being investigated, and detailed recreation and timber management planning have been completed. Motor vehicle use within the Forest is restricted, limiting the types and levels of recreational use.

PUBLIC ACCESS AND STAGING AREAS

Public access into the Forest continues to be a significant challenge. At this time, only one remote, legal access point provides a staging area with parking. This entry, the main entrance to SDSF, consists of an easement across private property and is subject to winter closures due to continuing landslides on Highland Way. Furthermore, those who do not use trails (e.g., physically challenged individuals) are limited to using the edge of the Forest because there is no public access by motor vehicles into the internal areas of SDSF.

Providing alternative access requires a complete analysis of access needs and an assessment of land available for easement or acquisition. Accommodating non-trail users should be a consideration in future recreation and education planning. Developing an access plan to address this issue is a high priority.

EXISTING FACILITIES

SIGNS AND MAPS

Directional signs with mileage have been installed at all trail intersections and at the entrance to the Forest. Large signs at each entrance have also been placed listing forest rules and regulations that are enforced. Sign boards are located at three locations on SDSF: the Highland Way entrance, Badger Spring, and the Ridge Trail entrance from The Forest of Nisene Marks State Park. The sign boards display large Forest maps and are updated periodically with educational materials, announcements, and public notices regarding rules and closures. Free brochures which include maps are also provided.

MULTI-USE ROAD AND TRAIL SYSTEM

The trail system of SDSF consists mostly of old logging roads that have been reopened, along with some single-track trail segments (Figure 19). Currently, a few of the trails are not

completely contained within the State Forest where they meander along the Forest boundary. Some trails lead from the State Forest to the State Park and others end at private land.

Loop trails keep visitors inside the Forest and direct them back through the State Park or to Highland Way, instead of onto private property. Most visitors use the interconnected trail system on the south side of the creek, typically descending the Ridge Trail and climbing Hihn's Mill Road and connecting between the two using the single-track trails to loop between them. This natural circulation pattern results in many trails being used in a de-facto one-way direction.

Trails are shared by all recreational users, except for one half-mile of Ridge Trail between Corral Trail and TFNMSP. Horses are not allowed on this upper section to keep them from entering the Park, as they are not permitted due to deed restrictions.

SDSF does not suffer the problems and conflicts experienced on other public lands that allow hiking, horseback riding, and mountain biking on the same trails. The remoteness of the entrance road, the length of Hihn's Mill Road that must be traversed before reaching the first trail junction and the steep and rough conditions of the single-track trails seem to filter out casual trail users who seek short and easy trail experiences. Consequently, most of the Forest users are more experienced and adventurous than typical park visitors. The main trails along the southern portion of the Forest are used extensively by mountain bikers, while the spur trails on the north and western side of Soquel Creek are used much less, mainly by neighbors of SDSF.


Forest staff have worked to effectively manage the trail system and to monitor the satisfaction of all user groups. Forest trails are open to all user groups and trail etiquette is stressed to Forest visitors. A Trail Use and Safety Guide has been developed to foster cooperative use and reduce trespass problems.

ROAD AND TRAIL DESCRIPTIONS

Hihn's Mill Road – 5.6 miles

Hihn's Mill Road is the main road through the Forest, used by nearly all recreational visitors as well as SDSF staff, emergency vehicles, timber operators, researchers and others. This rocky, all-season road traverses the length of the Forest. From Highland Way it crosses Soquel Creek at a bridge, passes through the main parking lot and staging area, and traverses private property for a half mile before entering SDSF at the eastern boundary. It descends over 1000 feet in elevation for 4 miles, passing the

Figure 19. Transportation System of SDSF.



bottom of each of the single-track trails before crossing Soquel Creek again. From the lower bridge it then climbs up past the intersection of Amaya Creek Road, and continues to the gate at Olive Springs Quarry. Public use is restricted by an administrative closure past Amaya Creek Road to limit trespass into the hazardous mining area.

Ridge Trail - 3.3 miles

Ridge Trail leads from TFNMSP at the southeast corner of the State Forest and follows Santa Rosalia Ridge to Sawpit Trail. It meanders along the common boundary between the State Forest and TFNMSP. This trail is the only other authorized public access route to the Forest aside from the main entrance at Highland Way.

Corral Road/Trail - 1.7 miles

Corral Trail starts at Sulphur Springs Trail and connects with Ridge Trail toward the southeast end of the Forest. It is named for an old corral, less than half way up the trail that was used by the Sulphur Springs Resort to house their livestock. The top 0.3 miles consists of a single-track trail.

Sulphur Springs Road - 1.5 miles

This trail is a road that starts on Hihn's Mill Road and goes up to Ridge Trail. It was constructed in the 1870s to access the Sulphur Springs Resort. The trail passes remnant cold springs, but all of the old resort buildings have been destroyed or removed. This route also serves as an emergency vehicular access to the lower helipad near Hihn's Mill Road and the upper helipad on Santa Rosalia Ridge.

Braille Trail – 1.5 miles

This is the most recently developed trail in the Forest. Connecting Hihn's Mill Road with Ridge Trail, Braille Trail is a good example of the extraordinary relationship that has grown between SDSF staff and the public. This trail began as an unsanctioned trail in the late 1990's. It was originally constructed by mountain bikers at night where they found their way "by Braille". Its route and construction methods were found to cause adverse impacts to water quality and unnecessary risks to public safety. Public support for correcting the inappropriate trail features led to an alliance between SDSF and recreational trail advocates and began formation of the Stewards of Soquel Forest. A compromise solution led to the challenging but environmentally sustainable trail alignment in place today that is now a sanctioned trail. Many of the volunteers involved in realigning Braille Trail have continued their association with SDSF to construct and maintain highly desirable and environmentally considerate trails.

Tractor Road/Trail - 1.5 miles

This trail was constructed in 1934 when logging methods changed from steam donkey to crawler tractor. It was the first logging road developed to access the slopes of Santa Rosalia Ridge. It starts at Hihn's Mill Road and ends at Ridge Trail.

Flow Trail (planned) - 4.5 miles

In 2013 a planning effort began to develop a new type of trail called a flow trail at SDSF, following the reopening of Tractor Trail previously used as a logging road during the Rim THP. This effort is sponsored by the Mountain Bikers of Santa Cruz. They have provided funding and expertise for planning and implementation. The flow trail is designed to be used by any skill-level bicyclist and emphasizes a low gradient with grade reversals that utilize the rider's momentum to minimize pedaling and braking. Since all the other single-track trails at SDSF are more appropriate for highly skilled, experienced bicyclists, the new flow trail will allow for a more diverse recreational experience. This trail will be appropriate for families with children and beginner mountain bikers.

Sawpit Trail - 1.2 miles

This trail starts at Hihn's Mill Road and connects with Ridge Trail. It is located one-half mile east of the picnic area at Badger Spring. There is a sawpit along the trail that was used before the turn of the century for cutting logs into lumber.

Longridge Road - 1.5 miles

This trail is a road that starts on Hihn's Mill Road and travels to the county's Longridge Road. The last 0.1 mile is on private property. This trail is primarily used by neighbors.

Amaya Basin Road – 0.7 mile

This trail is a road that begins midway along Longridge Road and extends along the eastern side of Amaya Creek. It ends at a landing last used for the Longridge THP.

Fern Gulch Spur – 0.2 mile

This trail is a logging road constructed in 2011 for the Fern Gulch THP and ends at a landing used for those harvest operations.

Amaya Creek Road – 2.0 miles

This road starts on Hihn's Mill Road, about 0.5 mile west of the bridge over Soquel Creek. It climbs up to the ridge separating Amaya Creek from Hester Creek and intersects with Comstock Mill Road near the Forest boundary.

Amaya Pond Trail – 0.5 mile

This trail is a road that starts about half way along Amaya Creek Road. It passes near Amaya Pond, and then descends down in a sinuous manner towards Amaya Creek, ending at a landing.

Amaya Bridge Trail (planned)

In 2003 planning began for this trail that would connect Amaya Pond Trail with Longridge Spur Trail, providing a loop on the north side of the Forest. Complex issues arose in designing the bridge needed to cross Amaya Creek and planning was put on hold until more resources could be provided.

Comstock Mill Road 0.7 mile

Public use of Comstock Mill Road is currently restricted to prevent trespass onto private property.

High Bridge Spur Trail (inactive) - 0.3 mile

This trail is the original route of Hihn's Mill Road into the Forest. The bridge was lost to fire in the 1970's but the road prism remains and was re-established as a logging road with temporary bridges for the neighboring CHY Olive Springs THP. This route is in the area closed administratively to prevent trespass at Olive Springs Quarry.

OTHER FACILITIES

Picnic Areas

The Badger Spring picnic site was the first such area in the Forest and was established long before the State assumed management. This area is very scenic, but relatively remote. Badger Spring is located along Hihn's Mill Road in the alluvial flats of the East Branch of Soquel Creek. The site has several picnic tables and a few hitching posts for horses. On the opposite side of the road is the Forest's ten-acre old-growth redwood grove.

Other picnic area locations are the Forest entrance, Sulphur Springs, the lower helipad area and on Ridge Trail between Braille Trail and Tractor Trail.

Portable Toilet

There is one portable toilet in the Forest, provided and maintained by a generous donation from a mountain bike manufacturer. It is located by the eastern property gate on Hihn's Mill Road, near the Forest entrance.

Parking Area

There is one parking area located between the two gates at the Highland entrance. This area is not in SDSF, but the State has a public-use easement with the owners, Roger and Michelle Burch. The parking area has been rocked and is generally accessible to the public year round. Work done in or around the parking area to maintain erosion control, reduce fire hazard, post new signage or change access through the common gates is coordinated with Burch family representatives. Recent upgrades include additional rocking, installation of new signage, redecking of the bridge over Soquel Creek and the installation of a steel gate on the bridge that replaces the old chain gate. SDSF staff will continue to coordinate with the Burch family representatives to determine the need to close the parking area due to a range of issues including saturated soil, illegal parties, high fire danger or timber operations.

Campground

There is one campground that is used by special permit on a limited basis by educational groups, researchers and personnel working on projects on the Forest.

RECREATION PROGRAMS

ORGANIZED EVENTS

Organized events are allowed through a fee-based Special Use Permit. Special Use Permits may allow uses which are otherwise restricted, such as use of motor vehicles and camping. These events are limited to the number that can be managed by the SDSF staff and are evaluated on a case by case basis.

FORESTRY EDUCATION

There is an encouraging level of interest in the Forest's educational potential. Local teachers are interested in bringing classes out to the Forest and local residents like the idea of educating the general public about the environment. Additional Forestry Education programs and interpretive information should be provided for recreational users. Outreach programs have been initiated with local schools and natural history organizations. Interpretive signs have been installed on the Forest. SDSF staff members offer verbal educational information to the public, both in the office and in the field.

FUTURE RECREATION

Additional opportunities are available to provide a more diverse recreational experience at SDSF. This includes the potential development of new facilities such as a Forestry Education Center and new uses such as an ADA-accessible trail, an interpretive trail, or possibly a disc golf course. Members of the public have expressed a desire for these types of opportunities at SDSF, as well as a larger and more interconnected trail system. Each of these possibilities needs further investigation and planning in order to evaluate the feasibility and compatibility with the goals and mandates of SDSF.

As progress is made toward securing new access for a Forestry Education Center and new facilities, a recreation and trails plan should be developed. The new recreation and trails plan will require careful consideration as to how changes in access and use patterns will affect the other areas of SDSF's mission, such as education, research, watershed protection, and demonstration of sustainable forestry.

SDSF is one of the few publicly-owned lands in Santa Cruz County that has the ability to provide a public hunting opportunity and may elect to do so in the future in conjunction with CDFW. Given existing constraints of access to the Forest, limited availability of CDFW and CAL FIRE staff time, the Santa Cruz County 'no shoot' zone, and multiple uses occurring on the Forest, a pilot program may initially be implemented to determine long-term feasibility of a hunting program. Game species may include, but not be limited to, deer, wild pig, band-tailed pigeon, California quail, and/or wild turkey. SDSF staff will work with local CDFW wildlife biologists and statewide program staff as necessary to determine appropriate levels of harvest in order to maintain population viability and to schedule special hunt events. Any hunting program which may be established for a particular species will be consistent with the appropriate CDFW statewide Environmental Impact Report or equivalent Strategic Plan developed by CDFW and will comply with all State and Federal laws pertaining to sport

hunting.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Many comments and opinions were gathered during the Recreation Study. Appendix A of the study's final report contains all of the results of the questionnaire as well as comments and concerns of users and neighbors. For detailed comments, refer to Soquel Demonstration State Forest Recreation Study Final Report (McNally and Hester, 1993).

MANAGEMENT GUIDELINES

1. Public safety and protection of the natural resources of the Forest will be emphasized in all patrol and management planning activities. Violators of Forest regulations will be cited.
2. Certain activities may be prohibited, and certain areas administratively closed by order of the Forest Manager. Such restrictions will be posted at all entrances and sign boards on the Forest. Special Use Permits may be granted at the discretion of the Forest Manager to allow otherwise unauthorized uses on a case-by-case basis. Restricted activities include use of recreational motor vehicles, camping, fires, entering closed areas, gathering mushrooms without a permit, use after sunset and before sunrise, large public gatherings, organized races, commercial events, shooting, hunting and fishing (per CDFW Regulations).
3. In accordance with Board of Forestry and Fire Protection policy, recreational facilities will be maintained with minimal development, preserving the rustic and informal characteristics of the Forest. Periodic assessments will be made to ensure that facilities meet users' needs while remaining as natural as possible.
4. Regular maintenance will be provided to ensure the upkeep and safe conditions of all existing facilities, including picnic tables, signboards, parking areas, and trails. Periodic inspection and maintenance of recreational facilities will be performed by Forest staff, Ben Lomond Conservation Camp crews, and volunteers.
5. Recreation will be coordinated to achieve integration and compatibility with timber management, resource protection, demonstration and education, and the neighboring community. Demonstrations will show how recreation and timber harvesting, two seemingly conflicting management objectives, can be integrated. Forestry education will be an integral part of the recreation experience. Interpretive resources will explain the basics of forest management, current research on the Forest, and the need to protect sensitive resources. In addition, visitors will be directed away from private property through circulation patterns (loop trails), use of proper signage (directional, regulation, no trespassing), and keeping Forest attractions away from private property boundaries.

6. SDSF will be managed to provide positive experiences for all recreational users. Proper trail etiquette between mountain bikers, equestrians, and hikers will be encouraged through signs, educational information, and patrol by staff and volunteers.
7. Management will strive to minimize recreational development and recreational use impacts in ecologically sensitive areas such as riparian zones. Any recreational improvements in riparian areas will be based on careful site-specific evaluations.

PLANNED ACTIONS

1. Record and compile descriptions of all reported violations or nuisances caused by public users at SDSF or on adjacent ownerships including, but not limited to, trespass, vandalism, littering, and noise. Implement restrictions on public use if needed.
2. Conduct ongoing patrols of riparian areas including those with existing facilities to enforce prohibitions on vandalism and other damage to riparian habitat related to public use. Implement restrictions on public use if needed.
3. Evaluate and grant requests for recreational Special Use Permits on a case-by-case basis.
4. Compile annual estimates of public use of SDSF in user days, using patroller reports, surveys, trail counters, and other information.
5. Continue to support, and work to expand volunteer programs to enhance recreation, interpretation and patrol.
6. Provide input to neighboring land managers to facilitate interconnected trail systems and regional trail development.
7. Develop an access plan, and acquire land or easements necessary to provide better public access to the Forest via Soquel-San Jose Road.
8. Evaluate the need for a new Recreation and Trails Master Plan for SDSF to improve the trail network and recreation opportunities in the Forest.

CHAPTER 12: RESOURCE PROTECTION

FIRE PROTECTION

The CAL FIRE San Mateo-Santa Cruz Unit Chief is responsible for fire protection in the State Forest. The Forest Manager, the Unit's Operations Division Chief for Santa Cruz County, and the local CAL FIRE Battalion Chief will work together to ensure an adequate fire protection program is in place for SDSF. In addition, the Forest staff will work with other agencies as needed to provide fire protection for the Forest.

FIRE HISTORY

The fire return interval for the area that is now SDSF in the era before fire suppression was roughly ten to twenty years. These fires were caused by lightning or set by Native Americans to manage vegetation.

A fire that occurred in 1903 spread from the east to Santa Rosalia Ridge down to the Sulphur Springs area (Powell, n.d. pg. 161). Another fire started in 1922 in Hinckley Creek and spread to Soquel Creek, burning a total of 7,000 acres.

A 50-year fire history (1929-1979) of the Santa Cruz Mountains was compiled by Jason Greenlee in 1981. The area which is now SDSF was included in the study, and the following fires occurred during that 50-year period:

- 1933 - 240 acres burned in the Amaya Creek drainage.
- 1936 - 54 acres burned along Santa Rosalia Ridge between Tractor Trail and Sawpit Trail.
- 1936 - 118 acres burned in the Hinckley Creek basin, with approximately 25 acres in the Forest, along the ridgeline at the top of Sulphur Springs Trail.
- 1938 - 87 acres burned along the upper portion of Sulphur Springs Trail.
- 1957 - 168 acres burned from Highland Way to Hihn's Mill Road, just east of Ashbury Gulch. Approximately 50 acres were in the Forest.

The exact causes of these fires are not known, but it is believed that logging practices of the past, coupled with high risk machinery and equipment, caused most of them. Phil Mason, a long-time local resident, remembers that the 1933 fire was caused by a steam donkey engine that caught fire, burning the Amaya Creek drainage.

Modern logging methods and equipment are much safer from a fire perspective, and the risk

of fire has been greatly reduced. This is evident from the lack of fires in the Forest during the last 30 years. The only known recent fire occurred in 1970 when the High Bridge, at the southwest edge of the Forest, was set on fire.


Recent large fires in the area (Figure 20) include the Lexington Fire in July, 1985; the Summit Fire in May, 2008; and the Loma Fire in October, 2009. Though not in the State Forest, the Lexington Fire's impact to the Summit mountain community was severe enough to mention here. An arsonist started the fire which burned 13,800 acres in the Los Gatos Creek Drainage (600 acres were in Santa Cruz County). Forty-five hundred people were evacuated from their homes, including residents of the Summit area. In the end, 44 structures were destroyed, including houses and outbuildings.

The Summit Fire burned 4,270 acres in the upper portion of the Soquel and Corralitos Creek watersheds from May 22–27, 2008. In total, 1,130 acres burned in the Soquel Creek watershed (34 percent high burn severity, 43 percent moderate, 20 percent low, and three percent unchanged). Thirty-five residences and 64 outbuildings were destroyed.

The Loma Fire burned 485 acres within the Soquel Creek watershed between October 25 and November 2, 2009. One trailer and two outbuildings were destroyed. The fire started on Loma Prieta Ridge and was pushed southwest by strong northeast winds. The fire slowed when it entered the 2008 Summit Fire boundary. Activity also slowed in unburned forested areas, due to higher humidity and higher fuel moisture. Existing roads provided access and control lines to aid the fire-fighting effort. The Summit Fire and the Loma Fire had very similar effects where portions of each area experienced significant damage and mortality to the understory herbaceous and shrub layers, particularly at higher elevations. The majority of the overstory canopy survived intact. Isolated pockets of mature trees, including hardwoods, redwoods, and Douglas-fir, were damaged or killed. Knobcone pine and chaparral in the upper portion of the watershed burned with high intensity and suffered significant mortality. Both fires burned through a number of swales and watercourses, but higher humidity and fuel moisture levels in these areas kept the intensity of the burn relatively low.

The largest recent fire in the Santa Cruz Mountains was the 2009 Lockheed Fire, which burned 7,819 acres, mainly in the Scotts Creek watershed (CAL FIRE, 2009). While located approximately 12 miles to the west of SDSF, this fire exhibited a similar pattern to the Summit Fire, burning in the chaparral and knobcone pine-covered portions of the burn area near ridges with considerably higher intensity than in the redwood stands located lower on slope near watercourses. Loganbill (2013) reported that the first winter's rainfall after the fire did not produce increased stormflow sediment and event sediment loads, likely due to the fact that near-stream sediment contribution was minimal, and the majority of hillslope-derived sources were not hydrologically connected.

Figure 20. Recent Fire History.




PRE-SUPPRESSION

Pre-suppression is defined as fire protection activities performed before fire occurrence to ensure reduced fire intensity and effective fire suppression. Pre-suppression plans discuss site-specific ways to minimize loss and to reduce hazard and risk. The current pre-suppression plan for SDSF will be updated by the local CAL FIRE Battalion Chief with assistance from the Forest Manager. The more comprehensive plan will include the definition and assessment of high risk and hazard areas within the watershed boundaries, maps of fire defense improvements, prevention techniques, and an evaluation of available resources. State Forest staff continue to work on these activities along with Unit fire protection personnel.

Fire Defense Improvements

Fire defense improvements will be strategically located to protect forest land and neighboring properties. Improvements in the State Forest include three water tanks, shaded fuel breaks, and two helipad locations (Figure 21). In addition, appropriate signing, fire hazard reduction, and adequate access to roads and trails will be added or maintained.

Figure 21. Fire Defense Improvements.



Fire hazard and prevention information, as well as Forest regulations, will be posted on all information boards. The parking area, information boards, and picnic areas will be treated to reduce fire hazards for safety and demonstration purposes. The major roads and trails in the Forest are maintained to provide access for fire protection purposes.

Shaded fuelbreaks protect high value areas such as forest land, historical sites, and neighboring property. Typically, they are areas 100 to 300 feet wide where vegetation and other forest fuels have been decreased in order to reduce the rate of spread of an advancing fire. Less wide shaded fuel breaks are also beneficial. Within SDSF, all main roads and prominent ridgelines will be treated as shaded fuelbreaks. Within these shaded fuelbreaks, dead trees and ladder fuels (shrubs and lower tree limbs) are removed and the overstory canopy is thinned to a level where shade will still retard the growth of new ground fuels. The understory is modified so that a low-growing ground cover is retained within the fuelbreak to provide fuels to start a backfire. Whenever possible, fuelbreaks should visually merge with the surrounding landscape, conforming to the natural features of the area. Periodic maintenance is needed to maintain fuelbreak specifications.

Shaded fuelbreaks in SDSF are being constructed, with the help of crews from Ben Lomond Conservation Camp, along ridges and high use roads and trails. This is to provide safe locations for fire control lines and backfiring; ridgelines are commonly used as control points. Shaded fuelbreaks and roads also aid in the compartmentalization of the property in order to contain a wildfire to its smallest size possible given existing infrastructure. This will minimize the need to install additional fuelbreaks with bulldozers in the event of a wildfire.

Regulations

Restrictions are in effect for hunting and shooting, smoking, and fires within the Forest. Fires are not permitted anywhere in the Forest, including parking areas. A possible exception to these regulations would be campfires in the permit-only group campground (to be considered on a case-by-case basis). Hunting and shooting are administratively prohibited in the Forest, although hunting may be allowed in the future consistent with a program specifically designed by CDFW for SDSF (See Recreation Chapter).

The periods of extreme fire danger for SDSF usually occur from July through October though these periods may be extended by severe weather. During these periods, SDSF will follow the Unit's Red Flag Alert Plan. This is consistent with the plan The Forest of Nisene Marks State Park follows during extreme fire danger conditions. The Forest Manager will coordinate with the Unit Operations Division Chief to determine necessary actions to be employed. The steps include increasing patrols of the Forest, posting red flag alert signs, providing more fire prevention information and awareness of current conditions to Forest visitors, and reducing the number of visitors in the Forest by posting the area as closed.

Education

SDSF staff will coordinate with the Unit Fire Prevention staff for educational purposes. Educational information will be used to reduce the number of human-caused fires within the State Forest. Target groups will include neighbors, visitors, school groups, and local organizations. In general, neighboring property owners pose a risk of human-caused

wildland fires to the Forest. They will be encouraged to meet with Burrell and Soquel Forest Fire Station personnel for information on CAL FIRE's Fire Prevention Program in order to minimize the risk of wildfire.

Signs will be posted on all information boards alerting Forest visitors to the dangers of fire and ways they can minimize these dangers. Dangerous Fire Area signs will be posted when appropriate. During regular patrols, visitors will be informed of fire hazards and prevention methods.

School groups and local organizations will be referred to Fire Prevention staff or local fire stations for fire safety information and presentations. Groups visiting the Forest for education programs will be informed of fire safety, hazards, and prevention techniques. Education programs will also cover the ecological role of fire in the environment and the importance of fire in maintaining biodiversity.

Enforcement

Forest patrol is an important part of fire protection and prevention. SDSF staff will coordinate with the Unit Fire Prevention staff for patrol purposes. Patrols will include public contact, fire detection, and patrol of roads and trails during the fire season. CAL FIRE personnel will be utilized for weekend patrols and major holidays, especially during periods of high fire danger. CAL FIRE peace officers will either provide direct supervision or lead these activities. Additional patrols may be conducted by volunteers as deemed appropriate and safe by CAL FIRE. SDSF staff foresters will also enforce fire-related California Forest Practice Rules at active logging sites on the Forest.

Fire suppression cost recovery will be pursued for damages resulting from deliberate and negligent acts of Forest users. Active investigations will be used to locate responsible parties and recover maximum legal damages.

SUPPRESSION

Suppression tactics are based on information from and implementation of the pre-suppression plan. SDSF staff can support initial attack fire control personnel by providing local expertise regarding current road conditions, fuelbreaks to be utilized for compartmentalization of the fire, vegetation conditions, and cultural resources. The staff may also evacuate visitors, close the fire area, perform law enforcement tasks, provide access through gates, provide information on the location of water tanks, helipads and water drafting sites and assist with media information as appropriate.

Detection

Detection strategies include daily patrols, searching for evidence of fires, and CAL FIRE air flights during extreme fire danger periods or after lightning storms. Also, the Unit's Emergency Command Center will check the Automatic Lightning Detection System (ALDS) for possible strikes in the Forest.

Communication

As part of communication, SDSF will maintain an adequate radio system and stay in close contact with local CAL FIRE forest fire stations (Burrell, Soquel, and Corralitos). Local CAL FIRE fire control personnel will become familiar with the Forest, its road and trail systems, water sources, and landmarks (for use as reference points) and be advised of any changes that occur.

CAL FIRE's resource tracking system, Computer Aided Dispatch (CAD), will be used to dispatch the appropriate personnel and equipment to any fires on SDSF. The State Forest is a defined response area within CAD. The staff will inform the Emergency Command Center of any changes or updates to the CAD database, including information on roads, access points, and fire defense improvements.

POST-SUPPRESSION

Post-suppression activities include the evaluation of pre-suppression information, suppression actions, and fire line suppression repair. Fire suppression repair involves actions needed to repair damage caused by suppression activities. Common suppression repair activities include fireline erosion control, watercourse crossing repair, slash treatment, road repair and drainage improvements, gate and fence repair, waterline repair, mapping of hazard trees, etc. Rehabilitation involves erosion control and other restoration activities not directly caused by fire suppression activities. Unit personnel will evaluate post-suppression activities on an individual fire basis.

To minimize increases in wildfire risks resulting from increased public use of the Forest, the staff will record and compile descriptions and locations of all wildfires occurring at SDSF. This information will be evaluated annually. If an increase in wildfire frequency occurs, appropriate measures will be implemented as needed to reduce wildfire risk.

PRESCRIBED FIRE

Prescribed fire is the controlled use of fire under specific weather and fuel moisture conditions within a predetermined area. Fire, under these conditions, produces the intensity of heat and rate of spread required to accomplish specific management objectives. These objectives could include fire hazard reduction, silvicultural research, and ecosystem enhancement.

A prescribed fire program that involves these objectives will be evaluated for SDSF. A risk and benefit assessment will be the first step in the evaluation. The greatest risk is the loss of fire control and resulting damage that may occur. Benefits include a reduction in fuel load, removal of exotic plant species, and improvements to vegetation and wildlife habitats. A prescribed fire program must consider the mountain community's concerns and address them effectively. An established prescribed fire program provides research opportunities for both resource management and fire behavior purposes.

If a prescribed fire program is established in SDSF, fuel loads and types will be sampled and

mapped, and burning prescriptions will be developed for the different vegetation types found in the Forest. A weather station was established at the lower helipad in 2013 to foster this program and other management and research activities.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

On April 29, 1993, SDSF held a workshop designed to gather information, comments, and concerns from the public. Based on the comments received, there are four major issues concerning fire protection and SDSF.

The first issue involves concern about logging equipment and its potential to cause fires. As stated earlier in this chapter, modern logging methods and equipment are much safer and less likely to generate fires than historic logging practices. Additionally, logging contractors are required by law to develop a fire protection plan for each THP and to follow the state's Forest Practice Rules regarding fire protection. During harvest activities in SDSF, periodic inspections will be performed by both CAL FIRE Forest Practice Officers and Forest staff.

The risk of fire as a result of recreational use is also a concern. Individuals commented on issues such as limiting the number of people allowed in the Forest during periods of high fire danger and the prohibition of fires and camping at all times. Forest policy is to inform users of fire danger and to discourage activities that may result in increased fire risk. Similarly, camping, hunting and shooting, and publicly owned motor vehicles are prohibited in SDSF though camping may occur by special permit and hunting may be allowed in the future (see the Recreation Chapter).

The third issue raised at the workshop concerns the use of prescribed fire in SDSF. Comments about the consequences of both using and refraining from prescribed burning were received. As mentioned above, a risk and benefit analysis will be completed and analyzed before any decisions are made in this area. Lastly, comments regarding fire defense improvements vary from concern about how improvements may change the character of SDSF to suggestions for road accessibility. Currently, roads in the Forest are kept clear of combustible vegetation by Ben Lomond Conservation Camp crews and Forest staff. Likewise, the condition of roadbeds are examined periodically and maintained as needed. Though fire defense improvements may change the appearance of some areas, their presence is needed in order to allow timely and effective response should a fire occur.

PLANNED ACTIONS

1. Coordinate with the Unit Battalion Chief and other fire control personnel to update the pre-suppression plan for SDSF.
2. Continue to identify fire defense improvements and continue their construction. Include the Unit Fire Protection staff in these decisions.
3. Provide patrols to enforce fire prevention policies, coordinating with the Fire

Prevention staff as needed. Forest staff and volunteers will patrol SDSF on weekends and holidays, especially during periods of high fire danger. The Forest will be closed to public use when fire risks become excessive.

4. Conduct fire prevention education for neighbors and Forest visitors utilizing SDSF and fire prevention staff members.
5. Work with the Department of Parks and Recreation personnel from The Forest of Nisene Marks State Park to ensure effective fire protection along Santa Rosalia Ridge.
6. Record and compile descriptions/locations of all wildfires occurring at SDSF including ambient weather and fire hazard conditions, ignition source, area and vegetation types burned, and estimated damage. Implement appropriate measures as needed to reduce wildfire risks.
7. Evaluate the needs and feasibility of a prescribed burn program.

FOREST PEST MANAGEMENT

Forest pests, such as insects, diseases, and vertebrates, have long been established in California's native timberlands. Populations of pests are dynamic and fluctuate in response to climatic and environmental changes such as drought, forest overstocking and windthrow, fire, and other site disturbances. The actual or potential effects of pests may reduce or threaten to reduce anticipated tree growth, species composition, or forest stocking. At the same time, other forest resources, such as wildlife habitat, may be impacted. Integrated forest pest management provides a means to address these issues.

The intent of integrated pest management (IPM) is to prevent or restrain forest pest problems using population suppression and the minimization of factors that predispose trees to infestation. IPM makes use of the benefits of cultural, mechanical, chemical, semiochemical (i.e., synthetic pheromone), and biological pest management alternatives.

Pests known to have caused tree mortality within or adjacent to SDSF are listed in Table 10. There may be other pests of local tree species that are seldom detected or reported, or are of minor significance. State Forest staff will continue to monitor the Forest for early signs of forest pests or conditions that may lead to infestation.

Monitoring is an essential part of detecting early signs of forest pests or scenarios that may lead to infestation. Part of pest monitoring in SDSF will involve forest health surveys to detect pest damage or infestations in standing trees, slash, windthrow, down woody debris, or stumps. SDSF will also assist the pest monitoring program of the California Department of Food and Agriculture by deploying and inspecting gypsy moth traps in high use areas of the Forest. Other efforts to reduce pest damage or predisposition will include:

- * The minimization of injuries to residual trees during forest management activities.

- * Reuse of old tractor roads or recreational trails where available to reduce soil compaction.
- * Retention of a diverse species composition in or adjacent to stands following forest management activities and within or near future regeneration units.

Table 10. Forest Pests of the Soquel Creek Watershed.

PEST	HOST
Fungal Diseases	
Black stain root disease	Douglas-fir
<i>Armillaria</i> root disease	Douglas-fir, oaks, tanoak
Madrone canker	Madrone
Sudden Oak Death	Primary hosts on SDSF are California bay-laurel and tanoak
Insects	
Flatheaded fir borer	Douglas-fir
Douglas-fir beetle	Douglas-fir
Redwood bark beetle	Coast redwood
Western oak bark beetle	Oaks, tanoak
California oakmoth	Oaks, tanoak
Mammals	
Tree squirrels	Coast redwood

- * Avoidance of non-native tree species which may be predisposed to pests with few local pest predators and parasites.
- * Use of CAL FIRE or other forest pest management specialists to train SDSF staff in forest pest recognition and management.

SUDDEN OAK DEATH (SOD):

Sudden oak death is a disease effecting several species of plants. It is caused by a water mold, *Phytophthora ramorum*, and is related to other *Phytophthora* species that cause potato blight and Port Orford cedar root disease. Sudden Oak Death kills some species and

individual plants rapidly. Other species may die back slowly, develop spots on the leaves and stems, or show almost no effects. Susceptible plants include members of the “red” oak group - coast and canyon live, black and Shreve oaks; as well as tanbark oak, California bay-laurel, rhododendron, and many other native and non-native species.

The disease was first found around Mt. Tamalpais in Marin county in 1995. It may have been imported to California from Europe on nursery shipments of rhododendrons. Rapid expansion of the number of dead and diseased trees and the geographical area affected occurred after wet winters and springs in the late 1990s and again from 2004 through 2006. The disease is now confirmed to occur in California throughout the San Francisco Bay Area, and in coastal counties from Monterey to Humboldt. For more information on SOD, please see <http://www.suddenoakdeath.org/>.

The disease is well-established at SDSF and in surrounding areas, and effects tanoaks, coast live oaks, and other susceptible species throughout the Forest. The death of trees in SOD disease centers has created openings in areas formerly dominated by tanoaks. Research conducted at SDSF and elsewhere indicates that over time tanoaks will no longer be a dominant species in the overstory in these locations. Douglas-fir and bay-laurel trees are likely to become the dominant overstory species in these areas in the future.

SDSF is within the declared SOD Zone of Infestation (ZOI) established by the California State Board of Forestry and Fire Protection and is within the “Regulated Area” for SOD as designated by the California Department of Food and Agriculture (CDFA). The ZOI and Regulated Area are identical and cover all portions of the 14 infested counties identified in the California Department of Food and Agriculture Section 3700: Oak Mortality Disease Control.

Federal regulations from the USDA-APHIS, and state regulations from CDFA address SOD concerns. CDFA regulations that limit the movement of host materials apply to forest management activities on SDSF including timber harvest, timber stand improvement activities, and harvest of minor forest products. Under the state regulation, host material cannot be transported from the Regulated Area unless accompanied by a compliance agreement. Wherever a ZOI applies, the Forest Practice Rules [14 CCR 917.9(a)] require that mitigations be included in Timber Harvesting Plans (THPs) to prevent the spread of the infestation. A federal quarantine for *P. ramorum* was issued as an interim rule by USDA – APHIS, with the most recent rule dated February, 2007. For more information on the Federal rule see <http://nature.berkeley.edu/comtf/pdf/APHIS-2005-0102-0001.pdf>.

SOD is known to occur at SDSF, and host material that is likely to be transported may consist of logs from host species produced as part of a harvest plan and minor forest products such as salvage sawlogs, firewood, and greenery.

SOD can spread via host material. Therefore, plants, plant parts, unprocessed wood and wood products, and other products of the above mentioned hosts cannot be moved from counties infested with SOD without authorization by the County Agricultural Commissioner or CAL FIRE's and USDA Forest Service's harvest document approval process. The term

"harvest document" refers to any document filed with the California Department of Forestry and Fire Protection that authorizes the removal of forest products for commercial purposes. See <http://pi.CDFCA.ca.gov/pqm/manual/htm/455.htm> for California Department of Food and Agriculture's regulations regarding commodities covered and restrictions of their movement.

Firewood sales

On State Forest lands, where the public is required to have a permit to collect firewood, CAL FIRE uses this public contact to educate the person(s) about SOD by providing a Pest Alert or other information on SOD along with the firewood permit. A provision is added to the permit explaining the current regulations and that compliance is required.

Hazard Reduction for SOD

Severely diseased or dying oaks and tanoaks with stem infections are often attacked by bark beetles (western oak bark beetle and ambrosia beetles) and infected with decay fungi such as *Hypoxylon* spp. Trees with these secondary insect attacks and decay fungi may be structurally weakened and will be removed if adjacent to roads, trails, or other high-use areas.

In addition, trees killed by SOD may lead to increased surface and crown fuels, thus potentially influencing fire behavior. In addition to addressing public safety issues, removal of dead and dying trees should also lessen fire intensity and reduce the risk of crown fires near disease centers.

PITCH CANKER

SDSF staff will incorporate the most current best management practices as identified by the California Pitch Canker Task Force for controlling the distribution and spread of Pitch Canker. Very few pine trees exist at SDSF.

TREE SQUIRRELS

Tree squirrels may damage sapling redwoods by stripping the bark from upper stems. This causes partial or complete girdling, leading to either top kill or sapwood decay. As damage from squirrels is associated with dense second growth redwood stands, thinning is the only recommended management action. Where trees are more widely spaced, squirrel damage will generally be reduced.

PLANNED ACTIONS

1. Continually monitor the Forest for signs of pests and notify the CAL FIRE Forest Pest Management Specialist of any findings. Take action as needed to minimize or eliminate any problems.
2. Thin dense stands of trees where necessary to maintain healthy growing conditions.

3. Provide applied forest pest research opportunities to interested agencies, institutions, or organizations.
4. Coordinate with Santa Cruz County in the detection of and protection against gypsy moths or other introduced pests.
5. Continue to remove dying and dead trees adjacent to high-use roads, trails and other facilities.

CHAPTER 13: ARCHAEOLOGY

HISTORY OF SDSF AREA

Until the early 1990s the archaeology of the Central Coast was not well understood due to the scarcity of studies. Many of the most serious gaps in our understanding of the cultural and environmental diversity of the Central Coast have now been filled. SDSF is located within the ethnographic territory of Native people who spoke a dialect of the Costanoan (now also known as Ohlone) family of languages. Although native people speaking various Costanoan languages occupied the South Coast Ranges between San Pablo Bay and Monterey prior to the arrival of Europeans, “Costanoans” [from the Spanish *costanos*, “coast people”] were neither a single ethnic group nor a political entity (Levy, 1978). Rather, Costanoan is a linguistic term designating a family of eight languages. One of these, Awaswas, was the language spoken by people living along the coast between Davenport and Aptos.

Although the Spanish sailed along the Central Coast as early as 1542, the Portola overland expedition in 1769 initiated long term contact. At that time most of this portion of the coast was occupied by a large number of small, autonomous tribelets (Kroeber, 1925). The native population was decimated during the Spanish, and later, American invasions of their territory. Early explorers, priests and settlers, and later ethnographers documented at least some aspects of Ohlone worldview and material culture. The documentation from these early contacts is very incomplete. Systematic anthropological description of the native people and their culture was not initiated until early in the twentieth century following at least 150 years of drastic population decline (Jones et al., 2007).

Utilization of the area now SDSF since 1542 is better understood, with a body of historical records and documents to supplement the archaeological record. In Santa Cruz County four chronological subdivisions have been defined for the Historic Period: Protohistoric (1542-1769), Spanish Colonial (1769-1822), Mexican (1822-1848) and Anglo-American (1848-present). Historic-era sites on SDSF span from the Mexican Period (1822-1848) to the Anglo-American Period (1848-present). Please see the “History of Ownership” section of Chapter 3 and “SDSF Facilities” section of Chapter 14 for a summary of the history of SDSF and neighboring properties. For a detailed account of the archaeology and history of SDSF see Dillon (1992).

ARCHAEOLOGICAL RESEARCH

An archaeological and historical field survey of SDSF was conducted during the summer of 1991. The chief investigator was Dr. Brian Dillon, a consulting archaeologist associated with the California State University at Northridge. In March of 1992, an archaeological and historical report was generated from the survey. In addition to study results, this report also includes information from archival records research and extensive oral history interviews. During the survey, Dr. Dillon and his crew discovered six archaeological sites within the State Forest boundaries: two prehistoric and three historic sites, as well as one site with both prehistoric and historic features. Additional studies covering SDSF history and archaeology have identified many additional sites that are documented in Confidential Archaeological Addendums to THPs

and archived with the Northwest Information Center.

The significance of each site was determined by its archaeological and historical value, as outlined in state and federal guidelines. Significance, as defined by these guidelines, is based on uniqueness and preservation, with both considered in the determination of a site's value. Uniqueness refers to how many other similar features exist (on other sites), while preservation refers to the condition of the features remaining on the site. A site is not considered significant if it, although unique, has been completely destroyed, as there is nothing left to protect or study.

A confidentiality policy exists which limits public disclosure of sensitive archaeological and historical resources. Consequently, site locations in SDSF with moderate to high levels of significance will not be revealed to the general public. The confidentiality policy protects the resources from artifact collection, site excavation, and vandalism. The policy was approved by the State Historical Resources Commission under authority of Public Resources Code Section 5020.4 (c).

Since 1991, several additional archaeological surveys have been conducted on SDSF. As a result of these surveys and the initial one conducted by Dr. Dillon, approximately 30 percent of the Forest has been examined at least once for archaeological resources. Surveys conducted since 1991 have resulted in the identification and recording of numerous additional prehistoric and historic-era sites.

PREHISTORIC SITES

The prehistoric era is believed to have begun on the central coast about 4,000 years ago. The most active times were during the Late Prehistoric Period, 1000 - 1600 AD, when hundreds of sites were established. The prehistoric sites found in SDSF are from this Late Prehistoric Period. Prehistoric sites found on SDSF include the following constituents: bedrock mortars (some including cupules and petroglyphs); portable size mortars; large lithic tools including manos, metates, a pestle and cooking stones; small lithic tools and debitage consisting of non-native rock types including chert; small stone manuports of unknown purpose; and midden soils.

HISTORIC SITES

The Historic Period began along the central coast in 1769 and continues to the present day. The historic sites found in SDSF span from the Mexican Period, 1822 - 1848, to the Anglo-American Period, 1848 to present. The most recent site in the Forest, however, dates back to World War II.

Historic sites found on SDSF include the following features and artifacts: Depression-era split products manufacturing sites; logging-related equipment and materials such as 55-gallon drums, steel snatch-bocks, wire rope/cable of various diameters and style, crosscut saws, oiler jugs, and related items; logging-related features such as modified stumps, cable roads, and skid roads; sawmill-related equipment and materials and a sawpit; a livestock corral made from redwood pickets; scrapped automobiles and parts; segments of wagon roads constructed between 1857 and 1870; several segments of old fence comprised of mostly split redwood posts and fragments of rusted barbed wire; a variety of components from a steam donkey and the log skids which it was

mounted upon; single-family wood-framed residences and various associated landscape features built between 1936 and 1959; remains of other structures of unknown age and purpose; and discarded cans, bottles and miscellaneous debris.

All of the sites and isolates described above were documented with records prepared in accordance with California Office of Historic Preservation (OHP) guidelines (CDPR 1995).

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Following completion of the archaeological study in SDSF, members of the public expressed a desire to learn about Dr. Dillon's discoveries and their significance. Individuals were interested in research or other studies that might result from the findings as well as seeing the archaeological sites. Eventually, all sites of moderate and high significance will be more thoroughly evaluated and, depending on the results, may be available for public viewing.

MANAGEMENT GUIDELINES

1. Heritage resources on SDSF will be protected, especially during planning and management activities including timber harvesting, recreation, and forestry education.
2. Sites determined eligible and/or listed on the California Register of Historical Resources will be preserved and kept confidential, as per applicable state statute and regulation . If, after thorough and careful study, it is determined that certain sites can endure limited public use, they may be made accessible to the community. Such determinations shall be made in consultation with the California State Historic Preservation Officer (SHPO) and the CAL FIRE Northern Region-Coast Area Archaeologist (or CAL FIRE State Forests Archaeologist).
3. Educational opportunities, including the display of resilient historic features, will be incorporated into SDSF's Demonstration and Forestry Education programs.

PLANNED ACTIONS

1. Provide the opportunity for scientific study and research on all archaeological and historical resources. Researchers working with significant sites will be expected to uphold all confidentiality policies and will conduct work only after a specific research design has been reviewed and approved by CAL FIRE Archaeology staff.
2. Identify and catalog additional historic information including maps, photographs, written documents, interviews, existing archaeological collections and archival materials.
3. Monitor and periodically inspect heritage resources on SDSF to ensure that existing policies are affording effective protection to those resources (cf. Executive Order W-26-92; PRC Sections 5020 through 5024).

CHAPTER 14: ROADS AND OTHER IMPROVEMENTS

SDSF FACILITIES

The first development of roads within SDSF can be traced back to the mid-nineteenth century. According to local historian Ron Powell (Rancho de Palo de Yesca: the Shoquel Augmentation Rancho, n.d.), the Santa Cruz Gap Turnpike (an arterial between Santa Cruz and the Santa Clara valley) was completed in May of 1856. This road included parts of what are now Hihn's Mill Road, Longridge Road, and Spanish Ranch Road. Further development of roads occurred when Frederick A. Hihn acquired the SDSF property in 1863.

In the 1870's, Hihn built Sulphur Springs Road to reach his Sulphur Springs Resort as well as a private road for logging and mill access. These roads, along with segments of the Santa Cruz Gap Turnpike, created what is now known as Hihn's Mill Road. This route extends from the Olive Springs Quarry to the parking area off Highland Way. Additionally, logging by Jared and Seth Comstock in 1878 led to the construction of Comstock Mill Road and portions of Robinwood Lane.

The remainder of the roads in SDSF were constructed by the three timber companies that owned the property after Hihn. This later road construction started with the Monterey Bay Redwood Company in 1924, continued through the CHY Company ownership in the 1960s and 1970s, and ended with the Pelican Timber Company in the 1980s. Figure 19 (see Recreation Chapter) shows the locations of all existing Forest and access roads.

The only portion of SDSF that does not contain roads is the area south of Badger Spring to Santa Rosalia Ridge. This area is bordered by Sawpit Trail (see the Recreation Chapter for information on trails) on the east, the East Branch of Soquel Creek on the northwest, and the Forest boundary on the south. Steam donkeys were used to log this section prior to the introduction of crawler tractors, so no major roads or trails were built. Scars on the slopes above Badger Spring are still evident from this type of logging.

The only bridges associated with SDSF cross the East Branch at the Highland Way entrance and a short distance downstream from the confluence with Amaya Creek on Hihn's Mill Road. The Highland Way bridge is a narrow nine-foot wide rail flatcar bridge supported by earthen abutments. Future plans call for replacing this bridge with a wider structure. This bridge and the nearby parking area are located on the Burch property. The second bridge, Hihn's Bridge, is a 90-foot-long rail flatcar bridge installed by CAL FIRE in 1999. Hihn's Bridge was built as part of the contract with the timber purchaser for the Amaya THP #1-98-027 SCR. Plans for the Hihn's Bridge were part of the THP and additional reports and plans are stored at the SDSF office. The load rating for this bridge as determined by the California Department of Transportation is MS-18 or HS-20. The point load capacity is 200,000 to 240,000 pounds. At the time the bridge was built it cost about \$120,000. Future plans include a permanent, year-round crossing of the East Branch of Soquel Creek at Longridge Road.

LEGAL ACCESS

When the State acquired the major portion of the Pelican Timber Company's holdings along the East Branch in 1988 (refer to the Administration Chapter for details), the property had limited accessibility from county roads. The Forest is surrounded by private property on three sides and The Forest of Nisene Marks State Park (largely undeveloped) to the south. To ensure access, two deeded right-of-ways through private property were granted at the time of acquisition, providing for both administrative and public access.

The first right-of-way is through the Burch property to the east of SDSF. This route includes 0.7 miles of Hihn's Mill Road out to Highland Way, and provides the only public vehicular access to SDSF (public vehicles are not allowed in the Forest but can drive to and park in a designated area just off of Highland Way). The second right-of-way is through the CHY Company property to the southwest of SDSF and includes one mile of Hihn's Mill Road from the Forest boundary to the Olive Springs Quarry. Olive Springs Road, which connects with the major arterial of Soquel-San Jose Road, can then be accessed through the quarry for administrative purposes only.

DRIVABLE ROADS

All drivable roads in SDSF have been evaluated for safety and stability and are open to varying degrees of seasonal administrative vehicular use. A number of old logging roads and constructed skid trails in the Forest need to be accurately mapped and to have their suitability for reuse determined. Table 11 is a summary of the drivable roads in SDSF, with information on the road name, length, and location.

PUBLIC ACCESS ROADS

County roads leading to SDSF access points include Olive Springs Road, Highland Way, and Eureka Canyon Road. Olive Springs Road provides reliable year-round administrative access through the Olive Springs Quarry. As mentioned above, this road connects with Hihn's Mill Road at the southwestern edge of the Forest. This route from the quarry has a rock surface and is generally open.

Highland Way, which is east of Soquel-San Jose Road, leads to the eastern portion of SDSF and, as previously mentioned, provides the only public vehicular access. Highland Way has been unreliable during winters, however, due to landslide closures. Eureka Canyon Road, which winds through the mountains from Corralitos, provides a longer alternate route to the Highland Way entrance. Eureka Canyon Road is typically used when closures block Highland Way but it is also subject to slides. The Aptos Creek Fire Road, located within The Forest of Nisene Marks State Park, provides administrative access as well as a public entrance (by foot or bicycle) at the east end of Ridge Trail.

ROAD BUILDING AND MAINTENANCE

Road building and maintenance are critical elements of forest management. Forest roads are usually the largest source of management-related sediment and they provide the means to recreate, harvest timber, access emergency situations, maintain facilities, and patrol the Forest. Therefore, they should be in the best possible condition at all times. Culverts, bridges, ford crossings, water bars, rolling dips, and drainage ditches (structures which divert water away from or off of roads) need to be regularly inspected and repaired as necessary.

Table 11. Drivable Road Segments of SDSF.

ROAD NAME	LENGTH	LOCATION
Hihn's Mill Road	7.0 miles	Highland Way to Olive Springs Road
Amaya Creek Road	2.0 miles	Hihn's Mill Road to Comstock Mill Road
Comstock Mill Road	0.7 miles	Robinwood Lane to private property
Sulphur Springs Road	1.5 miles	Hihn's Mill Road to Ridge Trail
Corral Trail	1.2 miles	Sulphur Springs Road to the end of roaded portion of Corral Trail
Tractor Trail	1.3 miles	Hihn's Mill Road to end of roaded portion of Tractor Trail
Longridge Road	1.5 miles	Hihn's Mill Road to private property
Amaya Basin Road	0.7 miles	
Amaya Pond Road	0.5 miles	
Amaya Spillway Road	0.1 miles	
Lower Helipad Road	0.1 miles	
Shortridge Road	0.1 miles	

Winter inspection is crucial as the majority of water and soil movement occurs during the rainy season. New roads or skid trails will not be constructed or old roads or skid trails rehabilitated without thorough evaluations of topography, intended use, soil stability, drainage capabilities, and construction costs.

There are approximately 19 miles of existing or abandoned roads within SDSF. According to the analysis done for the EIR, about 22 miles of new roads are needed to complete the Forest road network and approximately 15 miles of road will eventually be abandoned. The existing road density is about 4.6 miles per square mile.

California Forest Practice Rules specific to road building and maintenance will be applied during all forest management activities. This will be particularly emphasized during timber harvesting operations. These regulations, along with other site-specific strategies, will help minimize erosion and sediment delivery to watercourses, visual disturbance, and road construction/maintenance needs. Generally, all construction or rehabilitation of roads will include outslipping of road surfaces, rolling dips, and limited use of inside ditches to minimize hydrologic connectivity to stream channels. A road management plan will be developed which includes information from the 2003 Pacific Watershed Associates road inventory, which identified 82 sites with significant sediment delivery potential to watercourse channels.

THE FOLLOWING PUBLIC COMMENTS WERE RECEIVED DURING REVIEW OF THE 1998 GENERAL FOREST MANAGEMENT PLAN

Since the roads of SDSF will not be available for motorized use by the public, comments regarding them have been minimal. The issue of Forest access, however, is another matter. While most of the access concerns described in the Administration Chapter relate to the lack of public entrances, the inadequacy of existing public entrances is a problem as well. Individuals have expressed that the entrances are not only remote, but also difficult to maneuver in some cases. This is particularly true of equestrians who trailer their horses to the Highland Way parking area. As detailed in the Administration Chapter and Management Guideline Number 1 below, SDSF is actively seeking new legal access points. New entrance areas will be designed so that multiple uses can be accommodated.

Another concern is that excessive road building will take place in the Forest. Though SDSF does not have a detailed road construction plan, California Forest Practice Rules place restrictions on the amount and types of roads that can be built in forested areas. It is the intent of the Forest to adhere to these regulations in addition to the examination of all projects and possible impacts on a site-specific basis.

MANAGEMENT GUIDELINES

1. Work with neighboring ownerships to maintain and improve access to the Forest. Continue working with the Olive Springs Quarry and the County of Santa Cruz to ensure that both ends of Hihn's Mill Road remain open.
2. All roads and other improvements shall be monitored and maintained in good condition. This will provide for safety and help prevent surface and mass erosion. Ben Lomond Conservation Camp and California Conservation Corps crews, CAL FIRE heavy fire equipment operators, and volunteers will help with road maintenance and repair, as appropriate.
3. New roads and other improvements will be developed as needed for access, resource protection, and forest management activities. High standards and compliance with California Forest Practice Rules for new logging road construction and existing road rehabilitation to minimize soil damage will be emphasized. Roads located within or near

unstable areas will be designed with the assistance of CEG staff from the California Geological Survey. New construction and maintenance methods will be researched and incorporated to demonstrate their effectiveness. Methods to manage portions of the Forest without the use or construction of new roads will be investigated and may provide additional opportunities for research and demonstration.

4. Roads that are not needed to provide access, or are at high risk for sediment production, or are difficult to maintain and monitor will be abandoned.
5. Design watercourse crossings to reduce sediment inputs from vehicle use, bicyclists, and equestrians on roads and install foot bridges at stream crossings on trails.
6. Close the Highland Way parking area and Forest roads during periods of heavy precipitation and soil saturation when sediment delivery to a watercourse is a risk.

PLANNED ACTIONS

1. Determine the best approach for a road management plan and develop a plan for SDSF. This plan will include the inventory, assessment, and risk-rating of Forest roads that was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (PWA, 2003).
2. Road conditions will be continually assessed to determine which segments are no longer suitable for use.
3. Investigate and, if possible, acquire additional administrative and public access to SDSF. Alternative funding sources will be examined for these acquisitions as well as for construction and maintenance. (Refer to the Administration Chapter for more information on access issues.)
4. Continue to upgrade Hihn's Mill Road and the Highland Way entrance parking area for all-season use. This includes road rocking and replacing culverts at risk of failure with larger culverts. Coordinate work at the Highland Way parking area and along the Burch property right-of-way with the landowner representatives. Upgrade Longridge, Amaya Creek, and Sulphur Springs Roads for improved emergency access.
5. Widen and provide permanent abutments to the Highland Way entrance bridge in cooperation with Burch/Redwood Empire. Maintain trash racks above the inflow to culverts in the Forest. An inventory, assessment and risk-rating of culverts was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (PWA, 2003). Continue to upgrade and remediate watercourse crossings identified in the PWA report.
6. Continue to implement the mile-post identification system for roads and trails to facilitate management and emergency response. Maintain the gate identification system. Continue

to build upon the GIS layers and maps of roads and gates to be utilized in computer-aided dispatching for emergencies.

7. Obtain review of the location of all roads, landings, and skid trails on unstable areas by a Certified Engineering Geologist.
8. Implement the Streambank Failure/Hihn's Mill Road repair project along the portion of Hihn's Mill Road east of Hihn's Bridge using a fish friendly design.
9. Install a permanent, year-round crossing of Soquel Creek at Longridge Road.

REFERENCES

LITERATURE

- Alley, D.W. 2006. Comparison of Juvenile Steelhead Densities, Population Indices and Habitat Conditions in Soquel Creek, Santa Cruz County, California, 1997-2005; with an Index of Expected Adult Returns. Prepared for the Soquel Creek Water District.
- Alley, D.W. and Associates. 1992. Recommended Monitoring and Logging Guidelines for the Soquel Demonstration State Forest Related to Aquatic Resources and Flood Control. Unpublished report submitted to Thomas Sutfin, SDSF Forest Manager. 24 p.
- . 2001 and 2002. Soquel Lagoon Monitoring Reports. Prepared for the City of Capitola.
- . 2003. Soquel Creek Salmonid Assessment and Enhancement Plan. Santa Cruz County Resource Conservation District, Soquel Creek Watershed Assessment and Enhancement Project Plan. Capitola, California. 182 p.
- . 2004. 2004 Soquel Creek Lagoon Management and Enhancement Plan Update. Prepared for the City of Capitola. Brookdale, California. 172 p.
http://www.mpwmd.dst.ca.us/mbay_irwm/irwm_library/soq_cr_mgt_plan.pdf
- . 2011. Juvenile Steelhead Densities in the San Lorenzo, Soquel, Aptos and Corralitos Watersheds, Santa Cruz County, California; with San Lorenzo and Soquel Trend Analysis. Santa Cruz, California. 93 p.
- Anderson, K.R. 1995. A Status Review of the Coho Salmon (*Oncorhynchus kisutch*) in California South of San Francisco Bay. California Department of Fish and Game report to the Fish and Game Commission. 82 p.
- Arora, D. 1986. Mushrooms Demystified. Ten Speed Press. Berkeley, California. 959 p.
- Barnhart, R.A. 1986. Species Profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest)--steelhead. U.S. Fish and Wildlife Service Biological Report 82 (11.60). U.S. Army Corps of Engineers, TR EL-82-4. 31 p.
http://www.nwrc.usgs.gov/wdb/pub/species_profiles/82_11-060.pdf
- Bartlett, R.D. and P.P. Bartlett. 2009. Guide and Reference to the Amphibians of Western North America (North of Mexico) and Hawaii. University Press of Florida.
- . 2009. Guide and Reference to the Snakes of Western North America (North of Mexico) and Hawaii. University Press of Florida.
- . 2009. Guide and Reference to the Turtles and Lizards of Western North America (North of Mexico) and Hawaii. University Press of Florida.
- Becker, G.S. and I.J. Reining. 2008. Steelhead/rainbow trout (*Oncorhynchus mykiss*) resources

south of the Golden Gate, California. Cartography by D.A. Asbury. Center for Ecosystem Management and Restoration. Oakland, California. Prepared for California State Coastal Conservancy and The Resources Legacy Fund Foundation. 293 p. plus figures and tables. http://www.cemar.org/SSRP/pdfs/SSRP_textOnly.pdf

Bedrossian, T.L., and Sowma, J.A. 1990. Earthquake damage in Soquel Demonstration State Forest, Santa Cruz County, *in* McNutt, S.R., and Sydnor, R.H., eds., *The Loma Prieta (Santa Cruz Mountains), California, earthquake of 17 October 1989*: California Department of Conservation, Division of Mines and Geology Special Publication 104, pp. 105-112.

Benyus, J.M. 1989. *Northwoods Wildlife: A Watcher's Guide to Habitats*. NorthWord Press, Inc. Minocqua, Wisconsin. 365 p.

Berg, N.H., K.B. Roby, and B.J. McGurk. 1996. Cumulative watershed effects: applicability of available methodologies to the Sierra Nevada. In: *Sierra Nevada Ecosystem Project; final report to Congress*. Davis, California: University of California, Davis. Centers for Water and Wildland Resources: 39–78. Vol. 3. Chapter 2. http://pubs.usgs.gov/dds/dds-43/VOL_III/VIII_TOC.PDF

Berlekamp, B. 1994. 1994 SDSF Fish Habitat Survey Report. California Department of Forestry and Fire Protection, Soquel, California. Appendix A. 39 p. *in* Berlekamp, B. and T. Sutfin. *Draft Fisheries Management Plan*. Soquel Demonstration State Forest. Soquel, California. 18 p. plus Appendices. http://calfire.ca.gov/resource_mgt/downloads/Soquel/DRAFT-Fisheries_MgmtPlan.pdf

Berrill, J-P., and K.L. O'Hara. 2009. Simulating multiaged coast redwood stand development: interactions between regeneration, structure, and productivity. *Western Journal of Applied Forestry* 24(1): 24-32.

Bisson, P.A., J.L. Nielsen, and J. W. Ward. 1988. Summer production of coho salmon stocked in Mount St. Helens streams 3-6 years after the 1980 eruption. *Transactions of the American Fisheries Society* 117: 322-335. http://www.fs.fed.us/pnw/lwm/aem/docs/bisson/1988_bisson_summer_production_of_coho.pdf

Blazej, N. 1997. *Education Study for Soquel Demonstration State Forest*. Prepared for California Department of Forestry and Fires Protection, Soquel Demonstration State Forest Soquel, California. University of California, Berkeley. Berkeley, California. 77 p.

Breschini, G.S. 1972. *The Indians of Monterey County*. Monterey County Archaeological Society. Carmel, California. 48 p.

Broadbent, S.M. 1972. *The Rumsen of Monterey: an ethnography from historical sources*. *Contributions of the University of California Archaeological Research Facility* 14:45-94. <http://digitalassets.lib.berkeley.edu/anthpubs/ucb/text/arf014-005.pdf>

Butler, B., R. Bates and R. Hester. 2001. *Forestry Education Center Study for Soquel*

Demonstration State Forest An Analysis of California Environmental Education Centers. University of California, Berkeley. Berkeley, California. 76 p.

---. 2001. Forestry Education Center Study for Soquel Demonstration State Forest Listening Report. University of California, Berkeley. Berkeley, California. 30 p.

Butler, B. and R. Hester. 2001. Forestry Education Center Study for Soquel Demonstration State Forest Proposed Program. University of California, Berkeley. Berkeley, California. 35 p.

Cafferata, P. and C. Poole. 1993. Watershed Assessment for the East Branch of Soquel Creek. Final report submitted to the California Department of Forestry and Fire Protection. Sacramento, California. 59 p.

California Department of Fish and Game. 1996. Macroinvertebrate Sampling on Soquel Demonstration State Forest and Soquel Forest Species List. 7 p.

---. 2004. Recovery Strategy for California Coho Salmon. Report to the California Fish and Game Commission. Sacramento, California. 594 p.

---. 2008. Complete List of Amphibian, Reptile, Bird and Mammal Species in California. Sacramento, California. 25 p.

http://www.dfg.ca.gov/biogeodata/cwhr/pdfs/species_list.pdf

---. 2009. Special Animals List.

---. 2010. State and federally listed endangered and threatened animals of California. Department of Fish and Game, Biogeographic Data Branch, California Natural Diversity Database. Sacramento, California. 12 p.

California Department of Forestry and Fire Protection (CDF). 1993-1999, 2001-2012. Steelhead Trout Population Survey Report. Soquel Demonstration State Forest. Soquel, California.

---. 1997-1999, 2001-2003, 2005-2012. Instream Temperature Monitoring, East Branch of Soquel Creek, Fern Gulch Creek, and Amaya Creek. Soquel Demonstration State Forest. Soquel, California.

--- (CAL FIRE). 2009. Lockheed Fire Post Fire Risk Assessment. San Mateo-Santa Cruz Unit. 109 p.

---. 2010. Soquel Demonstration State Forest, Option A Plan for Achievement of Maximum Sustained Production of High Quality Timber Products. Soquel, California. 40 p.

---. 2014. California Forest Practice Rules, Title 14, California Code of Regulations, Chapters 4, 4.5 and 10. Sacramento, California. 339 p.

California Department of Parks and Recreation Office of Historic Preservation. 1990. Archaeological Resource Management Reports (ARMR): Recommended Contents and Format. Preservation Planning Bulletin Number 40. Sacramento, California. 29 p.
<http://ohp.parks.ca.gov/pages/1054/files/armr.pdf>

----. 1991. Guidelines For Archaeological Research Designs. Preservation Planning Bulletin Number 5. Sacramento, California. 17 p.
<http://ohp.parks.ca.gov/pages/1069/files/arch%20research%20design.pdf>

----. 1995. Instructions for Recording Historical Resources. Sacramento, California. 57 p.
<http://www.parks.ca.gov/pages/1054/files/manual95.pdf>

CDPR (California Department of Parks and Recreation) State Historic Resources Commission. 1993. Guidelines for the Curation of Archeological Collections. Sacramento, California. 20 p.
<http://ohp.parks.ca.gov/pages/1054/files/guide93.pdf>

CaliforniaHerps.com. A Guide to the Reptiles and Amphibians of California
<http://www.californiaherps.com/>

California State Board of Equalization. Timber yield taxes for 1994-2012. Timber Tax Office. Sacramento, California.

Chartrand, S., B. Hecht, and D. Shaw. (Balance Hydrologics, Inc.) 2003. Soquel Creek Watershed Assessment: Geomorphology and Baseflow Hydrology. In: Santa Cruz County Resource Conservation District. Soquel Creek Watershed Assessment and Enhancement Plan. Capitola, California. 89 p. plus Tables and Figures.
<http://www.balancehydro.com/pdf/soquelwatershed.pdf>

Clark, D.T. 1986. Santa Cruz County Place Names. Santa Cruz County Historical Society. Santa Cruz, California.

Clinton, N. 2009. Soquel Demonstration State Forest WHR Mapping. Berkeley Environmental Technology International, LLC. Berkeley, California. 23 p.

Conway, S. 1976. Logging practices: principles of timber harvesting systems. Miller Reeman Publications, Inc. San Francisco, California.

Dillon, B. 1992. Archaeological and Historical Survey of Soquel Demonstration Forest. CDF Archaeological Reports, Number 6. 177 p.

Elliott, M. 2013. Biological control of California bay laurel resprouts using *Chondrostereum purpureum* and other wood decay fungi. Project description. 2 p.

Elliott, M., G. Chastagner, S. Shamoum, G. Sumampong, E. Goheen, and A. Kanaskie. Biological control of tan oak and bay laurel resprouts using the fungus *Chondrostereum purpureum*. Poster. 1 p.

Erickson, L. 2011. Field Herp Report. 9 p.

---. 2011. Soquel Demonstration State Forest Herpetofauna Survey Report. 2 p.

---. 2011. Soquel Demonstration State Forest Herpetology Survey Result summary 12/2010-12/2011. 3 p.

---. 2014. SDSF Herpetological Survey Report. 4 p.

Federal Register. 1996. Endangered and Threatened Species: Threatened Status for Central California Coast Coho Salmon Evolutionarily Significant unit (ESU). Vol. 61, No. 212. 12 p. <http://www.gpo.gov/fdsys/pkg/FR-1996-10-31/pdf/96-27887.pdf>

---. 2005. Endangered and Threatened Species: Final Listing Determinations for 16 ESUs of West Coast Salmon, and Final 4(d) Protective Regulations for Threatened Salmonid ESUs. Vol. 70, No. 123. 45 p. <http://www.gpo.gov/fdsys/pkg/FR-2005-06-28/pdf/05-12351.pdf>

---. 2012. Endangered and Threatened Species: Range Extension for Endangered Central California Coast Coho Salmon. Vol. 77, No. 63. 12 p. <http://www.gpo.gov/fdsys/pkg/FR-2012-04-02/pdf/2012-7860.pdf>

Foley, J. 2011. Summary, Davis Tick-Borne Disease Activities through Spring 2011.

Foster, D.G. and M.V. Thornton. 2001. Management Plan for CDF's Historic Buildings and Archaeological Sites. Archaeological Report Number 22, California Department of Forestry and Fire Protection. Sacramento, California. http://www.fire.ca.gov/resource_mgt/archaeology/downloads/plan.pdf

Garbelotto, M. 2013. Progress Report SDSF Experiments 12/2013 on Sudden Oak Death. Memo to SDSF staff. 12 p.

Good, T.P., R.S. Waples, and P. Adams (editors). 2005. Updated status of federally listed ESUs of West Coast salmon and steelhead. US Department of Commerce, NOAA Tech memo. NMFS-NWFSC-66. 598 p. <http://swfsc.noaa.gov/publications/fed/00749.pdf>

Gray, A.E. 2014. Importance and spatial distribution of *Phytophthora ramorum* host species in a coast redwood forest. For Master of Science degree (Applied Geography), University of North Texas. 95 p.

Greene, M. 2011. 2011 Raptor Survey for the Fern Gulch THP 1-09-069-SCR. 13 p.

---. 2011. 2011 Raptor Survey for the Rim THP 1-09-107-SCR. 8 p.

---. 2012. 2012 Raptor Survey for the Fern Gulch THP 1-09-069-SCR. 13 p.

---. 2012. 2012 Raptor Survey for the Rim THP 1-09-107-SCR. 8 p.

Greening Associates. 2003. Soquel Creek Riparian Vegetation Assessment. Santa Cruz County Conservation District. Soquel Creek Watershed Assessment and Enhancement Project Plan. Capitola, California. 45 p. plus Appendices.

Greenlee, J.M. 1981. Santa Cruz County Fire History: (Maps--CA: J.M. Greenlee, 1981. 1:25,344). University of California, Santa Cruz. Maps References 6, 1528, 55, 673.

Griffith, K. 2014. Effects of Vegetation Structure and Canopy Exposure on Small-Scale Variation in Atmospheric Deposition Inputs to a Mixed Conifer Forest in California. For Master of Science degree (Geography), University of North Texas. 75 p.

Harrington, J.P. 1942. Culture element distributions: XIX Central California Coast. University of California Press, Berkeley and Los Angeles, California. Anthropological Records 7(1):1-46. <http://digitalassets.lib.berkeley.edu/anthpubs/ucb/text/ucar007-002.pdf>

Hayden, K.J., A. Netterl, R. Dodd, and M. Garbelotto. 2011. Will all the trees fall? Variable resistance to an introduced forest disease in a highly susceptible host. *Forest Ecology and Management* 26 (2011) 1781-1791. 11 p.

Haydon, W. 2001. Geologic and Geomorphic Features Related to Landsliding, Amaya and Fern Gulches, Soquel Demonstration State Forest, Santa Cruz County, California. (Draft) 1 p.

---. 2001. Relative Landslide Potential, Amaya and Fern Gulches, Soquel Demonstration State Forest, Santa Cruz County, California. (Draft) 1 p.

Heizer, R.F., ed. 1973. The Costanoan Indians. De Anza College, Local History Series 18. Cupertino, California.

Hester, R., M. McNally, G. Donaldson, and N. Nelson. 1993. Soquel Demonstration State Forest Recreation Study Final Report. Center for Environmental Design Research, University of California. Berkeley, California. 97 p.

---. Soquel Demonstration State Forest Recreation Study Supplemental Data Report. 80 p.

Hilton, S. and T.E. Lisle. 1993. Measuring the Fraction of Pool Volume Filled with Fine Sediment. USDA Forest Service, Pacific Southwest Research Station. Berkeley, California. 12 p.

Holland, V.L., M. Hanson, D. Keil, and L. Kelly. 1992. Preliminary Biological Assessment of Soquel Demonstration Forest. Santa Cruz County, California. Contract File 4904, California Polytechnic State University Sponsored Programs. 48 p.

Hunter, M.L. 1990. Wildlife, Forests and Forestry: Principles of Managing Forests for Biological Diversity. Regents/Prentice Hall, Englewood Cliffs, New Jersey.

Hyland, T. 2002. Botanical Survey of the Soquel Demonstration State Forest Timber Harvest Plan and Adjacent Areas. 7 p.

Hyland, T., and D. Neubauer. 2010. Fern Gulch THP Botanical Survey Soquel Demonstration State Forest. 10 p.

Jones and Stokes Associates, Inc. 1995. Environmental Impact Report for the Soquel Demonstration State Forest General Forest Management Plan. Draft. July 1995. (JSA 94-146.) Sacramento, California. Prepared for the California Department of Forestry and Fire Protection. Sacramento, California.

----. 1996. Environmental Impact Report for the Soquel Demonstration State Forest General Forest Management Plan. Final. June 1996. (JSA 94-146.) Prepared for the California Department of Forestry and Fire Protection. Sacramento, California.

Jones, T.L., N.E. Stevens, D.A. Jones, R.T. Fitzgerald, and M.G. Hylkema. 2007. The Central Coast: A Midlatitude Milieu. In *California Prehistory: Colonization, Culture, and Complexity*, edited by Terry L. Jones and Kathryn A. Klar, pp. 125-146. Altamira Press. New York.

Knopp, C. 1992. Monitoring plan for an assessment of indices for measuring the condition of cold water fish habitat. Unpublished report submitted to the North Coast Regional Water Quality Control Board as part of a contract with the California Department of Forestry and Fire Protection. 19 p.

Kroeber. A.L., 1925. *Handbook of the Indians of California*. Smithsonian Institution, Bureau of American Ethnology Bulletin 78. Reprint (1976). Dover Publications, New York.

Land Trust of Santa Cruz County. 2011. *A Conservation Blueprint: An Assessment and Recommendations from the Land Trust of Santa Cruz County*. Santa Cruz, California. 222 p.

Landers, R. 2013. Soquel Demonstration State Forest Bird Observations April 18, 2013, Amaya, Hihn's Mill and Sulphur Springs Roads. 1 p.

Lassette, N.S. and G.M. Kondolf. 2003. *Process Based Management of Large Woody Debris at the Basin Scale, Soquel Creek, California*. Final report prepared for the California Department of Forestry and Fire Protection and Soquel Demonstration State Forest. University of California, Berkeley, Department of Landscape Architecture and Environmental Planning. Berkeley, California. 130 p.

---. 2011. *Large Woody Debris in Urban Stream Channels: Redefining the Problem*. River Research and Applications 2011. 11 p.

Lee, S. 1991. *A Cruise Report of the Soquel Demonstration State Forest*. California Department of Forestry and Fires Protection. San Mateo-Santa Cruz Ranger Unit. 123 p.

- Levy, R.S. 1978. Costanoans in California, edited by R. F. Heizer, pp. 485-495. Handbook of North American Indians, Vol. 8. W. C. Sturtevant. Smithsonian Institution, Washington, DC.
- Lewis, J. and R.M. Rice. 1989. Critical sites erosion study, volume II: site conditions related to erosion on private timber lands in northern California. Final report submitted to the California Department of Forestry and Fire Protection. 95 p.
- Ligon, F., A. Rich, G. Rynearson, D. Thornburgh, and W. Trush. 1999. Report of the Scientific Review Panel on California Forest Practice Rules and Salmonid Habitat. Prepared for the Resources Agency of California and the National Marine Fisheries Service. Sacramento, California. 92 p. + appendices. http://resources.ca.gov/SRP_Rept.pdf
- Linsley and Kraeger Associates, Ltd. 1992. Summary report on stream flow and rainfall data from gauges in the Soquel Creek Watershed, July 1990-March 1992. Los Gatos, California.
- Lisle, T.E. and S. Hilton. 1992. The volume of fine sediment in pools: an index of sediment supply in gravel bed streams. *Water Resources Bulletin* 28(2): 371-383.
- Litschert, S.E. 2009. Predicting cumulative watershed effects in small forested watersheds. Ph.D. Dissertation. Colorado State University. Fort Collins, Colorado. 208 p.
- Loganbill, A.W. 2013. Post-fire Response of Little Creek Watershed: Evaluation of Change in Sediment Production and Suspended Sediment Transport. Master's Thesis, California Polytechnic State University. San Luis Obispo, California. 143 p.
- Manson, M.W. and J.A. Sowma-Bawcom. 1992. Geology, slope stability and earthquake damage in Soquel Demonstration State Forest. California Department of Conservation, Division of Mines and Geology. Open-file report 92-8. 65 p.
- Margolin, M. 1978. *The Ohlone Way: Indian Life in the San Francisco-Monterey Bay area.* Heyday Books. Berkeley, California.
- Mayer, K.E. and W.F. Laudenslayer, Jr., eds. 1988. *A guide to wildlife habitats of California.* Multiple-agency cooperative project for the California Department of Forestry and Fire Protection. 166 p.
- McPherson, B. no date. Soquel Demonstration State Forest Sudden Oak Death Research Study, transects and disease progression plots. 7 p.
- Milliken, R.T., 1995. *A Time of Little Choice: The Disintegration of Tribal Culture in the San Francisco Bay Area: 1769-1810.* Ballena Press. Menlo Park, California.
- Moratto, M.J., 1984. *California Archaeology.* Academic Press, New York.
- Moyle, P.B. 2002. *Inland Fishes of California.* University of California Press, Ltd. Berkeley, California.

National Marine Fisheries Service (NMFS). 2012. Recovery plan for central California coast coho salmon (*Oncorhynchus kisutch*) Evolutionary Significant Unit. National Marine Fisheries Service, Southwest Region. Santa Rosa, California. 338 p.
<http://swr.nmfs.noaa.gov/rcovey/index.htm>

Neubauer, D. 2013. Comstock Mill THP Botanical Survey Plant List, 14 May 2013. 12 p.

---. 2013. Comstock Mill THP Botanical Survey. 5 p.

Pacific Watershed Associates. 2003. Erosion Prevention Planning Project for County Roads and Roads in the Soquel Demonstration State Forest, in the Soquel Creek Watershed, Santa Cruz County, California. Summary Report prepared for the Santa Cruz County Resource Conservation District and the California Department of Fish and Game. Arcata, California. 31 p. plus appendices

Papa, M. 2011. Effects of Silviculture Management on Coast Redwood Forest Composition, Density and Structure in Santa Cruz and San Mateo Counties. Master's thesis for California Polytechnic University. San Luis Obispo, California. 275 p.

Payne, S. 1978. A Howling Wilderness: A History of the Summit Road Area of the Santa Cruz Mountains, 1850-1906. Loma Prieta Publishing. Los Gatos, California.

Poole, C. 1992. Stream channel inventory for the East Branch of Soquel Creek. Unpublished report submitted as a Senior Thesis, Department of Environmental Studies, University of California. Santa Cruz, California. 50 p.

Powell, R. n.d. Rancho de Palo de Yesca: the Shoquel Augmentation Rancho. Unpublished Soquel Demonstration State Forest Headquarters, Soquel, California. 168 p.

Punnett Brothers. 1906. Official Map of the County of Santa Cruz. State Lands Commission Grants of Land in California Made by Spanish or Mexican Authorities. Sacramento, California.

Reavis, B. 1991. Status of California Steelhead stocks. In Pacific States Marine Fisheries Commission and Association of Northwest Steelheaders (organizers), International Symposium on Steelhead Trout Management. Portland, Oregon. P. 57-63

Reynolds, S.D. 2013. Soquel Creek LWD Project: Background LWD Loading Rates. California Geologic Survey. 9 p.

---. 2013. Soquel Creek LWD Project – Site 1 As-Built Report. California Department of Conservation, California Geological Survey. Sacramento, California. 27 p.

---. 2013. Soquel Demonstration State Forest Watershed Enhancement Project Stream Habitat Improvement and Management; Large Woody Debris Augmentation, Large Woody Debris Site 1 and Related, Monitoring Summary – February, 2013. California Department of Conservation,

California Geological Survey. Sacramento, California. 11p.

---. 2014. Soquel Creek LWD Project – Site 2, 4, & 5 As-Built Report. California Department of Conservation, California Geological Survey. Sacramento, California. 78 p.

Rice, R.M. 1989. On-site effects: the necessary precursors of cumulative watershed effects. Unpublished paper presented at a Cumulative Impacts Training Session. Fort Bragg, California. 12 p.

---. 1993. A guide to data collection and analysis in support of an appraisal of cumulative watershed effects. Unpublished report prepared for private timber companies in California. Arcata, California. 29 p.

Rice, R.M., and J. Lewis. 1991. Estimating erosion risks associated with logging and forest roads in northwestern California. *Water Resources Bulletin* 27(5): 809-818.

----. 1992. Erosion risk and cumulative watershed effects associated with forest management in the upper Butano watershed, with special consideration of effects on the Pescadero Marsh. Unpublished report prepared for Big Creek Lumber Company. Davenport, California. 40 p.

----. 1992. A guide to data collection and analysis in support of an appraisal of cumulative watershed effects. Unpublished report prepared for private timber companies in California. 29 p.

Santa Cruz County Planning Department. 1994. County of Santa Cruz 1994 General Plan and Local Coastal Program Land Use Plan. Santa Cruz, California.

Santa Cruz County Resource Conservation District (SCCRCD). 2003. The Soquel Creek watershed assessment and enhancement project plan. Capitola, California. 92 p.

Sillett, S.C. et al. 2010. Increasing wood production through old age in tall trees. *Forest Ecology and Management*, 259 (5): 976-994.
<http://www.sciencedirect.com/science/article/pii/S037811270900872X>

Singer, S. and M.L. Swanson. 1983. Soquel Creek storm damage recovery plan: a reconnaissance level study with recommendations for watershed management. Unpublished report of USDA Soil Conservation Service. Aptos, California. 71 p.

Smith, J.J. 2005. Unpublished Data from Scale Analysis of Down-migrant Smolts on the San Lorenzo River, 1987-89. Department of Biological Sciences. San Jose State University.

----. 2008. Steelhead Growth in a Small Central California Watershed: Upstream and Estuarine Rearing Patterns. *Transactions of the American Fisheries Society*, 137:114-128.

Sogard, S.M., T.H. Williams, and H. Fish. 2009. Seasonal Patterns of Abundance, Growth, and Site Fidelity of Juvenile Steelhead in a Small Coastal California Stream. *Transactions of the American Fisheries Society*, 138: 549-563

State Emergency Assessment Team (SEAT). 2008 Summit Fire State Emergency Assessment Team Report affecting watershed in Santa Cruz and Santa Clara Counties. Sacramento, California 130 p.

State Lands Commission. 1984. Grants of Land in California Made by Spanish or Mexican Authorities. Sacramento, California.

Sturm, E., E. Gilbert-Horvath, J. Garza, and R. MacFarlane. 2009. Creation of a captive broodstock program for southern coho salmon (*Oncorhynchus kisutch*); results from the initial rearing and spawning of the first brood year. NOAA-TM-NMFS-SWFSC-436.

Suddjian, D.L. 2001. Spring 2001 Raptor survey at the proposed Fern Gulch THP area. 18 p.

----. 2001. A Tale of Ten Snags. A Forest Management Success Story. The Albatross, Santa Cruz Bird Club Newsletter Vol. 47, No 1: Sep/Oct 01. 2 p.

----. 2003. Results of 2003 Marbled Murrelet Surveys at Soquel Demonstration State Forest, Santa Cruz County, California. 7 p.

----. 2004. Results of 2003-2004 Marbled Murrelet Surveys at Soquel Demonstration State Forest, Santa Cruz County, California. 8 p.

----. 2005. Birds of Soquel Creek Riparian Habitat. Notes from presentation to Friends of Soquel Creek meeting January 26, 2005. 2 p.

----. 2005. 2005 Raptor Survey Fern Gulch THP, Soquel Demonstration State Forest, Santa Cruz County, California. 12 p. plus map 1 p.

----. 2006. 2006 Raptor Survey Fern Gulch THP, Soquel Demonstration State Forest, Santa Cruz County, California. 14 p. plus map 1 p.

Thomas, J. H. 1961. Flora of the Santa Cruz Mountains of California. Stanford University Press. Palo Alto, California. 433 p.

Titus, R.G., D.C. Erman, and W.M. Snider. 2005. History and status of steelhead in California coastal drainages south of San Francisco. In preparation. California Department of Fish and Game. Sacramento, California.

University of California, Berkeley Center for Environmental Design Research. 1996. Draft Education and Recreation Master Plan for Soquel Demonstration State Forest. Berkeley, California. 75 p.

U.S. Department of Agriculture, Natural Resources Conservation Service. 2004. Soil Survey Geographic (SSURGO) database for Santa Cruz County, California. 148 p.

U.S. Department of the Interior, National Park Service. 1991. How to Apply the National Register Criteria for Evaluation. National Register Bulletin 15. Washington, D.C.

Valentine, S. 1991. "Listening" for the Soquel Demonstration State Forest. Final Report for a University of California Berkeley course Interdepartmental Studies IDS 206. 22 p.

PERSONAL COMMUNICATION

Anderson, T. and J. Brown. July, 1994. Habitat Surveyors, California Polytechnic State University.

Mason, P. June, 1992. Local resident.

McCaslin, M. August, 1993. Representative of the Monterey Bay Salmon and Trout Project.

McCrary, B. May, 2003. Owner, Big Creek Lumber Company.

Smith, J. February, 1993. Biologist, San Jose State University.

Wilson, N. July, 1993. Representative of the Fungus Federation.

APPENDICES

APPENDIX A: TEXT OF ASSEMBLY BILL 1965

ASSEMBLY BILL NO. 1965

CHAPTER 1338

An act to add Article 4 (commencing with Section 4660) to Chapter 9 of Division 4 of the Public Resources Code, relating to state forests.

(Approved by Governor September 29, 1987.
Filed with Secretary of State September 29, 1987.)

LEGISLATIVE COUNSEL'S DIGEST

AB 1965, Farr. Soquel Demonstration State Forest.

Under existing law, the Department of Forestry and Fire Protection is authorized to engage in the management, protection, and restoration of state forests in accordance with plans approved by the State Board of Forestry and Fire Protection.

This bill would declare it the policy of the state to establish the Soquel Demonstration State Forest in Santa Cruz County. The department would be responsible for the development and establishment of the state forest and for maintenance and operations. The bill would authorize the department to permit a limited amount of commercial timber operations within the forest in order to provide funds for its maintenance and operation. The bill would require the department, in conjunction with a specified advisory committee, to adopt a general plan for the park by January 1, 1989, and would require the general plan to be approved by the advisory committee prior to adoption by the department.

The department's duties and authority under the bill would only arise if the state acquires the property comprising the demonstration forest.

The people of the State of California do enact as follows:

SECTION 1. Article 4 (commencing with Section 4660) is added to Chapter 9 of Division 4 of the Public Resources Code, to read:

ARTICLE 4. SOQUEL DEMONSTRATION STATE FOREST

4660. It is hereby declared to be the policy of the state to establish and preserve an intensively managed, multifaceted research forest which is representative of forest activities as a living forest in Santa Cruz County within northern California's coastal redwood belt. The coast redwoods, as the dominant tree species in this area, are a valuable natural resource and are unique in North America for their beauty, abundance, diversity, and public accessibility, and their extreme beauty and economic value requires special measures for their protection for the use, enjoyment, and education of the public.

It is the intent of the Legislature, in establishing the Soquel Demonstration State Forest, to provide an environment that will do all of the following:

- (a) Provide watershed protection for local communities and base-line monitoring and studies of the hazards, risks, and benefits of forest operations and watersheds to urban areas.
- (b) Provide public education and examples illustrating compatible rural land uses, including sustained yield timber production, as well as the historic development of timbering and forestry machinery, within the context of local community protection and nearby pressures.
- (c) Provide a resource for the public, environmental groups, elected officials, environmental planners, the educational community, and the media as an open environment for the inspection and study of environmental education, forestry practices, and effects thereof.
- (d) Protect old growth redwood trees.

4661. The department may permit a limited amount of commercial timber operations on the property within the Soquel Demonstration State Forest in order to provide funds for the maintenance and operation of the state forest and to allow fulfillment of the objectives of Section 4660. Income from the state forest property shall sustain all costs of operation and provide income for research and educational purposes.

4662. The department is responsible for the development and establishing of the Soquel Demonstration State Forest and for ongoing maintenance and operations. The director shall appoint an advisory committee to assist the department in planning future management of the forest. The advisory committee shall include representatives of the Santa Cruz County Board of Supervisors, the Department of Parks and Recreation, the Board of Forestry and Fire Protection, the Forest of Nisene Marks Advisory Committee, and the Department of Fish and Game.

4663. The department, in coordination with the advisory committee, shall adopt by January 1, 1989, a general plan for the state forest which reflects the long-range development and management plans to provide for the optimum use and enjoyment of the living forest, as provided in Section 4660, as well as the protection of its quality and the watershed within the Santa Cruz area. The general plan shall be approved by the advisory committee prior to adoption by the department.

4664. The duties and authority of the department pursuant to this article shall only arise if the state acquires the property comprising the Soquel Demonstration State Forest.

APPENDIX B: SPECIES LISTS FOR SDSF

FAUNA OF SDSF

From Preliminary Biological Assessment of Soquel Demonstration State Forest, Santa Cruz County, California (Holland et al., 1992) and David Suddjian, Biological Consultant.

SCIENTIFIC NAME

COMMON NAME

FISH

<i>Lampetra tridentata</i>	Pacific lamprey
<i>Oncorhynchus mykiss iridius</i>	steelhead trout
<i>Oncorhynchus mykiss</i>	rainbow trout
<i>Gasterosteus aculeatus</i>	northern threespine stickleback
<i>Cottus asper</i>	prickly sculpin
<i>Catostomus occidentalis</i>	Sacramento sucker

AMPHIBIANS

Lawrence Erickson, herpetologist and CAL FIRE Fire Captain, updated the nomenclature for amphibian and reptile species previously observed. He also conducted on-going surveys, resulting in observations of some new species on the Forest.

<i>Taricha torosa torosa</i>	California newt, coast range newt
<i>Taricha granulosa granulosa</i>	rough-skinned newt
<i>Dicamptodon ensatus</i>	Pacific giant salamander, California giant salamander
<i>Batrachoseps attenuatus</i>	California slender salamander
<i>Ensatina eschscholtzii xanthoptica</i>	yellow-eyed ensatina
<i>Aneides flavipunctatus niger</i>	Santa Cruz black salamander
<i>Aneides lugubris</i>	arboreal salamander
<i>Bufo boreas halophilus</i>	California toad, western toad
<i>Pseudacris sierra</i>	Sierran tree frog, Pacific chorus frog
<i>Rana boylei</i>	foothill yellow-legged frog
<i>Rana draytonii</i>	California red-legged frog

REPTILES

<i>Actinomyx marmorata pallida</i>	western pond turtle, southern Pacific pond turtle
<i>Sceloporus occidentalis boucurtii</i>	western fence lizard, coast range fence lizard
<i>Plestiodon skiltonianus skiltonianus</i>	Skilton's skink, western skink
<i>Elgaria coerulea coerulea</i>	northern alligator lizard, San Francisco alligator lizard

<i>Elgaria multicaranata multicaranata</i>	southern alligator lizard, California alligator lizard
<i>Charina bottae</i>	northern rubber boa
<i>Diadophis punctatus amabilis</i>	Pacific ring-necked snake
<i>Contia tenuis</i>	sharp-tailed snake
<i>Pituophis catenifer catenifer</i>	Pacific gopher snake
<i>Lampropeltis getula californiae</i>	California kingsnake
<i>Lampropeltis zonata multifasciata</i>	California mountain kingsnake, coast mountain kingsnake
<i>Thamnophis sirtalis infernalis</i>	California red-sided garter snake, red-spotted garter snake
<i>Thamnophis elegans terrestris</i>	western terrestrial garter snake, coast garter snake
<i>Thamnophis atratus atratus</i>	Santa Cruz garter snake
<i>Coluber constrictus flaviventris</i>	western yellow-bellied racer, western racer
<i>Crotalus oreganus oreganus</i>	northern Pacific rattlesnake, western rattlesnake

BIRDS

<i>Ardea herodias</i>	great blue heron
<i>Butorides virescens</i>	green heron
<i>Aix sponsa</i>	wood duck
<i>Mergus merganser</i>	common merganser
<i>Anas platyrhynchos</i>	mallard
<i>Cathartes aura</i>	turkey vulture
<i>Pandion haliaetus</i>	osprey
<i>Accipiter striatus</i>	sharp-shinned hawk
<i>Accipiter cooperii</i>	Cooper's hawk
<i>Buteo jamaicensis</i>	red-tailed hawk
<i>Buteo lineatus</i>	red-shouldered hawk
<i>Aquila chrysaetos</i>	golden eagle
<i>Falco columbarius</i>	merlin
<i>Callipepla californica</i>	California quail
<i>Actitis macularia</i>	spotted sandpiper
<i>Patagioenas fasciata</i>	band-tailed pigeon
<i>Zenaida macroura</i>	mourning dove
<i>Megascops kennicottii</i>	western screech owl
<i>Bubo virginianus</i>	great horned owl
<i>Glaucidium gnoma</i>	northern pygmy-owl
<i>Asio otus</i>	long-eared owl
<i>Aegolius acadicus</i>	northern saw-whet owl
<i>Calypte anna</i>	Anna's hummingbird
<i>Selasphorus rufus</i>	rufous hummingbird
<i>Selasphorus sasin</i>	Allen's hummingbird
<i>Megaderyle alcyon</i>	belted kingfisher
<i>Melanerpes formicivorus</i>	acorn woodpecker
<i>Sphyrapicus ruber</i>	red-breasted sapsucker
<i>Picoides pubescens</i>	downy woodpecker

<i>Picoides villosus</i>	hairy woodpecker
<i>Colaptes auratus</i>	northern flicker
<i>Empidonax difficilis</i>	Pacific-slope flycatcher
<i>Sayornis nigricans</i>	black phoebe
<i>Myiarchus cinerascens</i>	ash-throated flycatcher
<i>Contopus cooperi</i>	olive-sided flycatcher
<i>Contopus sordidulus</i>	western wood-peewee
<i>Cypseloides niger</i>	black swift
<i>Chaetura vauxi</i>	Vaux's swift
<i>Tachycineta thalassina</i>	violet-green swallow
<i>Stelgidopteryx serripennis</i>	northern rough-winged swallow
<i>Hirundo pyrrhonota</i>	cliff swallow
<i>Corvus corax</i>	common raven
<i>Cyanocitta stelleri</i>	Steller's jay
<i>Aphelocoma californica</i>	western scrub-jay
<i>Poecile rufescens</i>	chestnut-backed chickadee
<i>Baeolofus inornatus</i>	oak titmouse
<i>Psaltriparus minimus</i>	bushtit
<i>Certhia americana</i>	brown creeper
<i>Sitta pygmaea</i>	pygmy nuthatch
<i>Troglodytespacificus</i>	Pacific wren
<i>Thryomanes bewickii</i>	Bewick's wren
<i>Chamaea fasciata</i>	wrentit
<i>Cinclus mexicanus</i>	American dipper
<i>Regulus calendula</i>	ruby-crowned kinglet
<i>Regulus satrapa</i>	golden-crowned kinglet
<i>Polioptila caerulea</i>	blue-gray gnatcatcher
<i>Toxostoma redivivum</i>	California thrasher
<i>Catharus ustulatus</i>	Swainson's thrush
<i>Catharus guttatus</i>	hermit thrush
<i>Turdus migratorius</i>	American robin
<i>Ixoreus naevius</i>	varied thrush
<i>Bombycilla cedrorum</i>	cedar waxwing
<i>Sturnus vulgaris</i>	European starling
<i>Vireo cassinii</i>	Cassin's vireo
<i>Vireo huttoni</i>	Hutton's vireo
<i>Vireo gilvus</i>	warbling vireo
<i>Setophaga coronata</i>	yellow-rumped warbler
<i>Setophaga townsendi</i>	Townsend's warbler
<i>Dendroica occidentalis</i>	hermit warbler
<i>Setophaga nigrescens</i>	black-throated gray warbler
<i>Setophaga petechia</i>	yellow warbler
<i>Oreothlypis celata</i>	orange-crowned warbler
<i>Oreothlypis ruficapilla</i>	Nashville warbler
<i>Geothlypis tolmiei</i>	MacGillivray's warbler
<i>Cardellina pusilla</i>	Wilson's warbler

<i>Piranga ludoviciana</i>	western tanager
<i>Passerella iliaca</i>	fox sparrow
<i>Melospiza melodia</i>	song sparrow
<i>Melospiza lincolni</i>	Lincoln's sparrow
<i>Zonotrichia atricapilla</i>	golden-crowned sparrow
<i>Zonotrichia leucophrys</i>	white-crowned sparrow
<i>Pipilo maculatus</i>	spotted towhee
<i>Melospiza crissalis</i>	California towhee
<i>Junco hyemalis</i>	dark-eyed junco
<i>Pheucticus melanocephalus</i>	black-headed grosbeak
<i>Loxia curvirostra</i>	red crossbill
<i>Haemorhous purpureus</i>	purple finch
<i>Coccothraustes vespertinus</i>	evening grosbeak
<i>Spinus tristis</i>	American goldfinch
<i>Spinus psaltria</i>	lesser goldfinch
<i>Spinus pinus</i>	pine siskin

The following additional bird observations were reported by Matt Greene in his 2012 reports on "Raptor Surveys for Fern Gulch THP" and "Rim THP Raptor Survey". These reports also confirmed sightings of many of the species observed previously.

SCIENTIFIC NAME

COMMON NAME

Meleagris gallopavo
Dryocopus pileatus

wild turkey
pileated woodpecker

MAMMALS

<i>Didelphis virginiana</i>	opossum
<i>Sorex trowbridgii</i>	Trowbridge's shrew
<i>Neurotrichus gibbsii</i>	shrew-mole
<i>Scapanus latimanus</i>	broad-footed mole
<i>Myotis lucifugus</i>	little brown myotis
<i>Myotis evotis</i>	long-eared myotis
<i>Myotis volans</i>	long-legged myotis
<i>Myotis californicus</i>	California myotis
<i>Lasionycteris noctivagans</i>	silver-haired bat
<i>Lasiurus cinereus</i>	hoary bat
<i>Pipistrellus hesperus</i>	western pipistrelle
<i>Eptesicus fuscus</i>	big brown bat
<i>Sylvilagus bachmani</i>	brush rabbit
<i>Neotamias merriami</i>	Merriam's chipmunk
<i>Sciurus griseus</i>	western gray squirrel
<i>Thomomys bottae</i>	Botta's pocket gopher
<i>Chaetodipus californicus</i>	California pocket mouse
<i>Reithrodontomys megalotis</i>	western harvest mouse
<i>Peromyscus truei</i>	pinyon mouse
<i>Peromyscus californicus</i>	California mouse
<i>Peromyscus maniculatus</i>	deer mouse
<i>Neotoma fuscipes</i>	dusky-footed woodrat
<i>Urocyon cinereoargenteus</i>	gray fox
<i>Procyon lotor</i>	raccoon
<i>Mustela frenata</i>	long-tailed weasel
<i>Mephitis mephitis</i>	striped skunk
<i>Canis latrans</i>	coyote
<i>Puma concolor</i>	mountain lion
<i>Lynx rufus</i>	bobcat
<i>Sus scrofa</i>	feral pig
<i>Odocoileus hemionus</i>	black-tailed deer

FLORA OF SDSF

From Preliminary Biological Assessment of Soquel Demonstration State Forest, Santa Cruz County, California (Holland et al., 1992) and Randy Morgan; the report for the “botanical survey for the Fern Gulch THP”, submitted by Tim Hyland on May 2, 2010; and the report for the botanical survey for the Comstock Mill THP, submitted by Dylan Neubauer on May 14, 2013.

* non-native

** most invasive

<i>Acer macrophyllum</i>	bigleaf maple
<i>Acer negundo</i>	California boxelder
<i>Achillea millefolium</i>	yarrow
<i>Acmispon americanus</i>	Spanish clover
<i>Acmispon americanus</i> var. <i>a.</i>	Spanish trefoil
<i>Acmispon glaber</i> var. <i>g.</i>	deerweed
<i>Acmispon heermannii</i> var. <i>orbicularis</i>	Heermann's trefoil
<i>Acmispon parviflorus</i>	small-flowered trefoil
<i>Adenocaulon bicolor</i>	trail plant
<i>Adenostema fasciculatum</i>	chamise
<i>Adenostoma fasciculatum</i> var. <i>f.</i>	chamise
<i>Adiantum aleuticum</i>	five-finger fern
<i>Adiantum jordanii</i>	California maidenhair
<i>Aesculus californica</i>	California buckeye
** <i>Ageratina adenophora</i>	eupatorium
<i>Agoseris grandiflora</i>	large-flowered Agoseris
<i>Agrostis avenacea</i>	Pacific bentgrass
<i>Agrostis hallii</i>	Hall's bentgrass
<i>Agrostis stolonifera</i>	western bentgrass
<i>Agrostis pallens</i>	leafy bentgrass
* <i>Aira caryophyllea</i>	silvery hairgrass
<i>Allium unifolium</i>	one-leaved onion
<i>Allophyllum divaricatum</i>	straggling gilia
<i>Alnus rhombifolia</i>	white alder
<i>Alnus rubra</i>	red alder
<i>Amsinckia intermedia</i>	common fiddleneck
* <i>Anagallis arvensis</i>	scarlet pimpernel
<i>Anaphalus margaritacea</i>	pearly everlasting
<i>Anisocarpus madioides</i>	woodland Madia
* <i>Anthemis cotula</i>	mayweed
<i>Anthriscus caucalis</i>	knotted hedge parsley
<i>Anthoxanthum occidentale</i>	vanilla grass
<i>Aquilegia formosa</i>	northwest crimson columbine
<i>Aralia californica</i>	California spikenard
<i>Arbutus menziesii</i>	madrone
<i>Arctostaphylos andersonii</i>	Santa Cruz mountain manzanita
<i>Arctostaphylos crustacea</i>	brittle-leaved manzanita

<i>Arctostaphylos crustacea</i> subsp. <i>crinita</i>	crinite manzanita
<i>Arctostaphylos</i> sp.	manzanita
<i>Arctostaphylos tomentosa</i> ssp. <i>crinita</i>	brittle-leaved manzanita
<i>Artemisia douglasiana</i>	mugwort
<i>Asarum caudatum</i>	wild ginger
<i>Asyneuma prenanthoides</i>	bellflower
<i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	lady fern
* <i>Avena fatua</i>	common wild oats
<i>Baccharis glutinosa</i>	marsh baccharis
<i>Baccharis pilularis</i> ssp. <i>Consanguinea</i>	coyote brush
<i>Barbarea</i> sp.	wintercress
<i>Boisduvalia densiflora</i>	dense-flowered Boisduvalia
<i>Boykinia occidentalis</i>	coast Boykinia
* <i>Brassica nigra</i>	black mustard
* <i>Briza maxima</i>	rattlesnake grass
<i>Briza minor</i>	little quaking grass
<i>Brodiaea elegans</i>	harvest brodiaea
<i>Bromus carinatus</i>	California brome
<i>Bromus carinatus</i> var. <i>c.</i>	California brome
* <i>Bromus diandrus</i>	riggut brome
* <i>Bromus hordeaceus</i>	soft chess brome
<i>Bromus laevipes</i>	woodland brome
* <i>Bromus madritensis</i>	red brome
<i>Bromus vulgaris</i>	woodland brome
<i>Calamagrostis rubescens</i>	wild morning glory
<i>Calandrinia breweri</i>	Brewer's redmaids
<i>Callitriche marginata</i>	California water starwort
<i>Calochortus albus</i>	fairy lantern
<i>Calystegia occidentalis</i>	pine grass
* <i>Calystegia occidentalis</i>	coast morning glory
<i>Calystegia purpurata</i> subsp. <i>p.</i>	western morning glory
<i>Cardamine californica</i>	milkmaids
<i>Cardamine oligosperma</i>	few-seeded bittercress
** <i>Carduus pycnocephalus</i> subsp. <i>p.</i>	Italian thistle
* <i>Carduus pycnocephalus</i>	Italian thistle
* <i>Carduus tenuiflorus</i>	slender-flowered thistle
<i>Carex barbarae</i>	Santa Barbara sedge
<i>Carex bolanderi</i>	Bolander's sedge
<i>Carex globosa</i>	round-fruited sedge
<i>Carex harfordii</i>	Harford's sedge
<i>Carex tumulicola</i>	foothill sedge
<i>Ceanothus cuneatus</i>	buck brush
<i>Ceanothus papillosus</i>	warty-leaved ceanothus
<i>Ceanothus thyrsiflorus</i>	blue brush
<i>Ceanothus thyrsiflorus</i> var. <i>t.</i>	blue-blossom
* <i>Cedrus deodara</i>	Deodar cedar

* <i>Centaurea melitensis</i>	tocalote
* <i>Centaurea solstitialis</i>	yellow star thistle
<i>Cerastium glomeratum</i>	mouse-eared chickweed
<i>Chlorogalum pomeridianum</i>	soap plant
<i>Chlorogalum pomeridianum</i> var. <i>p.</i>	soap plant
<i>Cirsium brevistylum</i>	Indian thistle
* <i>Cirsium vulgare</i>	bull thistle
<i>Clarkia purpurea</i> ssp. <i>viminea</i>	large godetia
<i>Claytonia parviflora</i> subsp. <i>p.</i>	small-flowered claytonia
<i>Claytonia perfoliata</i>	miners' lettuce
<i>Clinopodium douglasii</i>	yerba buena
<i>Clintonia andrewsiana</i>	red clintonia
<i>Collomia heterophylla</i>	varied-leaved Collomia
* <i>Conium maculatum</i>	poison hemlock
<i>Convolvulus arvensis</i>	bindweed
<i>Corallorhiza maculate</i> var. <i>occidentalis</i>	unspotted spotted coralroot
* <i>Cordyline</i> sp.	cordyline
<i>Cornus sericia</i> ssp. <i>californica</i>	western red or creek dogwood
<i>Cornus sericia</i> ssp. <i>sericia</i>	red osier dogwood
* <i>Cortaderia jubata</i>	pampas grass
<i>Corylus cornuta</i>	California hazel
<i>Corylus cornuta</i> var. <i>californica</i>	California hazel
* <i>Cotoneaster pannosus</i>	silverleaf cotoneaster
<i>Cryptantha micromeres</i>	minute-flowered cryptantha
<i>Cynoglossum grande</i>	hound's-tongue
* <i>Cynosurus echinatus</i>	dog's-tail grass
<i>Cyperus eragrostis</i>	tall cyperus
* <i>Cytisus scoparius</i>	Scotch broom
* <i>Dactylis glomerata</i>	orchard grass
<i>Danthonia californica</i>	California oat grass
<i>Deschampsia elongata</i>	slender hairgrass
<i>Dichelostemma pulchellum</i>	blue dicks
* <i>Dichondra micrantha</i>	Asian pony's foot
<i>Drymocallis glandulosa</i>	cinquefoil
<i>Dryopteris arguta</i>	coastal wood fern
<i>Ehrharta erecta</i>	panic veldt grass
<i>Elymus glaucus</i>	western ryegrass
<i>Elymus glaucus</i> subsp. <i>g.</i>	blue wildrye
<i>Epilobium brachycarpum</i>	annual willowherb
<i>Epilobium ciliatum</i> ssp. <i>ciliatum</i>	California willow herb
<i>Epilobium ciliatum</i> subsp. <i>watsonii</i>	Watson's willow herb
<i>Epipactis helleborine</i>	eastern orchid
<i>Equisetum hiemale</i>	scouring rush
<i>Equisetum hyemale</i> ssp. <i>affine</i>	common scouring rush
<i>Equisetum talmateia</i> ssp. <i>Braunii</i>	giant horse tail
<i>Equisetum telmateia</i>	horsetail

<i>Ericameria arborescens</i>	golden fleece
<i>Erigeron canadensis</i>	horseweed
* <i>Erigeron sumatrensis</i>	wide-leave horseweed
<i>Eriodictyon californicum</i>	California mountain balm
<i>Eriophyllum confertiflorum</i> var. <i>c.</i>	golden yarrow
* <i>Euphorbia peplus</i>	petty spurge
<i>Eurybia radulina</i>	broad-leaved wood aster
<i>Festuca arundinacea</i>	tall fescue
<i>Festuca bromoide</i>	six-weeks fescue
<i>Festuca californica</i>	California fescue
<i>Festuca myuros</i>	rattail fescue
<i>Festuca occidentalis</i>	western fescue
<i>Festuca perennis</i>	annual rye grass
<i>Festuca rubra</i>	red fescue
<i>Festuca subuliflora</i>	crinkle-awned fescue
* <i>Filago gallica</i>	daggerleaf cottonrose
<i>Fragaria vesca</i>	California strawberry
<i>Frangula californica</i>	California coffeeberry
<i>Frangula californica</i> subsp. <i>c.</i>	California coffeeberry
<i>Fritillariaaffinis</i> var. <i>affinis</i>	checker lily
* <i>Galium aparine</i>	goosegrass
<i>Galium californicum</i>	California bedstraw
<i>Galium californicum</i> subsp. <i>c.</i>	California bedstraw
* <i>Galium murale</i>	tiny bedstraw
* <i>Galium parisiense</i>	Parisian bedstraw
<i>Galium porrigens</i>	climbing bedstraw
<i>Galium triflorum</i>	trifid bedstraw
<i>Gamochaeta ustulata</i>	California everlasting
* <i>Gastridium phleoides</i>	nitgrass
* <i>Genista monspessulana</i>	French broom
<i>Geranium dissectum</i>	cut-leaved geranium
<i>Geranium molle</i>	dove's-foot geranium
<i>Gnaphalium luteo-album</i>	weedy cudweed
<i>Gnaphalium purpureum</i>	purple cudweed
<i>Gnaphalium ramossissimum</i>	pink everlasting
* <i>Hedera helix</i>	English ivy
<i>Helenium puberulum</i>	sneezeweed
<i>Heracleum maximum</i>	cow parsnip
<i>Heteromeles arbutifolia</i>	toyon
<i>Heuchera micrantha</i>	small-flowered heuchera
<i>Hieracium albiflorum</i>	white-flowered hawkweed
<i>Hierochloe occidentalis</i>	vanilla grass
<i>Hirschfeldia incana</i>	perennial mustard
* <i>Holcus lanatus</i>	velvet grass
<i>Holodiscus discolor</i>	cream bush
<i>Holta macrostachya</i>	leather root

<i>Hordeum brachyantherum</i> subsp. <i>b.</i>	California barley
* <i>Hordeum murinum</i> ssp. <i>leporinum</i>	foxtail barley
* <i>Hypericum calycinum</i>	St. John's wort
* <i>Hypochaeris glabra</i>	smooth cat's-ear
* <i>Hypochaeris radicata</i>	hairy cat's ear
<i>Iris ferneldii</i>	Santa Cruz Mountain iris
<i>Iris macrosiphon</i>	ground iris
<i>Juglans</i> sp.	walnut
<i>Juncus bufonius</i>	toad rush
<i>Juncus effusus</i> var. <i>brunneus</i>	bog rush
<i>Juncus effusus</i> var. <i>pacificus</i>	Pacific rush
<i>Juncus patens</i>	common rush
<i>Juncus xiphioides</i>	iris-leaved rush
* <i>Lathyrus latifolius</i>	perennial sweet pea
<i>Lathyrus vestitus</i>	common Pacific pea
<i>Lathyrus vestitus</i> var. <i>vestitus</i>	wood-pea
* <i>Lepidium strictum</i>	wayside pepper grass
* <i>Linum bienne</i>	small-flowered flax
<i>Logfia filaginoides</i>	California Filago
<i>Lonicera hispidula</i>	hairy honeysuckle
<i>Lotus eriophorus</i>	wooly trefoil
<i>Lotus scoparius</i>	deerweed
<i>Lupinus albifrons</i>	bush lupine
<i>Lupinus latifolius</i>	broad-leaved lupine
<i>Lupinus latifolius</i> var. <i>l.</i>	broad-leaved lupine
<i>Lupinus nanus</i>	sky lupine
<i>Luzula comosa</i> var. <i>c.</i>	common wood rush
<i>Luzula multiflora</i>	wood rush
<i>Madia elegans</i>	common Madia
<i>Madia gracilis</i>	slender tarweed
* <i>Madia sativa</i>	coast tarweed
<i>Maianthemum racemosum</i>	western Solomon's seal
<i>Maianthemum stellatum</i>	slender Solomon's seal
* <i>Marah fabaceus</i>	wild cucumber
* <i>Matricaria discoidea</i>	pineapple weed
* <i>Medicago polymorpha</i>	bur clover
<i>Melica imperfecta</i>	small-flowered melic grass
<i>Melica subulata</i>	Alaska onion grass
<i>Melica torreyana</i>	Torrey's melic grass
* <i>Melilotus albus</i>	white sweetclover
* <i>Melilotus indicus</i>	yellow sweetclover
* <i>Melissa officinalis</i>	lemon balm
<i>Mimulus aurantiacus</i>	sticky monkeyflower
<i>Mimulus aurantiacus</i> var. <i>a.</i>	sticky monkeyflower
<i>Mimulus cardinalis</i>	scarlet monkeyflower
<i>Mimulus guttatus</i>	common monkeyflower

<i>Morella californica</i>	wax myrtle
* <i>Myosotis latifolia</i>	forget-me-not
* <i>Nasturtium officinale</i>	watercress
<i>Navarretia squarrosa</i>	skunkweed
<i>Nemophila parviflora</i>	small-flowered Nemophila
<i>Nemophila parviflora</i> var. <i>p.</i>	small-flowered nemophila
<i>Nemophila pedunculata</i>	meadow nemophila
<i>Notholithocarpus densiflora</i>	tanoak
<i>Oemeria cerasiformis</i>	oso berry
<i>Osmorhiza berteroi</i>	sweet cicely
<i>Oxalis albicans</i>	hairy wood sorrel
<i>Oxalis oregana</i>	redwood sorrel
<i>Oxalis pes-caprae</i>	sour grass
<i>Oxalis pilosa</i>	hairy wood sorrel
<i>Pellaea andromedifolia</i>	coffee fern
<i>Pentagramma triangularis</i>	
ssp. <i>triangularis</i>	goldenback fern
<i>Persicaria punctate</i>	smartweed
<i>Petasites frigidus</i> var. <i>palmatus</i>	coltsfoot
<i>Phalaris californica</i>	California canary grass
<i>Pinus attenuata</i>	knobcone pine
* <i>Pinus coulteri</i>	Coulter pine
* <i>Pinus pinea</i>	Italian stone pine
* <i>Pinus radiata</i>	Monterey pine
<i>Piperia elongata</i>	rein orchid
<i>Pityrogramma triangularis</i> ssp. <i>Triangularis</i>	gold-backed fern
* <i>Plantago lanceolata</i>	English plantain
<i>Plantanus racemosa</i>	California sycamore
<i>Poa howellii</i>	Howell's bluegrass
<i>Polygala californica</i>	California milkwort
<i>Polypodium californicum</i>	California polypody
<i>Polypodium calirhiza</i>	polypody
* <i>Polypogon interruptus</i>	beard grass
* <i>Polypogon monspeliensis</i>	rabbitsfoot grass
<i>Polypogon viridis</i>	water beard grass
<i>Polystichum dudleyi</i>	Dudley's shield fern
<i>Polystichum munitum</i>	western sword fern
<i>Populus trichocarpa</i>	black cottonwood
<i>Potentilla glandulosa</i>	sticky potentilla
<i>Prosartes hookeri</i>	Hooker's fairy bells
<i>Pseudognaphalium californicum</i>	California everlasting
<i>Psudeognaphalium luteoalbum</i>	weedy cudweed
<i>Pseudognaphalium</i> sp.	everlasting
<i>Pseudognaphalium ramossissimum</i>	pink everlasting
<i>Pseudognaphalium stramineum</i>	cotton-batting plant
<i>Pseudotsuga menziesii</i>	Douglas-fir

<i>Psilocarphus tenellus</i>	slender woolly marbles
<i>Pteridium aquilinum</i>	bracken fern
<i>Pteridium aquilinum</i> var. <i>pubescens</i>	bracken fern
<i>Quercus agrifolia</i>	coast live oak
<i>Quercus agrifolia</i> var. <i>a.</i>	coast live oak
<i>Quercus agrifolia</i> x <i>kelloggii</i>	hybrid oak
<i>Quercus kelloggii</i>	black oak
<i>Quercus parvula</i> var. <i>shrevei</i>	Shreve oak
<i>Quercus parvula</i> var. <i>shrevei</i> x <i>kelloggii</i>	hybrid oak
<i>Quercus wislizenii</i>	interior live oak
<i>Ranunculus muricatus</i>	prickle-fruited ranunculus
<i>Ribes menziesii</i>	canyon gooseberry
<i>Ribes menziesii</i> var. <i>senile</i>	Santa Cruz gooseberry
<i>Ribes sanguineum</i> var. <i>glutinosum</i>	flowering current
<i>Rorippa palustris</i>	yellow cress
<i>Rosa californica</i>	California wild rose
<i>Rosa gymnocarpa</i>	wood rose
<i>Rosa spithamea</i>	ground rose
<i>Rubus leucodermis</i>	western raspberry
<i>Rubus parviflorus</i>	thimble berry
<i>Rubus ursinus</i>	California blackberry
* <i>Rumex acetosella</i>	sour dock or sheep sorrel
* <i>Rumex conglomeratus</i>	clustered dock
<i>Rumex salicifolius</i>	willow dock
<i>Rupertia physodes</i>	Rupert's scurf-pea
<i>Sagina apetala</i>	sticky pearlwort
<i>Salix lasiandra</i>	yellow willow
<i>Salix lasiolepis</i>	sandbar willow
<i>Salix sitchensis</i>	velvet willow
<i>Sambucus nigra</i> ssp. <i>caerulea</i>	elderberry
<i>Sanicula crassicaulis</i>	Pacific sanicle
<i>Scirpus microcarpus</i>	panicked bulrush
<i>Scoliopus bigelovii</i>	California fetid adder's tongue
<i>Scrophularia californica</i>	California figwort
* <i>Senecio glomeratus</i>	fireweed
<i>Senecio minimus</i>	toothed fireweed
<i>Sequoia sempervirens</i>	coast redwood
<i>Silene gallica</i>	windmill pink
* <i>Silybum marianum</i>	milk thistle
<i>Sisyrinchium bellum</i>	blue-eyed grass
<i>Solanum umbelliferum</i>	blue witch
<i>Soliva sessilis</i>	common soliva
* <i>Sonchus asper</i>	prickly sow thistle
<i>Sonchus asper</i> subsp. <i>a.</i>	prickly sow thistle
* <i>Sonchus oleraceus</i>	common sow thistle
* <i>Spergularia</i> sp.	sand spurry

<i>Stachys ajugoides</i> var. <i>rigida</i>	rigid hedge nettle
<i>Stachys bullata</i>	hedge nettle
<i>Stachys chammisonis</i>	swamp hedge-nettle
<i>Stachys rigida</i>	hedge nettle
<i>Stachys rigida</i> var. <i>quercetorum</i>	rigid hedge nettle
* <i>Stellaria media</i>	common chickweed
<i>Stipa pulchra</i>	purple needlegrass
<i>Symphoricarpos albus</i>	tall snowberry
<i>Symphoricarpos albus</i> var. <i>laevigatus</i>	snowberry
<i>Symphoricarpos mollis</i>	snowberry
<i>Symphotrichum chilense</i>	common California sister
* <i>Torilis arvensis</i>	hedge parsley
* <i>Torilis nodosa</i>	knotted hedge parsley
<i>Toxocodendron diversilobum</i>	poison oak
<i>Toxicoscordion fremontii</i>	Fremont's star lily
<i>Trientalis latifolia</i>	Pacific starflower
* <i>Trifolium angustifolium</i>	narrow-leaved clover
<i>Trifolium bididum</i> var. <i>decipiens</i>	pinole clover
<i>Trifolium dubium</i>	shamrock
<i>Trifolium gracilentum</i>	pin-point clover
* <i>Trifolium hirtum</i>	rose clover
<i>Trifolium microcephalum</i>	small-headed clover
<i>Trifolium obtusiflorum</i>	creek clover
<i>Trifolium repens</i>	white clover
<i>Trifolium variegatum</i> var. <i>v.</i>	white-tipped clover
<i>Trifolium wildenovii</i>	tomcat clover
<i>Trillium ovatum</i>	coast trillium
<i>Triodanis biflora</i>	Venus's looking glass
<i>Trisetum canescens</i>	tall trisetum
<i>Trisetum</i> sp.	trisetum
<i>Triteleia laxa</i>	Ithuriel's spear
<i>Typha latifolia</i>	broad-leaved cattail
<i>Umbellularia californica</i>	California bay-laurel
<i>Urtica dioica</i> ssp. <i>holosericea</i>	stinging nettle
<i>Urtica urens</i>	dwarf nettle
<i>Vaccinium ovatum</i>	huckleberry
<i>Vancouveria planipetala</i>	inside-out flower
<i>Verbena lasiostachys</i>	western verbena
<i>Verbena lasiostachys</i> var. <i>lasiostachys</i>	verbena
<i>Veronica americana</i>	American speedwell
<i>Veronica arvensis</i>	common speedwell
<i>Veronica peregrina</i>	purslane speedwell
<i>Veronica persica</i>	Persian speedwell
<i>Vicia disperma</i>	two-seeded vetch
* <i>Vicia sativa</i>	common vetch
<i>Vicia sativa</i> subsp. <i>nigra</i>	narrow-leaved vetch

<i>Vicia</i> sp.	wild cucumber
* <i>Vinca major</i>	periwinkle
<i>Viola ocellata</i>	two-eyed violet
<i>Viola pedunculata</i>	Johnny jump-up
<i>Viola sempervirens</i>	redwood violet
<i>Whipplea modesta</i>	yerba de selva
<i>Woodwardia fimbriata</i>	western chain fern

FUNGI OF SDSF

From personal communication with Nathan Wilson of the Fungus Federation of Santa Cruz.

<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>
<i>Agricus hondensis</i>	felt-ringed <i>Agricus</i>
<i>Amanita gemata</i>	gemmed <i>Amanita</i>
<i>Amanita pantherina</i>	panther <i>Amanita</i>
<i>Amanita phalloides</i>	death cap
<i>Amanita vaginata</i>	grisette
<i>Armillaria mellea</i>	honey mushroom
<i>Auriscalpium vulgare</i>	
<i>Boletus edulis</i>	king bolete
<i>Calodera viscosa</i>	yellow tuning fork
<i>Camarophyllus borealis</i>	snowy waxy cap
<i>Camarophyllus russocoriaceus</i>	cedar waxy cap
<i>Cantherellus californicus</i>	chanterelle
<i>Cantharellus cibarius</i>	
<i>Caulorhiza umbonata</i>	redwood rooter
<i>Clitocybe deceptiva</i>	anise mushroom
<i>Clitocybe nebularis</i>	cloudy <i>Clitocybe</i>
<i>Coprinus micaceus</i>	mica cap
<i>Cortinarius</i> sp.	<i>Cortinarius</i>
<i>Cortinarius varius</i>	<i>Cortinarius</i>
<i>Craterellus cornucopioides</i>	horn of plenty
<i>Crucibulum laeve</i>	white-egg bird's nest
<i>Entoloma</i> sp.	
<i>Fistulina hepatica</i>	beefsteak polypore
<i>Fomitopsis cajanderi</i>	
<i>Fomitopsis pinicola</i>	red-belted conk
<i>Galerina</i> sp.	
<i>Gomphidius subrosea</i>	rosy <i>Gomphidius</i>
<i>Gymnopilus sapineus</i>	common <i>Gymnopilus</i>
<i>Gyromitra infula</i>	hooded false morel
<i>Hebeloma crustuliniforme</i>	poison pie
<i>Helvella compressa</i>	
<i>Helvella lacunosa</i>	fluted black elfin saddle

<i>Helvella maculata</i>	
<i>Hemimycena</i> sp.	
<i>Heterotextus alpinus</i>	
<i>Hygrocybe acutoconica</i>	a cute conic waxy cap
<i>Hygrocybe coccinea</i>	righteous red waxy cap
<i>Hygrocybe conica</i>	witch's hat
<i>Hygrocybe flavescens</i>	golden waxy cap
<i>Hygrocybe punicea</i>	scarlet waxy cap
<i>Inocybe citrifolia</i>	
<i>Inocybe fastigiata</i>	corn silk Inocybe
<i>Lactarius argillaceifolius</i>	vulgar milk cap
<i>Lactarius chrysorheus</i>	yellow-staining milk cap
<i>Lactarius fragilis</i>	candy cap
<i>Lactarius rubrilacteus</i>	bleeding milk cap
<i>Lenzites betulina</i>	gilled polypore
<i>Leotia lubrica</i>	
<i>Leptonia parva</i>	blue-black Leptonia
<i>Naematoloma fasciculare</i>	sulfur tuft
<i>Paxillus involutus</i>	poison pax
<i>Phylloporus rhodoxanthus</i>	gilled bolete
<i>Pleurocybella porrigens</i>	angel wings
<i>Pleurotus ostreatus</i>	oyster mushroom
<i>Pluteus cervinus</i>	deer mushroom
<i>Psathyrella longipes</i>	
<i>Pseudohydnum gelatinosum</i>	jelly tooth
<i>Ramaria</i> sp.	Ramaria
<i>Ramaria gelatinosa</i>	jellied-base coral
<i>Ramaria stricta</i>	straight-branched coral
<i>Russula brevipes</i>	short-stemmed Russula
<i>Russula cremoricolor</i>	creamy Russula
<i>Russula silvicola</i>	emetic Russula
<i>Stereum hirsutum</i>	hairy Stereum
<i>Trametes versicolor</i>	turkey tail
<i>Verpa conica</i>	thimble morel

APPENDIX C: SOQUEL DEMONSTRATION STATE FOREST MONITORING PLAN

Identified repair projects will be implemented as staffing and budget limitations allow. To finance their construction, these projects will be generally incorporated into the timber harvest planning process.

WATERSHED MONITORING

TIMBER OPERATIONS

To avoid adverse impacts on water quality and fisheries resulting from the discharge of sediment to watercourses attributable to timber operations, SDSF staff will monitor all timber operations (including all harvesting areas and new roads, skid trails, and landings) annually for five to seven years following completion of the operations. Occurrences of substantial surface erosion (i.e., gullies) or mass wasting (i.e., landslides or slumps) resulting from the operations will be identified and described by a Registered Professional Forester (RPF).

Each substantial gully or landslide will be evaluated to determine its cause and identify stabilization measures that would be most feasible, effective, and cost effective. Such measures will be implemented within 90 days from the date when the subject site is identified, unless due cause for delay is explained and a reasonable alternative schedule for implementation is proposed by the SDSF Forest Manager. If, based on the judgment of a Certified Engineering Geologist (CEG), no stabilization measures are feasible or reasonable to apply to the subject site, feasible offsite watershed remediation measures will be implemented as recommended by the CEG in conjunction with the next timber operation conducted at SDSF.

The SDSF Forest Manager will describe appropriate mitigation measures to be incorporated into future timber operations and specified in future timber harvesting plans (THPs) to avoid a recurrence of the observed erosion or mass wasting events.

ROADS

An inventory that delineated, described, and risk-rated forest roads at SDSF was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (Pacific Watershed Associates, 2003). This assessment included identification, prioritization, and design of feasible projects to remediate sediment-discharge risks associated with each high- and extreme-risk road segment and estimated the cost effectiveness (in dollars per cubic yard of sediment) of each such project. Such projects include recontouring of roads and/or installation of improved drainage structures.

SDSF staff will conduct forensic monitoring of roads during and/or following most large storms. Any repair needs will be addressed as quickly as possible within the constraints of funding, and legal and regulatory requirements.

CULVERTS

SDSF staff evaluated the culverts along all Forest roads to determine where trash racks were necessary and appropriate. Steel T-posts were installed above each culvert where a need was determined. This work was completed prior to December 31, 1998. Each culvert is inspected annually and after large storm events (defined as producing rainfall of about 2 inches in 24 hours), and cleaned as needed. An inventory to delineate, describe, and risk-rate culverts at SDSF was completed as part of the Santa Cruz County Resource Conservation District's Soquel Creek watershed assessment (Pacific Watershed Associates, 2003). This information was compiled into a database to be maintained on an ongoing basis. The risk rating was based on standard engineering criteria for adequate culvert sizing, including effective culvert diameter, extent of watershed area, and estimated discharge of a storm with a 100-year recurrence interval.

The inventory included identification, prioritization, and design of feasible projects to reduce the risk of failure of high- or extreme-risk culverts, and estimated the cost-effectiveness (in dollars per cubic yard of sediment) of each such project. Potential projects include replacement of existing culverts with more reliable drainage structures (e.g., rocked dips) or larger culverts.

SDSF staff will conduct forensic monitoring of drainage structures during and/or following most large storms. Any repairs needs will be addressed as quickly as possible within the constraints of funding and legal and regulatory requirements.

ACTIVE LANDSLIDES AND SEDIMENT STORES

CAL FIRE, in conjunction with a CEG, will continue to inventory, delineate, describe and risk-rate active landslides and substantial unconsolidated concentrations of sediment and debris at SDSF. This information will be compiled into a database to be maintained on an ongoing basis, with new landslides added as they occur. Unconsolidated concentrations of sediment and debris and active landslides identified by Manson and Sowma-Bawcom (1992) will be included in the database. The description and risk-rating of segments will focus on conditions affecting the likelihood that runoff or seismic activity will cause mass movement resulting in sediment discharge to a watercourse.

CAL FIRE, in conjunction with a CEG, will continue to identify and design feasible projects to reduce the risk of failure of high- or extreme-risk landslides and sediment/debris accumulations, and estimate the cost-effectiveness (in dollars per cubic yard of sediment) of each such project. Landslide risk-reduction projects could include installation of drainage structures on roads above landslides to reduce the discharge of concentrated runoff onto the landslide, or armoring or buttressing of landslide toes. Sediment/debris accumulation risk-reduction projects could include recontouring or removal of material to reestablish original watercourses, bucking of logs in debris piles, and using the logs to armor eroding bank faces, or revegetation.

In 2004, the California Geological Survey submitted an Engineering Geologic Report for the portion of Fern Gulch that was included in the Fern Gulch timber sale. This report includes a detailed landslide map that assisted with timber sale development.

AMAYA BASIN

The California Geological Survey compiled a map of landslides triggered by the 1989 Loma Prieta earthquake for Soquel Demonstration State Forest (Bedrossian, 1989). This map was focused on earthquake-triggered landslides and does not identify other unstable areas. Additional reconnaissance-level slope stability mapping, which includes descriptions of the geology and geomorphologic features related to slope stability was prepared by Manson and Sowma-Bawcom (1992), including all of the Amaya Creek and Fern Gulch watersheds. This mapping was based upon aerial photographic interpretations with little field work and is recognized as being useful only for preliminary review of regional slope stability.

Additional review and mapping of the Amaya Basin was completed in 2001 by Wayne Haydon of the California Geological Survey. Two maps were produced titled *Relative Landslide Potential, Amaya and Fern Gulches* and *Geologic and Geomorphic Features Related to Landsliding, Amaya and Fern Gulches*. These maps were used as a basis for the preparation of the Fern Gulch THP as well as the more recent Comstock Mill THP in the Amaya basin.

REMEDIATION PROJECTS

CAL FIRE will prioritize all sediment discharge projects identified and designed to remediate high- or extreme-risk conditions, as described above. Such projects will be implemented in conjunction with future timber operations according to their priority as available funding permits, to ensure, to the extent practicable, that no significant increase in sedimentation results from the timber operations proposed in each THP.

SDSF staff will continue to monitor roads and drainage structures following large storm events. When possible, inspections will be conducted during storm events. Any repair needs will be addressed as quickly as possible. To avoid adverse impacts of implementing remediation projects on water quality and fish habitat, CAL FIRE will evaluate the performance of each previously-implemented remediation project to determine whether the project was successful in reducing risk of large-scale sedimentation. If CAL FIRE determines that any project is not meeting its intended objective, CAL FIRE will redesign and modify the project as needed.

AQUATIC RESOURCE MONITORING

FISHERIES MANAGEMENT PLAN

A Draft Fisheries Management Plan for SDSF was completed in August, 1995 (Berlekamp and Sutfin). The aquatic-resource monitoring program includes the items described below.

FISH POPULATIONS

In cooperation with the California Department of Fish and Wildlife (CDFW), fish population surveys were conducted at four separate sites on SDSF from 1993 to 2001. From 2002 to the present, SDSF has continued these annual surveys in cooperation with the National Oceanic and Atmospheric Administration's National Marine Fisheries Service (NOAA Fisheries). NOAA

Fisheries added a fifth permanent station, increased the monitoring frequency, and incorporated additional research components.

HABITAT

One formal and comprehensive aquatic habitat survey was conducted on SDSF in 1994 (Berlekamp). Several additional surveys have been conducted by various fisheries biologists in conjunction with other research studies.

POOL SEDIMENTATION AND STREAM TEMPERATURE

Aquatic resources monitoring includes a pool sedimentation component to assess and monitor pool substrate embeddedness. When the Draft Fisheries Management Plan was completed, the V* methodology was a popular assessment procedure to quantitatively measure sediment impairment. Fisheries biologists and hydrologists later advised SDSF staff that measuring sediment in a small number of pools, particularly using the V* method, would not be as useful as measuring other parameters, such as water temperature. To date, no formal survey has been conducted to assess pool sedimentation on watercourses within the Forest. The 1994 Fish Habitat Survey Report and other limited-scope investigations provide descriptions of the channel substrate composition.

On the advice of fisheries biologists, SDSF staff initiated a long-term program to monitor stream temperature at seven sites on the Forest. Continuous water temperature data has been collected from 1997 to the present (CAL FIRE, 1997-2012).

AQUATIC INVERTEBRATES

A macroinvertebrate sampling study was performed by SDSF staff in October, 1995. The California Department of Fish and Wildlife analyzed the samples and produced a report of their findings in May, 1996. Since then, NOAA Fisheries has conducted additional assessments in conjunction with other research projects. Additional funding and agency coordination will be investigated for future aquatic invertebrate studies.

HABITAT ENHANCEMENT STRUCTURES

Fish habitat enhancement structures will be reviewed and approved by CDFW and an inter-agency team before installation.

To avoid adverse impacts of habitat-enhancement structures on fish habitat, CAL FIRE, in conjunction with CDFW, will evaluate structures annually for three years following installation to determine whether they are performing as intended and whether they are causing any unintended adverse impacts on fish habitat. If CAL FIRE and CDFW determine that any structure is functioning improperly or is adversely affecting aquatic habitat, CAL FIRE will redesign, repair, or remove the structure, as needed.

FISH POACHING AND HARASSMENT

To avoid adverse impacts of public use on fish, CAL FIRE and CDFW will conduct ongoing patrols in SDSF to enforce prohibitions on fishing and fish harassment. Incidents of fishing or harassment will be recorded and compiled. CAL FIRE and CDFW will evaluate such information annually in conjunction with fish population estimates to determine whether poaching or harassment have had a significant adverse effect on SDSF fisheries. If so, CAL FIRE will respond by closing streamside trails and intensifying law enforcement (e.g., increased patrols of streamside roads) as needed to ensure rapid fishery recovery and avoid additional adverse fishery effects.

RIPARIAN RESOURCE MONITORING

To avoid adverse impacts of public use on riparian habitat, CAL FIRE will conduct ongoing patrols in SDSF to enforce prohibitions on vandalism and other damage to riparian habitat related to public use. Incidents of damage will be recorded and compiled. A qualified wildlife biologist will evaluate such information annually to determine whether the damage constitutes a significant adverse effect on wildlife. If so, CAL FIRE will implement additional restrictions on public use (e.g., prohibiting camping or weekday recreation use or cordoning off sensitive areas) as needed to ensure rapid habitat recovery and to avoid additional adverse wildlife effects.

WILDFIRE MONITORING

To minimize increases in wildfire risks resulting from increased public use at SDSF, CAL FIRE will record and compile descriptions of all wildfires occurring at SDSF, including ambient weather and fire hazard conditions, ignition source, area and vegetation types burned, and estimated damage. Such information will be evaluated annually to determine frequency and severity of wildfire incidents. CAL FIRE will implement appropriate measures (e.g., prohibiting all fires or weekday recreation use, or allocating additional fire-suppression resources for SDSF) as needed to reduce wildfire risks.

EMERGENCY SERVICES MONITORING

To minimize adverse effects on emergency response performance for residents of Santa Cruz County resulting from increased demands for such services at SDSF, CAL FIRE will record and compile all requests for emergency responses, including requests for police, fire, medical, or search and rescue services. Descriptions of emergency responses will include response times and will be evaluated annually. Improvements to emergency response will be evaluated and, as resources allow, CAL FIRE will enhance its emergency response capability by upgrading roads; developing additional helispots along remote trails; and, either directly or through a management agreement with another qualified entity, providing additional human and equipment resources for emergency response at SDSF.

This information has not been compiled into a comprehensive report, but the data is examined by staff to evaluate any trends. All incident reports must be requested through the Fire Prevention/Law Enforcement Bureau of CAL FIRE to insure that legally confidential information

(such as ongoing law enforcement actions, and the identities of juveniles or injured Forest visitors) is not disclosed.

NUISANCE MONITORING

To minimize nuisance impacts on SDSF neighbors, CAL FIRE will record and compile descriptions of all reported nuisances caused by SDSF users at SDSF or on adjacent ownerships including, but not limited to, trespass, vandalism, littering, and noise. This information has not been compiled into a comprehensive report, but the data is examined by staff to evaluate any trends.

SDSF staff work closely with individuals and groups of volunteers to assist in this effort. The Stewards of Soquel Forest and a local National Mountain Bike Patrol group have both contributed information to the nuisance monitoring database. SDSF staff have developed good relationships with neighbors to prevent incidents that constitute a nuisance.

PUBLIC-USE MONITORING

CAL FIRE will use camping records, surveys and other information to compile annual estimates of public use of SDSF in user days. Use will be estimated for specific activities including, but not limited to, mountain biking, hiking, equestrian use, and educational activities.

CULTURAL RESOURCE MONITORING

CAL FIRE will monitor and periodically inspect heritage resources on SDSF to ensure that existing policies are affording effective protection. The identification and protection of cultural resources are important components of forestry in California today. Registered Professional Foresters are required to attend archaeological training classes to acquire the ability to recognize cultural materials, and to be able to develop effective protection measures. In its role as a demonstration forest, SDSF can serve as a proving ground for the development and implementation of effective heritage resource management strategies.

APPENDIX D: BEST MANAGEMENT PRACTICES

INVASIVE PLANTS

- Scout the work site and access roads for presence of invasive plants prior to starting project work. Flag the location(s) of invasive plants to be avoided. If avoidance is not possible, treat or remove any invasive plants prior to allowing equipment to enter the work area.
- Schedule activities for periods when the risk of introducing or spreading invasive plants will be low (such as prior to flowering or when plant is dormant).
- Include methods to reduce the spread of invasive plants in training session prior to starting work on project.
- Clean clothing, tools, vehicles, equipment, and gear prior to entering the project site. Require the use of an inspection form to document that cleaning occurred before leaving an infested area or arriving at non-infested sites.
- Use a weed-free source for materials (including sand, gravel, seed, mulch, and other erosion control materials) brought in from outside locations. If no weed-free source is available, inspect, sterilize and/or treat material as needed.
- Prevent contamination of materials by invasive plants during storage and transport. Cover stockpiles of top soil, mulch, etc. with impermeable material (such as a tarp).
- Revegetate and/or mulch disturbed soils as soon as possible to reduce the likelihood of invasive plant establishment.
- Include not introducing invasive plants as a performance requirement in contracts.
- Minimize soil disturbance when conducting fuels reduction projects by using techniques such as chainsaw thinning and piling slash by hand.

AQUATIC ORGANISMS

- Disinfect all research equipment brought in from other watersheds. Include wading, diving, survey, electrofishing, and bug collecting gear, as well as in-stream equipment including ropes, fencing, and fish traps. Begin the decontamination procedure by scrubbing off equipment so it is clean and free of mud and other debris.
- Equipment can then be placed in a freezer overnight (at least four hours); soaked in hot water (at least five minutes in water at least 120 degrees); or soaked for at least 20 minutes in a 10% solution of bleach. If using diluted bleach, allow equipment to drip dry before rinsing. (Two other disinfectants, Roccal and iodophore, are acceptable substitutes for diluted bleach.)
- Vehicles carrying gear from areas infested with New Zealand mudsnail should also be cleaned to remove dirt and debris that might contain the snails.

SUDDEN OAK DEATH (SOD)

- Minimize travel, equipment operation, and/or collection of plants, firewood, or water in SOD disease centers.
- Minimize working in disease centers during muddy conditions, and during periods that are warm and rainy.
- If possible, start work in healthy stands before entering infected areas. Clean cutting tools (loppers, chainsaws, etc.) used on infected trees before working on healthy ones.
- Clean tools, equipment, tires, and foot wear after working in infected areas. Brush off mud, leaves and other debris, and spray shoes and any cutting tools with a disinfectant, such as Lysol or a 10% bleach solution. Power washers or car washes are effective means for cleaning cars, trucks, or large pieces of equipment.
- Movement of soil or plant parts (including firewood) may require a permit from the County Agricultural Commissioner. See <http://www.suddenoakdeath.org/diagnosis-and-management/regulations/> for detailed information on requirements.

HAZARDOUS MATERIALS

- Material Safety Data Sheets (MSDS), material labels and any additional handling and emergency instruction of hazardous materials are kept on file at the SDSF Headquarters.
- Any state employee handling hazardous materials are made aware of the potential hazards, given proper training and instruction, and also made aware of the location of the MSDS and any other documentation for the material.
- All contractors used in the application or use of these hazardous materials shall have the appropriate licenses and be able to read and understand the MSDS labels, appropriate recommendations and application instructions.
- The storage of the potentially hazardous materials on SDSF is in accordance with the MSDS and any buildings that are used for storage will display appropriate placards.